



STABILITY OF A FUNCTIONAL EQUATION COMING FROM THE CHARACTERIZATION OF THE ABSOLUTE VALUE OF ADDITIVE FUNCTIONS

ATTILA GILÁNYI^{1*}, KAORI NAGATOU² AND PETER VOLKMANN³

Communicated by T. Riedel

ABSTRACT. In the present paper, we prove the stability of the functional equation

$$\max\{f((x \circ y) \circ y), f(x)\} = f(x \circ y) + f(y)$$

for real valued functions defined on a square-symmetric groupoid with a left unit element. As a consequence, we obtain the known result about the stability of the equation

$$\max\{f(x + y), f(x - y)\} = f(x) + f(y)$$

for real valued functions defined on an abelian group.

1. INTRODUCTION

The topic of this paper is related to the study of the functional equation

$$\max\{f(x + y), f(x - y)\} = f(x) + f(y). \quad (1.1)$$

The general solution of this equation has been determined in the class of real valued functions defined on an abelian group G in a joint paper with Alice Simon [19]. It has been proved that a function f of this type satisfies (1.1) if and only if it has the form

$$f(x) = |a(x)| \quad (x \in G),$$

where $a : G \rightarrow \mathbb{R}$ is an additive function.

In the following, we will investigate the stability of a generalization of the equation above in the sense of Pólya–Szegő ([15]) and Hyers–Ulam ([20, 10]). Based on

Date: Received: 15 October 2010; Accepted: 3 December 2010

* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 39B82; Secondary 39B52, 46E99.

Key words and phrases. Stability of functional equations, square-symmetric groupoids.

the simple observation that replacing $x - y$ by \bar{x} in equation (1.1), it becomes

$$\max\{f((\bar{x} + y) + y), f(\bar{x})\} = f(\bar{x} + y) + f(y) \quad (\bar{x}, y \in G),$$

we may consider it on more general structures than groups. We will prove the stability of this equation for real valued functions defined on a square-symmetric groupoid with a left unit element. As a consequence of our main theorem, we get the stability of equation (1.1) on abelian groups, which has been proved in a joint paper with Witold Jarczyk [8].

2. NOTATION AND TERMINOLOGY

Throughout the paper, (S, \circ) denotes a *groupoid*, that is, a nonempty set S with a binary operation $\circ : S \times S \rightarrow S$. If $x \in S$, we define $x^1 = x$ and, for a nonnegative integer n , $x^{2^{n+1}} = x^{2^n} \circ x^{2^n}$.

A groupoid (S, \circ) (or the operation \circ on S) is called *square-symmetric* if

$$(x \circ y) \circ (x \circ y) = (x \circ x) \circ (y \circ y)$$

holds for all $x, y \in S$.

Obviously, a commutative and associative binary operation is also square-symmetric. However, neither commutativity nor associativity follow from square-symmetry. For example, the operation $x \circ y = y$ is associative, square-symmetric but not commutative on an arbitrary set S with at least two elements (concerning the connection between commutativity and square-symmetry, we refer to [9]). Furthermore, the commutative operation $x \circ y = |x - y|$ on $S = [0, \infty[$ is also square-symmetric but not associative. Moreover, power-symmetry is a “weaker property” than bisymmetry: the commutative and square-symmetric operation $x \circ y = |x - y|$ above is not bisymmetric. (An operation on S is called bisymmetric if $(x \circ \bar{x}) \circ (y \circ \bar{y}) = (x \circ y) \circ (\bar{x} \circ \bar{y})$ for $x, \bar{x}, y, \bar{y} \in S$; cf. [1]).

The stability of functional equations on square-symmetric groupoids was investigated by several authors. Among others, Karol Baron, Gian Luigi Forti, Zoltán Kaiser, R. Duncan Luce, Zenon Moszner, Zsolt Páles and Jürg Rätz proved such type of results in [2, 3, 4, 5, 6, 7, 11, 12, 13, 14, 18, 21, 22, 23]. Let us mention recent results of Barbara Przebieracz ([16, 17]) on the stability of some equations connected with the topic of the present paper (cf. also the references in [8]).

3. STABILITY THEOREMS

Theorem 3.1. *Let (S, \circ) be a square-symmetric groupoid and let δ be a nonnegative real number. If a function $f : S \rightarrow \mathbb{R}$ satisfies the inequality*

$$|f(x \circ x) - 2f(x)| \leq \delta \quad (x \in S)$$

then the function $g : S \rightarrow \mathbb{R}$ defined by

$$g(x) = \lim_{n \rightarrow \infty} \frac{1}{2^n} f(x^{2^n}) \quad (x \in S), \quad (3.1)$$

solves the equation

$$g(x \circ x) = 2g(x) \quad (x \in S) \quad (3.2)$$

and fulfils

$$|f(x) - g(x)| \leq \delta \quad (x \in S). \quad (3.3)$$

Moreover, there exists only one solution $g : S \rightarrow \mathbb{R}$ of (3.2) for which the left hand side in (3.3) is bounded.

Proof. The statement is a simple consequence of the results presented in [23] and [7]. \square

Theorem 3.2. *Let (S, \circ) be a square-symmetric groupoid with a left unit element and let $f : S \rightarrow \mathbb{R}$ be a function. If there exists a nonnegative real number ε such that*

$$|\max\{f((x \circ y) \circ y), f(x)\} - f(x \circ y) - f(y)| \leq \varepsilon \quad (x, y \in S) \quad (3.4)$$

then there exists a solution $g : S \rightarrow \mathbb{R}$ of the functional equation

$$\max\{g((x \circ y) \circ y), g(x)\} = g(x \circ y) + g(y) \quad (x, y \in S) \quad (3.5)$$

for which

$$|f(x) - g(x)| \leq 3\varepsilon \quad (x \in S). \quad (3.6)$$

Moreover, there exists only one solution $g : S \rightarrow \mathbb{R}$ of (3.5) for which the left hand side in (3.6) is bounded.

Proof. Replacing x and y by the unit element e of S in (3.4), we get that

$$|f(e)| \leq \varepsilon. \quad (3.7)$$

Writing e instead of x in (3.4), we obtain

$$|\max\{f(y \circ y), f(e)\} - 2f(y)| \leq \varepsilon \quad (y \in S). \quad (3.8)$$

Inequality (3.7) gives $f(e) \geq -\varepsilon$, thus, $\max\{f(y \circ y), f(e)\} \geq -\varepsilon$, which together with (3.8) implies

$$f(y) \geq -\varepsilon \quad (y \in S). \quad (3.9)$$

A simple calculation based on inequalities (3.8) and (3.9) yields

$$|f(y \circ y) - 2f(y)| \leq 3\varepsilon \quad (y \in S). \quad (3.10)$$

In fact, if $\max\{f(y \circ y), f(e)\} = f(y \circ y)$ for a $y \in S$ then $|f(y \circ y) - 2f(y)| \leq \varepsilon$, thus, (3.10) follows from (3.8). On the other hand, if $\max\{f(y \circ y), f(e)\} = f(e)$ for a $y \in S$ then, by (3.7), we have $f(y \circ y) \leq \varepsilon$, (3.8) gives $f(y) \leq \varepsilon$, (3.9) implies $f(y) \geq -\varepsilon$ and $f(y \circ y) \geq -\varepsilon$, and these properties yield (3.10).

Based on inequality (3.10) and applying Theorem 3.1 with $\delta = 3\varepsilon$, we obtain that the function $g : S \rightarrow \mathbb{R}$ defined by (3.1) satisfies properties (3.2) and (3.6). Writing x^{2^n} instead of x and y^{2^n} instead of y in (3.4), we obtain

$$|\max\{f((x^{2^n} \circ y^{2^n}) \circ y^{2^n}), f(x^{2^n})\} - f(x^{2^n} \circ y^{2^n}) - f(y^{2^n})| \leq \varepsilon \quad (3.11)$$

for all $x, y \in S$ and $n \in \mathbb{N} \cup \{0\}$, where $\mathbb{N} = \{1, 2, 3, \dots\}$. It can be shown by induction that the square-symmetry of S implies

$$x^{2^n} \circ y^{2^n} = (x \circ y)^{2^n} \quad (x, y \in S, n \in \mathbb{N} \cup \{0\}),$$

which yields

$$(x^{2^n} \circ y^{2^n}) \circ y^{2^n} = ((x \circ y) \circ y)^{2^n} \quad (x, y \in S, n \in \mathbb{N} \cup \{0\}).$$

Thus, dividing inequality (3.11) by 2^n and letting n approach infinity, we obtain that the function g satisfies (3.5).

In order to prove uniqueness, suppose that $\bar{g} : S \rightarrow \mathbb{R}$ is a solution of (3.5) which fulfils

$$|f(x) - \bar{g}(x)| \leq K \quad (x \in S)$$

with a nonnegative constant $K \in \mathbb{R}$. The triangle inequality, with (3.6) and the previous inequality, implies

$$|g(x) - \bar{g}(x)| \leq K + 3\varepsilon \quad (x \in S).$$

Since g and \bar{g} are solutions of inequality (3.4) with $\varepsilon = 0$, inequality (3.10) gives

$$g(x \circ x) = 2g(x) \quad (x \in S)$$

and

$$\bar{g}(x \circ x) = 2\bar{g}(x) \quad (x \in S).$$

Based on the last three formulas, Theorem 3.1 yields $g(x) = \bar{g}(x)$, $(x \in S)$, that is, the uniqueness part of our theorem. \square

As a corollary of Theorem 3.2, we obtain the stability result of [8].

Theorem 3.3. *Let $(G, +)$ be an abelian group and $f : G \rightarrow \mathbb{R}$ be a function. If there exists a nonnegative real number ε such that*

$$|\max\{f(x+y), f(x-y)\} - f(x) - f(y)| \leq \varepsilon \quad (x, y \in G) \quad (3.12)$$

then there exists a solution $g : G \rightarrow \mathbb{R}$ of the functional equation

$$\max\{g(x+y), g(x-y)\} = g(x) + g(y) \quad (x, y \in G) \quad (3.13)$$

for which

$$|f(x) - g(x)| \leq 3\varepsilon \quad (x \in G). \quad (3.14)$$

Moreover, there exists only one solution $g : G \rightarrow \mathbb{R}$ of (3.13) for which the left hand side in (3.14) is bounded.

Proof. Substituting $x - y$ by \bar{x} in inequality (3.12), we obtain that

$$|\max\{f((\bar{x} + y) + y), f(\bar{x})\} - f(\bar{x} + y) - f(y)| \leq \varepsilon \quad (\bar{x}, y \in G).$$

Since the abelian group G is a square-symmetric groupoid with a left unit element, Theorem 3.2 implies the existence of a function $g : G \rightarrow \mathbb{R}$ which solves the equation

$$\max\{g((\bar{x} + y) + y), g(\bar{x})\} = g(\bar{x} + y) + g(y) \quad (\bar{x}, y \in G)$$

and satisfies inequality (3.14). Writing $x = \bar{x} + y$ in the last equation, we get (3.13), which yields the existence part of our statement. The uniqueness can be proved based on Theorem 3.2 using the substitution $\bar{x} = x - y$ again. \square

Remark 3.4. According to the argumentation in the proof above, equations (3.5) and (3.13) are equivalent on abelian groups (i.e., they have the same solutions there). However, as it is well-known (cf., e.g., [24]), the equivalence of two functional equations does not imply that they are stable or unstable at the same time. This is the reason why we formulated our statement concerning the stability of equation (3.13) in a separate theorem.

Finally, we note that, in the class of real valued functions defined on a square-symmetric groupoid with a left unit element, the general solution of equation (3.5) has not been determined yet.

Acknowledgement. Some parts of the results presented here were achieved during a stay of the first author at Kyushu University, Fukuoka, Japan in 2008. He is thankful for the invitation and the kind hospitality of his host during his visit.

The research of the first author has been supported by the Hungarian Scientific Research Fund (OTKA) Grant NK-81402 and by TÁMOP 4.2.1./B-09/1/KONV-2010-0007/IK/IT project. The project is implemented through the New Hungary Development Plan co-financed by the European Social Fund, and the European Regional Development Fund. This work was partially supported by PRESTO, Japan Science and Technology Agency and a Grant-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan (No. 20224001).

REFERENCES

1. J. Aczél, *Vorlesungen über Funktionalgleichungen und ihre Anwendungen*, Birkhäuser Verlag, Basel, 1961.
2. K. Baron, P. Volkman, *On functions close to homomorphisms between square symmetric structures*, Sem. LV, no. 14 (2002), 12pp., <http://www.math.us.edu.pl/smdk>
3. G.L. Forti, *An existence and stability theorem for a class of functional equations*, Stochastica **4** (1980), 23–30.
4. G.L. Forti, *Continuous increasing weakly bisymmetric groupoids and quasi-groups in \mathbb{R}* , Math. Pannonica **8** (1997), 49–71.
5. G.L. Forti, *Comments on the core of the direct method for proving Hyers–Ulam stability of functional equations*, J. Math. Anal. Appl. **295** (2004), 127–133.
6. A. Gilányi, *Hyers–Ulam stability of monomial functional equations on a general domain*, Proc. Natl. Acad. Sci. USA **96** (1999), 10588–10590.
7. A. Gilányi, Z. Kaiser, Zs. Páles, *Estimates to the stability of functional equations*, Aequationes Math. **73** (2007), 125–143.
8. W. Jarczyk, P. Volkman, *On functional equations in connection with the absolute value of additive functions*, Ser. Math. Catoviciensis et Debreceniensis, no. 32 (2010), 11 pp., <http://www.math.us.edu.pl/smdk>
9. E.C. Johnson, D.L. Outcalt, A. Yagub, *An elementary commutativity theorem for rings*, Amer. Math. Monthly **75** (1968), 288–289.
10. D.H. Hyers, *On the stability of the linear functional equation*, Proc. Natl. Acad. Sci. USA **27** (1941), 222–224.
11. Z. Moszner, *On the stability of functional equations*, Aequationes Math. **77** (2009), 33–88.
12. Zs. Páles, *Hyers–Ulam stability of the Cauchy functional equation on square-symmetric groupoids*, Publ. Math. Debrecen **58** (2001), 651–666.
13. Zs. Páles, *Hyers–Ulam stability of generalized monomial functional equations on a general domain*, manuscript.
14. Zs. Páles, P. Volkman, R. D. Luce, *Hyers–Ulam stability of functional equations with a square-symmetric operation*, Proc. Natl. Acad. Sci. USA **95** (1998), 12772–12775.
15. Gy. Pólya, G. Szegő, *Aufgaben und Lehrsätze aus der Analysis, Vol. I*, Springer, Berlin, 1925.
16. B. Przebieracz, *Superstability of some functional equation*, Ser. Math. Catoviciensis et Debreceniensis, no. 31 (2010), 4pp., <http://www.math.us.edu.pl/smdk>
17. B. Przebieracz, *Stability of the Baron–Volkman functional equations*, Math. Inequal. Appl. (to appear).

18. J. Rätz, *On approximately additive mappings*, General Inequalities 2 (Ed. W. Walter), Birkhäuser, Basel, 1980, 233–251.
19. A. Simon (Chaljub-Simon), P. Volkman, *Caractérisation du module d'une fonction additive à l'aide d'une équation fonctionnelle*, Aequationes Math. **47** (1994), 60–68.
20. S. Ulam, *A Collection of Mathematical Problems*, Interscience Tracts in Pure and Applied Mathematics, no. 8, Interscience Publishers, New York, London, 1960.
21. P. Volkman, *On the stability of the Cauchy equation*, Proceedings of the International Conference Numbers, Functions, Equations '98 (Ed. Zs. Páles), Leaflets in Mathematics, Janus Pannonius University, Pécs, 1998, 150–151.
22. P. Volkman, *Zur Rolle der ideal konvexen Mengen bei der Stabilität der Cauchyschen Funktionalgleichung*, Sem. LV, no. 6 (1999), 6pp., <http://www.math.us.edu.pl/smdk>
23. P. Volkman, *O stabilności równań funkcyjnych o jednej zmiennej*, Sem. LV, no. 11 (2001), 6pp., <http://www.math.us.edu.pl/smdk>
24. P. Volkman, *Instabilität einer zu $f(x+y) = f(x) + f(y)$ äquivalenten Funktionalgleichung*, Sem. LV, no. 23 (2006), 1p., <http://www.math.us.edu.pl/smdk>

¹ FACULTY OF INFORMATICS, UNIVERSITY OF DEBRECEN, PF. 12, 4010 DEBRECEN, HUNGARY.

E-mail address: gilanyi@math.klte.hu

² FACULTY OF MATHEMATICS, KYUSHU UNIVERSITY, 744 MOTOOKA, NISHI-KU, FUKUOKA, 819-0395, JAPAN;

PRESTO, JAPAN SCIENCE AND TECHNOLOGY AGENCY

E-mail address: nagatou@math.kyushu-u.ac.jp

³ INSTITUT FÜR ANALYSIS, KIT, 76128 KARLSRUHE, GERMANY;

INSTYTUT MATEMATYKI, UNIWERSYTET ŚLĄSKI, BANKOWA 14, 40-007 KATOWICE, POLAND.