Acta Math. Univ. Comenianae
Vol. LXXI, 2(2002), pp. 221–230

POINTWISE WEIGHTED VECTOR ERGODIC THEOREM IN $L^1(X)$

K. EL BERDAN

Abstract. In this paper we prove the almost everywhere convergence of weighted multiparameter averages of linear surjective isometries in $L^1(X)$ and power bounded in $L^p(X), 1 < p \leq \infty$.

Throughout this paper X will be denoted a Banach space with norm $\|.\|$ and (Ω, β, μ) a σ-finite measure space. For $1 \leq p < \infty$, $L^p(X) = L^p(\Omega, X) = L^p((\Omega, \beta, \mu), X)$ denoted the usual Banach space of $X-$valued strongly measurable functions f on Ω with the norm given by

$$\begin{aligned} \|f\|_p &= \left(\int |f|^p \, d\mu\right)^{\frac{1}{p}} < \infty \text{ if } 1 \leq p < \infty, \\ \|f\|_\infty &= ess \sup \{|f(\omega)| \, ; \omega \in \Omega\} < \infty \text{ if } p = \infty. \end{aligned}$$

Let $d \geq 1$ be an integer, and let $T_1, \ldots, T_d$ be linear surjective isometries on $L^1(\Omega, X)$ such that each T_i is power bounded in $L^\infty(\Omega, X)$. Thus T_i, $1 \leq i \leq d$, can be considered to be power bounded in $L^p(\Omega, X)$ for each $1 < p < \infty$, by the Riesz convexity theorem.We will be concerned with classes of weights $\left\{a(\mathbf{k}); \mathbf{k} \in \mathbf{Z}_d^+\right\}$ such that the limit of averages

$$\lim_{|N| \to \infty} \frac{1}{|\mathbf{N}|} \sum_{k=0}^{N-1} a(\mathbf{k}) \mathbf{T}^{\mathbf{k}} f$$

exists a.e. for all $f \in L^1(X)$, where $\mathbf{T}^{\mathbf{k}} = T_1^{k_1} \ldots T_d^{k_d}$ with $\mathbf{N} = (N_1, \ldots, N_d)$ and $\mathbf{k} = (k_1, \ldots, k_d)$, $|\mathbf{N}| = N_1 \ldots N_d$, etc.....

The class of weights we will consider are the Besicovich sequences in $\mathbf{Z}_d^+$. In the case $d = 1$, Besicovitch sequences are defined to be the class of sequences $a(k)$ such that given $\varepsilon > 0$, there is a trigonometric polynomial ψ_ε such that

$$\limsup_n \frac{1}{n} \sum_{k=0}^{n-1} |a(k) - \psi_\varepsilon(k)| < \varepsilon$$

Received July 07, 2001.
1991 *Mathematics Subject Classification.* Primary 47A35, Secondary 28D05.
Key words and phrases. norm resolution, ergodic theorem, Banach space.
Thanks for Antoine Brunel and Sylvie Delabriere for helpful suggestions and the refree for kind advices.

and bounded Besicovitch weights are bounded weights in this class. Let us define the d-dimensional analogs of Besicovitch sequences: We say that the sequence $\{a(\mathbf{k})\}$ to be r-Besicovich if for every $\varepsilon > 0$ there is a sequence of trigonometric polynomials in d variables such that

$$\lim \sup_{N\to\infty} \frac{1}{|\mathbf{N}|} \sum_{\mathbf{k}=0}^{\mathbf{N}-1} |a(\mathbf{k}) - \psi_\varepsilon(\mathbf{k})|^r < \varepsilon.$$

We denote this class by $B(r)$. We say that $\{a(\mathbf{k})\}$ to be r-bounded Besicovitch sequence if $\{a(\mathbf{k})\} \in B(r) \cap l^\infty$. Let $\alpha = \sup_{\mathbf{k}} a_k$. We call that the Banach space X is without 1-projections iff there is no projection P on X such that $\|x\| = \|Px\| + \|x - Px\|$ for all $x \in X$.

In the vector case, R. V. Chacon proved the a.e. convergence of the averages $\frac{1}{n}\sum_{k=0}^{n-1} T^k f$ when X being reflexive Banach space, T is linear operator on $L^1(X)$ contraction in both $L^1(X)$ and in $L^\infty(X)$ and $f \in L^1(X)$. Yoshimoto [**13**] remarked that Chacon's theorem remains true if the operator T is contraction in $L^1(\Omega, X)$ but power bounded in $L^\infty(\Omega, X)$.

In [**8**] it was shown that Chacon's theorem remains true for the weighted averages $\frac{1}{n}\sum_{k=0}^{n-1} a(k)T^k f$ where $a(k)$ is a 1-Besicovitch bounded sequence.

In the real case ($X = \mathbf{R}$) and using linear modulus of non positive operator, R. Jones and J. Olsen proved [**11**] the a.e convergence of the averages $\frac{1}{|\mathbf{N}|}\sum_{\mathbf{k}=0}^{\mathbf{N}-1} a(\mathbf{k})\mathbf{T}^{\mathbf{k}} f$ with $\mathbf{N} = (n, \ldots, n)$ and $f \in L^1$ for Dunford-Schwartz operators. Akcoglu and Chacon proved [**1**] the almost everywhere convergence of the Cesaro average $A_n(T)f = \frac{1}{n}\sum_{i=0}^{n-1} T^i f$ for all $f \in L^1(\Omega, \mathbf{R})$, when T is a linear operator on $L^1(\Omega, \mathbf{R})$ contrction in $L^1(\Omega, \mathbf{R})$ and in $L^q(\Omega, \mathbf{R})$ for some $q \in]1, +\infty[$.

Using the linear modulus A. Brunel proved [**4**] the almost everywhere convergence of the multiparameter averages $A_n(T_1, \ldots, T_d)f = A_n(T_1)\ldots A_n(T_d)f$ when $T_1, \ldots, T_d$ are linear commuting operators on $L^1(\Omega, \mathbf{R})$ contraction in both $L^1(\Omega, \mathbf{R})$ and in $L^\infty(\Omega, \mathbf{R})$.

Our aim is to prove that if the operators $T_1, \ldots, T_d$ (possibly not commuting) are linear surjective isometries on $L^1(X)$ and contractions (or power bounded) in $L^\infty(X)$ then, we have a.e. convergence of the averages $\frac{1}{|\mathbf{N}|}\sum_{k=0}^{N-1} a(\mathbf{k})\mathbf{T}^{\mathbf{k}} f$ with $\mathbf{N} = (n, \ldots, n)$ for all $f \in L^1(X)$.

This result maybe considered to be a multidimensional version of Chacon's theorem [**5**] for surjective isometries in $L^1(X)$ and for the weighted averages, of course, the extension of Chacon's theorem to multidimensional case remains an open problem.

Using a representation of surjective isometries due to S. Guerre and Y. Raynaud [**10**] we give a vector multiparameter version of Jones-Olsen's result. In fact, the difficulty in vector case is that: for a vector operator T on $L^1(X)$ we cannot always find a DPO (dominated positive linear operator) τ, which is a contraction on $L^1 = L^1(\mathbf{R})$ (analog to the linear modulus) and verify that for all $f \in L^1(X)$

$$\|Tf\| \leq \tau(\|f\|). \tag{1}$$

(We give a counter example proving that the operators τ does not exist in general). In [**10**] Guerre and Raynaud proved in proposition 6.1 that if an isometry T admits a DPO, τ then we have $\|Tf\| = \tau(\|f\|)$ for all $f \in L^p(X)$.

Our main result is the following:

Theorem 1. *Let $T_1, \ldots, T_d$ be commuting linear surjective isometries on $L^1(X)$ and power bounded in $L^\infty(X)$. $a(\mathbf{k})$ is a d-dimensional r-Besicovitch bounded sequences. Then, for $f \in L^1(X)$ we have the almost everywhere convergence of the averages $\frac{1}{|\mathbf{N}|}\sum_{k=0}^{N-1} a(\mathbf{k})\mathbf{T}^{\mathbf{k}} f$ with $\mathbf{N} = (n, \ldots, n)$ as $n \to \infty$.*

We need the following lemma:

Lemma 1. *Let $T_1, \ldots, T_d$ be commuting linear surjective isometries on $L^1(X)$ and power bounded in $L^\infty(X)$. Let $F^* = \sup_N \left\| \frac{1}{|\mathbf{N}|}\sum_{\mathbf{k}=0}^{N-1} a(\mathbf{k})\mathbf{T}^{\mathbf{k}} f \right\|$ then, for all $f \in L^1(X)$ and any $a > 0$, there exists a positive real χ_d such that*

$$a\mu\{F^* > a\} \leq \chi_d \int_\Omega \|f\| \, d\mu.$$

Proof. We shall use the norm resolution [**10**, p. 368]. Let Z is Boole algebra (they are commuting). We define a measure μ_Z on $\mathcal{P}_1(Z)$ and a surjective mapping $N_Z : Z \to \mathcal{P}_1(Z)$ which is the norme resolution on $\mathcal{P}_1(Z)$ verifying

(*i*) $\forall z \in Z$, $\|z\| = |N_Z(z)|_{L^1_+(\mathcal{P}_1(Z))}$.

(*ii*) $\forall (u, v) \in Z^2, \forall (\alpha, \beta) \in R^2$

$$N_Z(\alpha u + \beta v) \leq |\alpha| \, N_Z(u) + |\beta| \, N_Z(v).$$

Guerre and Raynaud showed [**10**] that for each linear surjective isometry $T : Z \to Z$ that there is a positive surjective isometry

$$\tau : L^1_+(\mathcal{P}_1(Z)) \to L^1_+(\mathcal{P}_1(Z))$$

such that

$$N_Z(Tf) = \tau(N_Z f). \tag{2}$$

Let $Z = L^1(X) = L(\Omega, \beta, \mu, X)$. In [**9**] it was proved that if X is separable space (without loss of generality we can suppose that X is separable) then

$$\mathcal{P}_1(Z) = \beta \otimes \mathcal{P}_1(X).$$

Let $\mu_Z = \mu \otimes \mu_X$ with these notations we can write for $f \in L^1(X)$

$$N_Z(f) \in L^1(\beta \otimes \mathcal{P}_1(X)) = L^1_\beta\left(L^1_{\mathcal{P}_1(X)}\right)$$
$$N_Z(f)(\omega) = N_X(f(\omega))$$

and

$$\|f(\omega)\|_X = |N_X(f(\omega))|_{L^1_+(\mathcal{P}_1(Z))} = |N_z(f)(\omega)|_{L^1_+(\mathcal{P}_1(Z))}.$$

Denote by $N = N_Z = N_{L^1(X)}$, (2) shows that for all $j \in \mathbf{N}$ we have

$$\tau^j(Nf) = N(T^j f)$$

and since X is reflexive Banach space then it has a finite number of 1-projections (because otherwise $X \supset l^1$). Let

$$\mathcal{P}_1(X) = \Pi = \{1, \ldots, K\}.$$

We can decompose the space X as

$$X = X_1 \oplus^1 X_2 \oplus^1 \ldots \oplus^1 X_K$$

where X_i is a Banach space without 1-projections for $i = 1, \ldots, K$. If $x \in X$ then $x = (x_i)_{1 \leq i \leq K}$ and theorem of x will be

$$\|x\|_X = \sum_{i=1}^{K} \|x_i\|_{X_i}$$

where$\|x_i\|_{X_i}$ is the norm in the space X_i. We can write that the measure μ_X on the set $\{1, \ldots, K\}$ is a countable measure The space $L^1(\Omega \times \{1, \ldots, K\})$ identifies to the space $L^1(\Omega, l^1_K)$ and the norm resolutions $N = N_Z : L^1(X) \to L^1(\Omega \times \times \{1, \ldots, K\})$ and $N_X : X \to L^1_+(\{1, \ldots, K\}) = l^1_K$ are related by

$$N_X\left(f(\omega)\right)(i) = N(f)(\omega, i) = \|f_i(\omega)\|_{X_i}$$

and as $N_X\left(x\right))\left(i\right) = \|x_i\|_{X_i}$ we obtain

$$\begin{aligned} \|f_i(\omega)\|_{X_i} &= \sum_{i=1}^{K} \|f_i(\omega)\|_{X_i} = \sum_{i=1}^{K} N_X\left(f(\omega)\right)(i) \\ &= \sum_{i=1}^{K} Nf(\omega, i) = \|Nf(\omega, .)\|_{L^1(\Pi)} \end{aligned} \tag{3}$$

(3) gives the norm in X in terms of the norm resolution on $L^1(\Omega \times \{1, \ldots, K\})$. In its definition the operator τ acts in $L^1(\Omega \times \{1, \ldots, K\}) = L^1(\Omega, l^1_K)$ and verifies $\forall j \in \mathbf{N}$

$$\tau^j(Nf) = \tau^j\left[\left(\|f_i\|_{X_i}\right)_{1 \leq i \leq K}\right] = \left(\left\|T^j f_i\right\|_{X_i}\right)_{1 \leq i \leq K} = N(T^j f).$$

By (2) we have for $\varphi = Nf \in L^1(\Omega' = \Omega \times \{1, \ldots, K\})$ with $f \in L^1(\Omega, X)$

$$\begin{aligned} \|\tau\varphi\|_{L^1(\Omega')} &= \|\tau\left(Nf\right)\|_{L^1(\Omega')} = \|N(Tf)\|_{L^1(\Omega, X)} \\ &= \|Nf\|_{L^1(\Omega')} = \|f\|_{L^1(\Omega, X)} = \|\varphi\|_{L^1(\Omega')} \end{aligned}$$

which proves that τ is isometry on $L^1(\Omega, l^1_K)$. In what follows we will prove that $\sup_N \left\| \frac{1}{|\mathbf{N}|} \sum_{k=0}^{N-1} a(\mathbf{k}) \mathbf{T}^{\mathbf{k}} f \right\|$ is finite a.e. for all $f \in L^1(\Omega, X)$. □

To complete the proof we need the following proposition:

Proposition 1. *Let T be a surjective isometry in $L^1(X)$ and contracting in $L^q(X)$ $(1 \leq q \leq \infty)$, then its DPO τ is a contraction on $L^q(l^1_k)$ and if T is power bounded in $L^q(X)$ then is power bounded in $L^q(\Omega, \{1, \ldots, k\}) = L^q(\Omega, l^1_k)$.*

Proof. Let $q \in]1,+\infty[$. If T is a contraction in $L^q(X)$,we can write

$$\begin{aligned}
\|\tau(Nf)\|_{L^q(\Omega')} &= \left[\int\left(\sum_{i=1}^{K}\tau(Nf)(\omega,i)\right)^q d\omega\right]^{1/q} \\
&= \left[\int\left(\sum_{i=1}^{K}N(Tf)(\omega,i)\right)^q d\omega\right]^{1/q} \\
&= \left[\int\left(\sum_{i=1}^{K}\|(Tf(\omega))_i\|_{X_i}\right)^q d\omega\right]^{1/q} = \left[\int\|Tf(\omega)\|_X^q\, d\omega\right]^{1/q} \\
&= \|Tf\|_{L^q(X)} \le \|f\|_{L^q(X)} = \|\|f\|_X\|_{L^q} \\
&= \left\|\sum_{i=1}^{K}\|f\|_i\right\|_{L^q} = \|Nf\|_{L^q(l_k^1)}.
\end{aligned}$$

If T is power bounded in $L^q(X)$ we can write

$$\begin{aligned}
\left\|\tau^j(Nf)\right\|_{L^q(\Omega')} &= \left\|N(T^jf)\right\|_{L^q(l_K^1)} = \left\|T^jf\right\|_{L^q(X)} \\
&\le \|f\|_{L^q(X)} = \|Nf\|_{L^q(l_K^1)}.
\end{aligned}$$

Clearly

$$\|(.)\|_{L^q(l_K^q)} \le \|(.)\|_{L^q(l_K^1)} \le K^{\frac{q-1}{q}}\|(.)\|_{L^q(l_K^q)}$$

and then we get

$$\left\|\tau^j(Nf)\right\|_{L^q(l_K^q)} \le \left\|\tau^j(Nf)\right\|_{L^q(l_K^1)} \le cK^{\frac{q-1}{q}}\|Nf\|_{L^q(l_K^q)}$$

which proves that τ is power bounded in $L^q(\Omega')$, $1 < q \le \infty$.

We have two cases to consider:

Case 1. The space X is without 1-projections.

By [**10**] if T_j, $j = 1,\dots,d$, are surjective isometries on $L^1(X)$ and contraction (resp. power bounded) in $L^q(X)$, $1 < q \le \infty$, then, by Proposition 1.6 in [**10**] (resp. Proposition 2) they are majorizable by (DPO) τ_j, $j = 1,\dots,d$, which are isometries in L^1 and contraction in L^q (resp. power bounded in $L^q(\Omega')$, $1 < q \le \infty$.

To the operators $\tau_1,\dots,\tau_d$ we associate the Brunel operator U which is a contraction in L^1 and a contraction (resp. a power bounded) in $L^q, 1 < q \le \infty$. Moreover, the operator U verifies [**12**, p. 213]

$$\begin{aligned}
&\sup_n\left\|\frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1}a(\mathbf{k})\mathbf{T}^{\mathbf{k}}f\right\| = \sup_n\left\|\frac{1}{\mathbf{n}^d}\sum_{\mathbf{k}=0}^{\mathbf{N}-1}a(\mathbf{k})\mathbf{T}^{\mathbf{k}}f\right\| \\
&\le \sup_n\left\|\frac{1}{\mathbf{n}^d}\sum_{k=0}^{\mathbf{N}-1}a(\mathbf{k})\tau^{\mathbf{k}}f\right\| \le \alpha\sup_n\left\|\frac{1}{\mathbf{n}^d}\sum_{k=0}^{\mathbf{N}-1}\tau^{\mathbf{k}}\|f\|\right\| \\
&\le \alpha\chi_d\sup_n A_n(U)\|f\|.
\end{aligned}$$

In this case we obtain by applying the maximal weak inequality to the operator U (which is Dunford-Schwarz operators)

$$a\mu\left\{F^*>a\right\}\leq a\mu\left\{\sup_n A_n(U)\left\|f\right\|>a/\alpha\chi_d\right\}\leq\alpha\chi_d\int_\Omega\left\|f\right\|d\mu.$$

Case 2. The space X is reflexive Banach space:

The operators $\tau_1,\ldots,\tau_d$ are isometries in $L^1(\Omega'=\Omega\otimes\Pi)$ and power bounded in $L^\infty(\Omega')$ apply the Proposition 2 with $q=\infty$ to obtain that Brunel's operator U associated to the family $\tau_1,\ldots,\tau_d$ is also a contraction in L^1 and power bounded in L^∞. Using (3) and the fact that μ_X is a countable measure we get

$$\begin{aligned}\sup_n\left\|\frac{1}{\mathbf{n}^d}\sum_{k=0}^{N-1}a(\mathbf{k})\mathbf{T}^{\mathbf{k}}f(\omega)\right\|&\leq\sup_n\sum_{i=1}^{K}N\left(\frac{1}{\mathbf{n}^d}\sum_{k=0}^{N-1}a(\mathbf{k})\mathbf{T}^{\mathbf{k}}f\right)(\omega,i)\\&\leq\sum_{i=1}^{K}\sup_n\left(\frac{1}{\mathbf{n}^d}\sum_{k=0}^{N-1}a(\mathbf{k})\tau^{\mathbf{k}}\right)(Nf)(\omega,i)\\&\leq\chi_d\sum_{i=1}^{K}\sup_n A_n(U)Nf(\omega,i).\end{aligned}\tag{4}$$

By Yoshimoto [**13**], we can write

$$\begin{aligned}a\mu\left\{F^*>a\right\}&\leq a\mu\left\{\sum_{i=1}^{K}\sup_n A_n(U)Nf(\omega,i)>a/\chi_d\right\}\\&\leq a\mu\left\{\bigcup_{i=1}^{K}\left\{\sup_n A_n(U)Nf(\omega,i)>a/k\chi_d\right\}\right\}\\&\leq\sum_{i=1}^{K}a\mu\left\{\sup_n A_n(U)Nf(\omega,i)>a/k\chi_d\right\}\\&\leq\chi_d\sum_{i=1}^{K}\int_{\Omega'}Nf(\omega,i)d\left(\mu\otimes\mu_X\right)\\&=\chi_d\left\|Nf\right\|_{L^1(\Omega')}\\&=\alpha\chi_d\left\|f\right\|_{L^1(\Omega,X)}=\alpha\chi_d\int_\Omega\left\|f\right\|d\mu.\end{aligned}$$

□

Before giving the proof of our main result, we state a result of Jones-Olsen in [**11**, pp. 351]

Theorem 2. *For all $r\geq1$ we have $B(r)\cap l^\infty=B(1)\cap l^\infty$.*

By this theorem it suffices to prove theorem 1 for the 1-Besicovitch bounded sequences.

Proof of Theorem 1. We have to prove that the averages $\frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} a(\mathbf{k})\mathbf{T}^{\mathbf{k}}f$ converge a.e. on a dense set on $L^1(X)$. Let $L^\infty(X)$ such a set. For every $\varepsilon > 0$ we have

$$\frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} a(\mathbf{k})\mathbf{T}^{\mathbf{k}}f = \frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \left(a(\mathbf{k}) - \psi_\varepsilon(\mathbf{k})\right)\mathbf{T}^{\mathbf{k}}f + \frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \psi_\varepsilon(\mathbf{k})\mathbf{T}^{\mathbf{k}}f$$

and then for all $f \in L^\infty(X)$ we have

(5)

$$\begin{aligned}\left\|\frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} a(\mathbf{k})\mathbf{T}^{\mathbf{k}}f\right\| &\leq \left\|\frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \left(a(\mathbf{k}) - \psi_\varepsilon(\mathbf{k})\right)\mathbf{T}^{\mathbf{k}}f\right\| + \left\|\frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \psi_\varepsilon(\mathbf{k})\mathbf{T}^{\mathbf{k}}f\right\| \\ &\leq \frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \left|a(\mathbf{k}) - \psi_\varepsilon(\mathbf{k})\right| \left\|\mathbf{T}^{\mathbf{k}}f\right\| + \left\|\frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \psi_\varepsilon(\mathbf{k})\mathbf{T}^{\mathbf{k}}f\right\|.\end{aligned}$$

In the case 1 we have

$$\begin{aligned}\frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \left|a(\mathbf{k}) - \psi_\varepsilon(\mathbf{k})\right| \left\|\mathbf{T}^{\mathbf{k}}f\right\| &\leq \frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \left|a(\mathbf{k}) - \psi_\varepsilon(\mathbf{k})\right| \tau^k \|f\| \\ &\leq \|f\|_\infty \frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \left|a(\mathbf{k}) - \psi_\varepsilon(\mathbf{k})\right| < \varepsilon \|f\|_\infty.\end{aligned}$$

In the case 2 we have

$$\begin{aligned}\frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \left|a(\mathbf{k}) - \psi_\varepsilon(\mathbf{k})\right| \left\|\mathbf{T}^{\mathbf{k}}f\right\| &\leq \frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \left|a(\mathbf{k}) - \psi_\varepsilon(\mathbf{k})\right| \left[\sum_{i=1}^{K} N\left(\mathbf{T}^{\mathbf{k}}f\right)(.,i)\right] \\ &= \frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \left|a(\mathbf{k}) - \psi_\varepsilon(\mathbf{k})\right| \left[\sum_{i=1}^{K} \tau^{\mathbf{k}} N f(.,i)\right] \\ &\leq \varepsilon \left[\sum_{i=1}^{K} \tau \|Nf(.,i)\|_\infty\right].\end{aligned}$$

We have seen in [**7**, pp. 28] that a. e. convergence holds for the averages $\frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} \mathbf{T}^{\mathbf{k}}f$. Let $(\lambda_1, \ldots, \lambda_d) \in \mathbb{C}^d$, with $|\lambda_i| = 1$, $i = 1, 2, \ldots, d$ then the theorem holds when $a(k) = \lambda_1^{k_1} \ldots \lambda_d^{k_d}$ since $\widetilde{\mathbf{T}}^{\mathbf{k}} = \lambda_1^{k_1} \ldots \lambda_d^{k_d}\mathbf{T}^{\mathbf{k}}$ is also a d-parameter sequences of surjective isometries operators on $L^1(X)$ and power bounded in $L^\infty(X)$ when $a(\mathbf{k}) = 1$. Clearly the a.e. convergence holds for finite linear combinations of such sequences, and hence holds for trigonometric ploynomial in d variables, which proves the convergence a.e. of the second term of (5) and then we have a.e. convergence of

$$\frac{1}{|\mathbf{N}|}\sum_{k=0}^{\mathbf{N}-1} a(\mathbf{k})\mathbf{T}^{\mathbf{k}}f$$

for all $f \in L^{\infty}(X)$. The Banach principle combining with Lemma 1 end the proof of Theorem 1 in the contraction (resp. power bounded) case by applying Akcoglu-Chacon's Theorem (resp. Yoshimoto's) Theorem to the operator U.

Remark 1. In the case when X is without 1-projections we obtain that Akcoglu-Chacon's theorem [**2**] can be extended to vector case and for linear surjective isometries.

Remark 2. If $p = 2$, Burkholder [**12**] constructed a surjective isometry in L^2 for which the pointwise ergodic theorem is false. For this reason we can prove that if X is reflexive Banach lattice, and if $T_1, \ldots, T_d$ are (non-commuting) surjective isometries on $L^p(X)$, $1 < p \neq 2 < \infty$, then $\forall f \in L^p(X)$:

$$\begin{aligned}
\left\| \sup_{\overrightarrow{N}} \left\| \frac{1}{|\mathbf{N}|} \sum_{k=0}^{N-1} a(\mathbf{k}) \mathbf{T}^{\mathbf{k}} f \right\| \right\|_{L^p(\Omega,X)} &\leq \left\| \left\| \sup_{\overrightarrow{N}} N_X \left(\frac{1}{|\mathbf{N}|} \sum_{k=0}^{N-1} a(\mathbf{k}) \mathbf{T}^{\mathbf{k}} f \right) \right\|_{L^p(\Pi)} \right\|_{L^p(\Omega,X)} \\
&= \left\| \sup_{\overrightarrow{N}} N_X \left(\frac{1}{|\mathbf{N}|} \sum_{k=0}^{N-1} a(\mathbf{k}) \mathbf{T}^{\mathbf{k}} f \right) \right\|_{L^p(\Pi \otimes \Omega)} \\
&\leq \left\| \sup_{\overrightarrow{N}} \frac{1}{|\mathbf{N}|} \sum_{k=0}^{N-1} a(\mathbf{k}) \tau^{\mathbf{k}} (Nf) \right\|_{L^p(\Pi \otimes \Omega)} \\
&\leq \alpha \left(\frac{p}{p-1} \right)^d \|Nf\|_{L^p(\Pi \otimes \Omega)} \\
&= \alpha \left(\frac{p}{p-1} \right)^d \|f\|_{L^p(\Omega,X)} .
\end{aligned}$$

The least inequality is true by applying Akcoglu's theorem d times successively on the operators $\tau_1, \ldots, \tau_d$.

For the commuting case. Using Brunel operator we obtain the following strong estimates

$$\left\| \sup_{\overrightarrow{N}} \left\| A_{\overrightarrow{N}}(T_1, \ldots, T_d) f \right\| \right\|_{L^p(\Omega,X)} \leq \chi_d \frac{p}{p-1} \|f\|_{L^p(\Omega,X)} .$$

Remark 3. 1. If T is power bounded in $L^q(X)$ then its DPO; τ is power bounded in $L^q(\Omega')$. But we don't know if τ is a contraction in $L^q(\Omega') = L^q(l^q_K)$ when T is in $L^q(X)$. For this reason in the case "X has 1-projection", we cannot obtain an extension of Akcoglu-Chacon's theorem.

2. If $\mathcal{P}_1(X)$ is singleton and if T is a contraction in $L^q(X)$, then τ is contraction in $L^q(\Omega')$, $1 < q < \infty$.

Now, we can find the results of [**7**] and [**8**] as corollary:

Corollary 1. *Let $T_1, \ldots, T_d$ be d commuting linear surjective isometries on $L^1(X)$ and power bounded in $L^{\infty}(X)$. Then, for $f \in L^p(X)$ we have the almost*

everywhere convergence of the averages

$$A_n(T_1,\dots,T_d)f=\frac{1}{n^d}\sum_{i_1=0}^{n-1}\dots\sum_{i_d=0}^{n-1}T^{i_1}\dots T^{i_d}f$$

as $n\to\infty$.

Corollary 2. *Let* X *be a reflexive Banach space and let* $\theta_1,\dots,\theta_d$ *be* d *preserving measure transformations on a* $\sigma-$*finite space. If* $T_jf=fo\theta_j$, $j=1,\dots,d$, *and if the transformations are commuting then the averages* $A_n(T_1,\dots,T_d)f$ *converge a.e. for all* $f\in L^1(X)$.

In the following we give an example of vector operator which is not majorizable by a dominated positive operator contraction in L^1.

Example 1. Let $\Omega=\{1,2\}$ be probability space, $\mu(1)=\mu(2)=1/2$, and let $X=\mathbf{R}^2$ (reflexive Banach space) with norm $\|(x,y)\|=|x|+|y|$. The Banach space $L^1\left(\{1,2\},\mathbf{R}^2\right)$ will be of dimension 4. The operator T will be represented by a square matrix of order 4. Let $T=(a_{ij})_{1\le i,j\le 4}$ if such an operator (DPO) τ exists then it will verify for all $\varphi\in L^1$

$$\tau_0\varphi=\sup\{\|Tf\|;\ \ \|f\|\le\varphi\}\le\tau\varphi.$$

We shall prove that for an operator T, τ_0 is not contracting in L^1. From this we deduce that τ is also not contracting. The condition $\|Tf\|_1\le\|f\|_1$ implies that $\sum_{i=1}^4|a_{ij}|\le 1$ for $j=1,\dots,4$. The operator τ_0 is contracting in L^1 if

$$(*)\qquad |a_{11}|+|a_{21}|+|a_{32}|+|a_{42}|\le 1$$

$$(**)\qquad |a_{12}|+|a_{22}|+|a_{31}|+|a_{41}|\le 1.$$

But if we take $a_{11}=1/2$; $a_{21}=a_{31}=a_{41}=10^{-1}$ and $a_{12}=a_{22}=a_{32}=a_{42}=1/4$ and 0 elsewhere then while the operator T is $L^1(X)-$contraction we have the conditions $(*)$ and $(**)$ are not satisfied. This implies that τ_0 is not L^1-contraction. Consequently, τ is not L^1-contraction. A similar calculation shows that τ_0 is not L^p-contraction although the operator T is. Since $X=l_2^1$ has two projections, by its definition, DPO and as τ_0 is not contraction in $L^1(\Omega')$, we then get that T does not have a DPO. By Chacon's theorem the Cesaro averages $A_n(T)f$ converge a.e. for all $f\in L^1(X)$ while the operator T is not majorilized by a DPO.

Remark 4.

1. By a similar argument of that of Brunel [**4**] Theorem 1 remains true when X is a reflexive Banach lattice space and the operators $T_1,\dots,T_d$ are positive linear operators contraction in both $L^1(X)$ and in $L^\infty(X)$.
2. The commutation of $\tau_1,\dots,\tau_d$ is not needed to the construction of the Brunel operator U.
3. If X is without1-projections the contraction of the operator T in $L^\infty(X)$ assumed in [**5**] can be replaced by the contraction in some $L^q(X)$ with $q\in]1,+\infty[$.

References

1. Akcoglu M. A., *A pointwise Ergodic Theorem in L^p-Space,* Canad. J. Math., **27** (1975), 1075–1082.

2. Akcoglu M. A. and Chacon R. V., *A convexity theorem for positive operators,* ZW **3** 328–332.

3. Beherend E., *Structure in real Banach space,* Lecture Notes in Math. **613** (Spring Verlage).

4. Brunel A., *Theoreme ergodiques ponctuel pour un semi groupe commutatif finiment engendre de contractions* AIHP B **9** (1973), 327–343.

5. Chacon R. V., *A vector ergodic theorem for operators satisfying norm conditions.* J. Math. **11** (1962), 165–172.

6. El Berdan K., *Theoreme ergodique vectoriel pour les transformations ponctuelles preservant la mesure,* C.R.A.S. Paris t. **317**, serie I (1993), 311–313.

7. _____, *Theoremes ergodiques a plusieurs parametres dans les espaces de Banach,* These d'Universite Paris 6. 1995.

8. _____, *The Vector Valued Ergodic Theorem for Bounded Besicovich sequences,* Acta Math. Univ. Comenianae Vol. **68**, 1(1999), 1–15.

9. Greim P., *Isometries and Structure of separability valued Bochner spaces Conference on measure Theory and Applications,* (Sherbook 1982) Lecture Notes (Springer Verlag).

10. Guerre S. and Ranaud Y., *Sur les isometries de $L^p(X)$ et le theoreme ergodique vectoriel.* Canad. J. Math. Vol. **XL**, N. 2 (1988), 306–391.

11. Jones R. L. and Olsen J., *Multiparameter Weighted Ergodic Theorems,* Can. J. Math. Vol. **46** (2) (1994), 343–35.

12. Krengel U., *Ergodic Theorems,* Walter de Guyter Berlin – New York 1985.

13. Yoshimoto T., *Vector valued ergodic theorem for operators satisfying norm conditions,* Pac. JM **85** (1980), 485–499.

K. El Berdan, Lebanese University Faculty of Sciences (I) Departement of Mathematics Hadeth Beirut, Lebanon, *e-mail*: kberdan@terra.net.lb