

Research Article

Principal Functions of Non-Selfadjoint Sturm-Liouville Problems with Eigenvalue-Dependent Boundary Conditions

Nihal Yokuş

Department of Mathematics, Karamanoğlu Mehmetbey University, 70100 Karaman, Turkey

Correspondence should be addressed to Nihal Yokuş, nyokus@kmu.edu.tr

Received 8 March 2011; Accepted 5 April 2011

Academic Editor: Narcisa C. Apreutesei

Copyright © 2011 Nihal Yokuş. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider the operator L generated in $L^2(\mathbb{R}_+)$ by the differential expression $l(y) = -y'' + q(x)y$, $x \in \mathbb{R}_+ := [0, \infty)$ and the boundary condition $y'(0)/y(0) = \alpha_0 + \alpha_1\lambda + \alpha_2\lambda^2$, where q is a complex-valued function and $\alpha_i \in \mathbb{C}$, $i = 0, 1, 2$ with $\alpha_2 \neq 0$. In this paper we obtain the properties of the principal functions corresponding to the spectral singularities of L .

1. Introduction

Let T be a nonselfadjoint, closed operator in a Hilbert space H . We will denote the continuous spectrum and the set of all eigenvalues of T by $\sigma_c(T)$ and $\sigma_d(T)$, respectively. Let us assume that $\sigma_c(T) \neq \emptyset$.

Definition 1.1. If $\lambda = \lambda_0$ is a pole of the resolvent of T and $\lambda_0 \in \sigma_c(T)$, but $\lambda_0 \notin \sigma_d(T)$, then λ_0 is called a spectral singularity of T .

Let us consider the nonselfadjoint operator L_0 generated in $L^2(\mathbb{R}_+)$ by the differential expression

$$l_0(y) = -y'' + q(x)y, \quad x \in \mathbb{R}_+, \quad (1.1)$$

and the boundary condition $y(0) = 0$, where q is a complex-valued function. The spectrum and spectral expansion of L_0 were investigated by Naïmark [1]. He proved that the spectrum of L_0 is composed of continuous spectrum, eigenvalues, and spectral singularities. He showed that spectral singularities are on the continuous spectrum and are the poles of the resolvent kernel, which are not eigenvalues.

Lyance investigated the effect of the spectral singularities in the spectral expansion in terms of the principal functions of L_0 [2, 3]. He also showed that the spectral singularities play an important role in the spectral analysis of L_0 .

The spectral analysis of the non-self-adjoint operator L_1 generated in $L^2(\mathbb{R}_+)$ by (1.1) and the boundary condition

$$\int_0^{\infty} K(x)y(x)dx + \alpha y'(0) - \beta y(0) = 0, \quad (1.2)$$

in which $K \in L^2(\mathbb{R}_+)$ is a complex valued function and $\alpha, \beta \in \mathbb{C}$, was investigated in detail by Krall [4–8]. In [4] he obtained the adjoint L_1^* of the operator L_1 . Note that L_1^* deserves special interest, since it is not a purely differential operator. The eigenfunction expansions of L_1 and L_1^* were investigated in [5].

In [9] the results of Naimark were extended to the three-dimensional Schrödinger operators.

The Laurent expansion of the resolvents of the abstract non-self-adjoint operators in the neighborhood of the spectral singularities was studied in [10].

Using the boundary uniqueness theorems of analytic functions, the structure of the eigenvalues and the spectral singularities of a quadratic pencil of Schrödinger, Klein-Gordon, discrete Dirac, and discrete Schrödinger operators was investigated in [11–20]. By regularization of a divergent integral, the effect of the spectral singularities in the spectral expansion of a quadratic pencil of Schrödinger operators was obtained in [13]. In [19, 20] the spectral expansion of the discrete Dirac and Schrödinger operators with spectral singularities was derived using the generalized spectral function (in the sense of Marchenko [21]) and the analytical properties of the Weyl function.

Let L denote the operator generated in $L^2(\mathbb{R}_+)$ by the differential expression

$$l(y) = -y'' + q(x)y, \quad x \in \mathbb{R}_+ \quad (1.3)$$

and the boundary condition

$$\frac{y'(0)}{y(0)} = \alpha_0 + \alpha_1 \lambda + \alpha_2 \lambda^2, \quad (1.4)$$

where q is a complex-valued function and $\alpha_i \in \mathbb{C}$, $i = 0, 1, 2$ with $\alpha_2 \neq 0$. In this work we obtain the properties of the principal functions corresponding to the spectral singularities of L .

2. The Jost Solution and Jost Function

We consider the equation

$$-y'' + q(x)y = \lambda^2 y, \quad x \in \mathbb{R}_+ \quad (2.1)$$

related to the operator L .

Now we will assume that the complex valued function q is almost everywhere continuous in \mathbb{R}_+ and satisfies the following:

$$\int_0^\infty x|q(x)|dx < \infty. \tag{2.2}$$

Let $\varphi(x, \lambda)$ and $e(x, \lambda)$ denote the solutions of (2.1) satisfying the conditions

$$\begin{aligned} \varphi(0, \lambda) &= 1, & \varphi'(0, \lambda) &= \alpha_0 + \alpha_1\lambda + \alpha_2\lambda^2, \\ \lim_{x \rightarrow \infty} e(x, \lambda)e^{-i\lambda x} &= 1, & \lambda &\in \overline{\mathbb{C}}_+, \end{aligned} \tag{2.3}$$

respectively. The solution $e(x, \lambda)$ is called the Jost solution of (2.1). Note that, under the condition (2.2), the solution $\varphi(x, \lambda)$ is an entire function of λ and the Jost solution is an analytic function of λ in $\mathbb{C}_+ := \{\lambda : \lambda \in \mathbb{C}, \text{Im } \lambda > 0\}$ and continuous in $\overline{\mathbb{C}}_+ = \{\lambda : \lambda \in \mathbb{C}, \text{Im } \lambda \geq 0\}$.

In addition, Jost solution has a representation ([22])

$$e(x, \lambda) = e^{i\lambda x} + \int_x^\infty K(x, t)e^{i\lambda t} dt, \quad \lambda \in \overline{\mathbb{C}}_+, \tag{2.4}$$

where the kernel $K(x, t)$ satisfies

$$\begin{aligned} K(x, t) &= \frac{1}{2} \int_{(x+t)/2}^\infty q(s)ds + \frac{1}{2} \int_x^{(x+t)/2} \int_{t+x-s}^{t+s-x} q(s)K(s, u)du ds \\ &+ \frac{1}{2} \int_{(x+t)/2}^\infty \int_s^{t+s-x} q(s)K(s, u)du ds \end{aligned} \tag{2.5}$$

and $K(x, t)$ is continuously differentiable with respect to its arguments. We also have

$$|K(x, t)| \leq cw \left(\frac{x+t}{2} \right), \tag{2.6}$$

$$|K_x(x, t)|, |K_t(x, t)| \leq \frac{1}{4} \left| q \left(\frac{x+t}{2} \right) \right| + cw \left(\frac{x+t}{2} \right), \tag{2.7}$$

where $w(x) = \int_x^\infty |q(s)|ds$ and $c > 0$ is a constant.

Let $\widehat{e}^\pm(x, \lambda)$ denote the solutions of (2.1) subject to the conditions

$$\lim_{x \rightarrow \infty} e^{\pm i\lambda x} \widehat{e}^\pm(x, \lambda) = 1, \quad \lim_{x \rightarrow \infty} e^{\pm i\lambda x} \widehat{e}_x^\pm(x, \lambda) = \pm i\lambda, \quad \lambda \in \overline{\mathbb{C}}_\pm. \tag{2.8}$$

Then

$$\begin{aligned} W[e(x, \lambda), \widehat{e}^\pm(x, \lambda)] &= \mp 2i\lambda, \quad \lambda \in \mathbb{C}_\pm, \\ W[e(x, \lambda), e(x, -\lambda)] &= -2i\lambda, \quad \lambda \in \mathbb{R} = (-\infty, \infty), \end{aligned} \tag{2.9}$$

where $W[f_1, f_2]$ is the Wronskian of f_1 and f_2 , ([23]).

We will denote the Wronskian of the solutions $\varphi(x, \lambda)$ with $e(x, \lambda)$ and $e(x, -\lambda)$ by $E^+(\lambda)$ and $E^-(\lambda)$, respectively, where

$$\begin{aligned} E^+(\lambda) &:= e'(0, \lambda) - (\alpha_0 + \alpha_1\lambda + \alpha_2\lambda^2)e(0, \lambda), \quad \lambda \in \overline{\mathbb{C}}_+, \\ E^-(\lambda) &:= e'(0, -\lambda) - (\alpha_0 + \alpha_1\lambda + \alpha_2\lambda^2)e(0, -\lambda), \quad \lambda \in \overline{\mathbb{C}}_-, \end{aligned} \quad (2.10)$$

and $\overline{\mathbb{C}}_- = \{\lambda : \lambda \in \mathbb{C}, \text{Im } \lambda \leq 0\}$. Therefore E^+ and E^- are analytic in \mathbb{C}_+ and $\mathbb{C}_- = \{\lambda : \lambda \in \mathbb{C}, \text{Im } \lambda < 0\}$, respectively, and continuous up to real axis.

The functions E^+ and E^- are called Jost functions of L .

3. Eigenvalues and Spectral Singularities of L

Let

$$G(x, t; \lambda) = \begin{cases} G^+(x, t; \lambda), & \lambda \in \mathbb{C}_+, \\ G^-(x, t; \lambda), & \lambda \in \mathbb{C}_- \end{cases} \quad (3.1)$$

be the Green function of L (obtained by the standard techniques), where

$$\begin{aligned} G^+(x, t; \lambda) &= \begin{cases} -\frac{\varphi(t, \lambda)e(x, \lambda)}{E^+(\lambda)}, & 0 \leq t \leq x \\ -\frac{\varphi(x, \lambda)e(t, \lambda)}{E^+(\lambda)}, & x \leq t < \infty \end{cases} \\ G^-(x, t; \lambda) &= \begin{cases} -\frac{\varphi(t, \lambda)e(x, -\lambda)}{E^-(\lambda)}, & 0 \leq t \leq x \\ -\frac{\varphi(x, \lambda)e(t, -\lambda)}{E^-(\lambda)}, & x \leq t < \infty. \end{cases} \end{aligned} \quad (3.2)$$

We will denote the set of eigenvalues and spectral singularities of L by $\sigma_d(L)$ and $\sigma_{ss}(L)$, respectively. From (3.1)–(3.2)

$$\begin{aligned} \sigma_d(L) &= \{\lambda : \lambda \in \mathbb{C}_+, E^+(\lambda) = 0\} \cup \{\lambda : \lambda \in \mathbb{C}_-, E^-(\lambda) = 0\}, \\ \sigma_{ss}(L) &= \{\lambda : \lambda \in \mathbb{R}^*, E^+(\lambda) = 0\} \cup \{\lambda : \lambda \in \mathbb{R}^*, E^-(\lambda) = 0\}, \end{aligned} \quad (3.3)$$

where $\mathbb{R}^* = \mathbb{R} \setminus \{0\}$.

From (3.3) we obtain that to investigate the structure of the eigenvalues and the spectral singularities of L , we need to discuss the structure of the zeros of the functions E^+ and E^- in $\overline{\mathbb{C}}_+$ and $\overline{\mathbb{C}}_-$, respectively.

Definition 3.1. The multiplicity of zero of the function E^+ (or E^-) in $\overline{\mathbb{C}}_+$ (or $\overline{\mathbb{C}}_-$) is called the multiplicity of the corresponding eigenvalue and spectral singularity of L .

We see from (2.9) that the functions

$$\begin{aligned}\psi^+(x, \lambda) &= \frac{\widehat{E}^+(\lambda)}{2i\lambda} e(x, \lambda) - \frac{E^+(\lambda)}{2i\lambda} \widehat{e}^+(x, \lambda), \quad \lambda \in \mathbb{C}_+, \\ \psi^-(x, \lambda) &= \frac{\widehat{E}^-(\lambda)}{2i\lambda} e(x, -\lambda) - \frac{E^-(\lambda)}{2i\lambda} \widehat{e}^-(x, \lambda), \quad \lambda \in \mathbb{C}_-, \\ \psi(x, \lambda) &= \frac{E^+(\lambda)}{2i\lambda} e(x, -\lambda) - \frac{E^-(\lambda)}{2i\lambda} e(x, \lambda), \quad \lambda \in \mathbb{R}^*\end{aligned}\tag{3.4}$$

are the solutions of the boundary value problem

$$\begin{aligned}-y'' + q(x)y &= \lambda^2 y, \quad x \in \mathbb{R}_+, \\ \frac{y'(0)}{y(0)} &= \alpha_0 + \alpha_1 \lambda + \alpha_2 \lambda^2,\end{aligned}\tag{3.5}$$

where

$$\widehat{E}^\pm(\lambda) = \widehat{e}^\pm(0, \lambda) - (\alpha_0 + \alpha_1 \lambda + \alpha_2 \lambda^2) \widehat{e}^\pm(0, \lambda).\tag{3.6}$$

Now let us assume that

$$q \in AC(\mathbb{R}_+), \quad \lim_{x \rightarrow \infty} q(x) = 0, \quad \sup_{x \in \mathbb{R}_+} [e^{\varepsilon \sqrt{x}} |q'(x)|] < \infty, \quad \varepsilon > 0.\tag{3.7}$$

Theorem 3.2 (see [24]). *Under the condition (3.7) the operator L has a finite number of eigenvalues and spectral singularities, and each of them is of a finite multiplicity.*

4. Principal Functions of L

In this section we assume that (3.7) holds. Let $\lambda_1, \dots, \lambda_j$ and $\lambda_{j+1}, \dots, \lambda_k$ denote the zeros of E^+ in \mathbb{C}_+ and E^- in \mathbb{C}_- (which are the eigenvalues of L) with multiplicities m_1, \dots, m_j and m_{j+1}, \dots, m_k , respectively. It is obvious that from definition of the Wronskian

$$\left\{ \frac{d^n}{d\lambda^n} W[\psi^+(x, \lambda), e(x, \lambda)] \right\}_{\lambda=\lambda_p} = \left\{ \frac{d^n}{d\lambda^n} E^+(\lambda) \right\}_{\lambda=\lambda_p} = 0\tag{4.1}$$

for $n = 0, 1, \dots, m_p - 1, p = 1, 2, \dots, j$, and

$$\left\{ \frac{d^n}{d\lambda^n} W[\psi^-(x, \lambda), e(x, -\lambda)] \right\}_{\lambda=\lambda_p} = \left\{ \frac{d^n}{d\lambda^n} E^-(\lambda) \right\}_{\lambda=\lambda_p} = 0\tag{4.2}$$

for $n = 0, 1, \dots, m_p - 1, p = j + 1, \dots, k$.

Theorem 4.1. *The following formulae:*

$$\left\{ \frac{\partial^n}{\partial \lambda^n} \psi^+(x, \lambda) \right\}_{\lambda=\lambda_p} = \sum_{m=0}^n A_m(\lambda_p) \left\{ \frac{\partial^m}{\partial \lambda^m} e(x, \lambda) \right\}_{\lambda=\lambda_p}, \quad (4.3)$$

$n = 0, 1, \dots, m_p - 1, p = 1, 2, \dots, j$, where

$$A_m(\lambda_p) = \binom{n}{m} \left\{ \frac{\partial^{n-m}}{\partial \lambda^{n-m}} \widehat{E}^+(\lambda) \right\}_{\lambda=\lambda_p}, \quad (4.4)$$

$$\left\{ \frac{\partial^n}{\partial \lambda^n} \psi^-(x, \lambda) \right\}_{\lambda=\lambda_p} = \sum_{m=0}^n B_m(\lambda_p) \left\{ \frac{\partial^m}{\partial \lambda^m} e(x, -\lambda) \right\}_{\lambda=\lambda_p}, \quad (4.5)$$

$n = 0, 1, \dots, m_p - 1, p = j + 1, \dots, k$, where

$$B_m(\lambda_p) = \binom{n}{m} \left\{ \frac{\partial^{n-m}}{\partial \lambda^{n-m}} \widehat{E}^-(\lambda) \right\}_{\lambda=\lambda_p} \quad (4.6)$$

holds.

Proof. We will proceed by mathematical induction, we prove first (4.3). Let $n = 0$. From (4.1) we get

$$\psi^+(x, \lambda_p) = a_0(\lambda_p) \cdot e(x, \lambda_p), \quad (4.7)$$

where $a_0(\lambda_p) \neq 0$. Let us assume that for $1 \leq n_0 \leq m_p - 2$, (4.3) holds; that is,

$$\left\{ \frac{\partial^{n_0}}{\partial \lambda^{n_0}} \psi^+(x, \lambda) \right\}_{\lambda=\lambda_p} = \sum_{m=0}^{n_0} A_m(\lambda_p) \left\{ \frac{\partial^m}{\partial \lambda^m} e(x, \lambda) \right\}_{\lambda=\lambda_p}. \quad (4.8)$$

Now we will prove that (4.3) holds for $n_0 + 1$. If $y(x, \lambda)$ is a solution of (2.1), then $(\partial^n / \partial \lambda^n) y(x, \lambda)$ satisfies

$$\left[-\frac{d^2}{dx^2} + q(x) - \lambda^2 \right] \frac{\partial^n}{\partial \lambda^n} y(x, \lambda) = 2\lambda n \frac{\partial^{n-1}}{\partial \lambda^{n-1}} y(x, \lambda) + n(n-1) \frac{\partial^{n-2}}{\partial \lambda^{n-2}} y(x, \lambda). \quad (4.9)$$

Writing (4.9) for $\psi^+(x, \lambda)$ and $e(x, \lambda)$, and using (4.8), we find

$$\left[-\frac{d^2}{dx^2} + q(x) - \lambda^2 \right] f_{n_0+1}(x, \lambda_p) = 0, \quad (4.10)$$

where

$$f_{n_0+1}(x, \lambda_p) = \left\{ \frac{\partial^{n_0+1}}{\partial \lambda^{n_0+1}} \psi^+(x, \lambda) \right\}_{\lambda=\lambda_p} - \sum_{m=0}^{n_0+1} A_m(\lambda_p) \left\{ \frac{\partial^m}{\partial \lambda^m} e(x, \lambda) \right\}_{\lambda=\lambda_p}. \quad (4.11)$$

From (4.1) we have

$$W[f_{n_0+1}(x, \lambda_p), e(x, \lambda_p)] = \left\{ \frac{d^{n_0+1}}{d\lambda^{n_0+1}} W[\psi^+(x, \lambda), e(x, \lambda)] \right\}_{\lambda=\lambda_p} = 0. \quad (4.12)$$

Hence there exists a constant $a_{n_0+1}(\lambda_p)$ such that

$$f_{n_0+1}(x, \lambda_p) = a_{n_0+1}(\lambda_p) e(x, \lambda_p). \quad (4.13)$$

This shows that (4.3) holds for $n = n_0 + 1$.

Similarly we can prove that (4.5) holds. \square

Definition 4.2. Let $\lambda = \lambda_0$ be an eigenvalue of L . If the functions

$$y_0(x, \lambda_0), y_1(x, \lambda_0), \dots, y_s(x, \lambda_0) \quad (4.14)$$

satisfy the equations

$$l(y_0) - \lambda_0 y_0 = 0, \quad l(y_j) - \lambda_0 y_j - y_{j-1} = 0, \quad j = 1, 2, \dots, s, \quad (4.15)$$

then the function $y_0(x, \lambda_0)$ is called the eigenfunction corresponding to the eigenvalue $\lambda = \lambda_0$ of L . The functions $y_1(x, \lambda_0), \dots, y_s(x, \lambda_0)$ are called the associated functions corresponding $\lambda = \lambda_0$. The eigenfunctions and the associated functions corresponding to $\lambda = \lambda_0$ are called the principal functions of the eigenvalue $\lambda = \lambda_0$.

The principal functions of the spectral singularities of L are defined similarly.

Now using (4.3) and (4.5) define the functions

$$U_{n,p}(x) = \left\{ \frac{\partial^n}{\partial \lambda^n} \psi^+(x, \lambda) \right\}_{\lambda=\lambda_p} = \sum_{m=0}^n A_m(\lambda_p) \left\{ \frac{\partial^m}{\partial \lambda^m} e(x, \lambda) \right\}_{\lambda=\lambda_p}, \quad (4.16)$$

$n = 0, 1, \dots, m_p - 1, \dots, p = 1, 2, \dots, j$
and

$$U_{n,p}(x) = \left\{ \frac{\partial^n}{\partial \lambda^n} \psi^-(x, \lambda) \right\}_{\lambda=\lambda_p} = \sum_{m=0}^n B_m(\lambda_p) \left\{ \frac{\partial^m}{\partial \lambda^m} e(x, -\lambda) \right\}_{\lambda=\lambda_p}, \quad (4.17)$$

$n = 0, 1, \dots, m_p - 1, p = j + 1, \dots, k$.

Then for $\lambda = \lambda_p, p = 1, 2, \dots, j, j + 1, \dots, k,$

$$\begin{aligned} l(U_{0,p}) &= 0, \\ l(U_{1,p}) + \frac{1}{1!} \frac{\partial}{\partial \lambda} l(U_{0,p}) &= 0, \\ l(U_{n,p}) + \frac{1}{1!} \frac{\partial}{\partial \lambda} l(U_{n-1,p}) + \frac{1}{2!} \frac{\partial^2}{\partial \lambda^2} l(U_{n-2,p}) &= 0, \end{aligned} \quad (4.18)$$

$n = 2, 3, \dots, m_p - 1,$

hold, where $l(u) = -u'' + q(x)u - \lambda^2 u$ and $(\partial^m / \partial \lambda^m)l(u)$ denotes the differential expressions whose coefficients are the m -th derivatives with respect to λ of the corresponding coefficients of the differential expression $l(u)$. Equation (4.18) shows that $U_{0,p}$ is the eigenfunction corresponding to the eigenvalue $\lambda = \lambda_p$; $U_{1,p}, U_{2,p}, \dots, U_{m_p-1,p}$ are the associated functions of $U_{0,p}$ ([25, 26]).

$U_{0,p}, U_{1,p}, \dots, U_{m_p-1,p}, p = 1, 2, \dots, j, j + 1, \dots, k$ are called the principal functions corresponding to the eigenvalue $\lambda = \lambda_p, p = 1, 2, \dots, j, j + 1, \dots, k$ of L .

Theorem 4.3. *One has*

$$U_{n,p} \in L^2(\mathbb{R}_+), \quad n = 0, 1, \dots, m_p - 1, \quad p = 1, 2, \dots, j, j + 1, \dots, k. \quad (4.19)$$

Proof. Let $0 \leq n \leq m_p - 1$ and $1 \leq p \leq j$. Using (2.6) and (3.7) we obtain that

$$|K(x, t)| \leq ce^{-\varepsilon\sqrt{(x+t)/2}}. \quad (4.20)$$

From (2.4) we get

$$\left| \left\{ \frac{\partial^n}{\partial \lambda^n} e(x, \lambda) \right\}_{\lambda=\lambda_p} \right| \leq x^n e^{-x \operatorname{Im} \lambda_p} + c \int_x^\infty t^n e^{-\varepsilon\sqrt{(x+t)/2}} e^{-t \operatorname{Im} \lambda_p} dt, \quad (4.21)$$

where $c > 0$ is a constant. Since $\operatorname{Im} \lambda_p > 0$ for the eigenvalues $\lambda_p, p = 1, \dots, j$, of L , (4.21) implies that

$$\left\{ \frac{\partial^n}{\partial \lambda^n} e(x, \lambda) \right\}_{\lambda=\lambda_p} \in L^2(\mathbb{R}_+), \quad n = 0, 1, \dots, m_p - 1, \quad p = 1, 2, \dots, j. \quad (4.22)$$

The proof of theorem is obtained from (4.16) and (4.22). In a similar way using (4.17) we may also prove the results for $0 \leq n \leq m_p - 1$ and $j + 1 \leq p \leq k$. \square

Let μ_1, \dots, μ_v , and μ_{v+1}, \dots, μ_l be the zeros of E^+ and E^- in $\mathbb{R}^* = \mathbb{R} \setminus \{0\}$ (which are the spectral singularities of L) with multiplicities n_1, \dots, n_v and n_{v+1}, \dots, n_l , respectively.

Similar to (4.3) and (4.5) we can show the following:

$$\left\{ \frac{\partial^n}{\partial \lambda^n} \psi(x, \lambda) \right\}_{\lambda=\mu_p} = \sum_{m=0}^n C_m(\mu_p) \left\{ \frac{\partial^m}{\partial \lambda^m} e(x, \lambda) \right\}_{\lambda=\mu_p}, \quad (4.23)$$

$n = 0, 1, \dots, n_p - 1, p = 1, 2, \dots, v,$
where

$$C_m(\mu_p) = - \binom{n}{m} \left\{ \frac{\partial^{n-m}}{\partial \lambda^{n-m}} E^-(\lambda) \right\}_{\lambda=\mu_p}, \quad (4.24)$$

$$\left\{ \frac{\partial^n}{\partial \lambda^n} \psi(x, \lambda) \right\}_{\lambda=\mu_p} = \sum_{m=0}^n D_m(\mu_p) \left\{ \frac{\partial^m}{\partial \lambda^m} e(x, -\lambda) \right\}_{\lambda=\mu_p},$$

$n = 0, 1, \dots, n_p - 1, p = v + 1, \dots, l,$
where

$$D_m(\mu_p) = \binom{n}{m} \left\{ \frac{\partial^{n-m}}{\partial \lambda^{n-m}} E^+(\lambda) \right\}_{\lambda=\mu_p}. \quad (4.25)$$

Now define the generalized eigenfunctions and generalized associated functions corresponding to the spectral singularities of L by the following:

$$v_{n,p}(x) = \left\{ \frac{\partial^n}{\partial \lambda^n} \psi(x, \lambda) \right\}_{\lambda=\mu_p} = \sum_{m=0}^n C_m(\mu_p) \left\{ \frac{\partial^m}{\partial \lambda^m} e(x, \lambda) \right\}_{\lambda=\mu_p}, \quad (4.26)$$

$n = 0, 1, \dots, n_p - 1, p = 1, 2, \dots, v,$

$$v_{n,p}(x) = \left\{ \frac{\partial^n}{\partial \lambda^n} \psi(x, \lambda) \right\}_{\lambda=\mu_p} = \sum_{m=0}^n D_m(\mu_p) \left\{ \frac{\partial^m}{\partial \lambda^m} e(x, -\lambda) \right\}_{\lambda=\mu_p}, \quad (4.27)$$

$n = 0, 1, \dots, n_p - 1, p = v + 1, \dots, l.$

Then $v_{n,p}, n = 0, 1, \dots, n_p - 1, p = 1, 2, \dots, v, v + 1, \dots, l,$ also satisfy the equations analogous to (4.18).

$v_{0,p}, v_{1,p}, \dots, v_{n_p-1,p}, p = 1, 2, \dots, v, v + 1, \dots, l$ are called the principal functions corresponding to the spectral singularities $\lambda = \mu_p, p = 1, 2, \dots, v, v + 1, \dots, l$ of L .

Theorem 4.4. *One has*

$$v_{n,p} \notin L^2(\mathbb{R}_+), \quad n = 0, 1, \dots, n_p - 1, p = 1, 2, \dots, v, v + 1, \dots, l. \quad (4.28)$$

Proof. If we consider (4.21) for the principal functions corresponding to the spectral singularities $\lambda = \mu_p, p = 1, 2, \dots, v, v + 1, \dots, l,$ of L and consider that $\text{Im } \lambda_p = 0$ for the spectral singularities, then we have (4.28), by (4.26) and (4.27). \square

Now introduce the Hilbert spaces

$$\begin{aligned} H_n &= \left\{ f : \int_0^\infty (1+x)^{2n} |f(x)|^2 dx < \infty \right\}, \quad n = 1, 2, \dots, \\ H_{-n} &= \left\{ g : \int_0^\infty (1+x)^{-2n} |g(x)|^2 dx < \infty \right\}, \quad n = 1, 2, \dots, \end{aligned} \quad (4.29)$$

with

$$\|f\|_n^2 = \int_0^\infty (1+x)^{2n} |f(x)|^2 dx; \quad \|g\|_{-n}^2 = \int_0^\infty (1+x)^{-2n} |g(x)|^2 dx, \quad (4.30)$$

respectively. Then

$$H_{n+1} \subsetneq H_n \subsetneq L^2(\mathbb{R}_+) \subsetneq H_{-n} \subsetneq H_{-(n+1)}, \quad n = 1, 2, \dots, \quad (4.31)$$

and H_{-n} is isomorphic to the dual of H_n .

Theorem 4.5. *One has*

$$v_{n,p} \in H_{-(n+1)}, \quad n = 0, 1, \dots, n_p - 1, \quad p = 1, 2, \dots, v, v+1, \dots, l. \quad (4.32)$$

Proof. From (2.4) we have

$$\begin{aligned} \int_0^\infty (1+x)^{-2(n+1)} \left| (ix)^n e^{i\mu_p x} \right|^2 dx &< \infty, \\ \int_0^\infty (1+x)^{-2(n+1)} \left| \int_x^\infty (it)^n K(x,t) e^{i\mu_p t} dt \right|^2 dx &< \infty. \end{aligned} \quad (4.33)$$

Using (4.26), (4.33) we obtain

$$v_{n,p} \in H_{-(n+1)}, \quad n = 0, 1, \dots, n_p - 1, \quad p = 1, 2, \dots, v. \quad (4.34)$$

In a similar way, we find

$$v_{n,p} \in H_{-(n+1)}, \quad n = 0, 1, \dots, n_p - 1, \quad p = v+1, \dots, l. \quad (4.35)$$

□

Let us choose n_0 so that

$$n_0 = \max\{n_1, n_2, \dots, n_v, n_{v+1}, \dots, n_l\}. \quad (4.36)$$

By Theorem 4.5 and (4.31) we get following theorem

Theorem 4.6. *One has*

$$v_{n,p} \in H_{-n_0}, \quad n = 0, 1, \dots, n_p - 1, \quad p = 1, 2, \dots, v, v+1, \dots, l. \quad (4.37)$$

Acknowledgment

The author would like to thank Professor E. Bairamov for his helpful suggestions during the preparation of this work.

References

- [1] M. A. Naïmark, "Investigation of the spectrum and the expansion in eigenfunctions of a non-selfadjoint differential operator of the second order on a semi-axis," *American Mathematical Society Translations*, vol. 16, pp. 103–193, 1960.
- [2] V. E. Lyance, "A differential operator with spectral singularities I," *American Mathematical Society Transactions Series 2*, vol. 60, pp. 185–225, 1967.
- [3] V. E. Lyance, "A differential operator with spectral singularities II," *American Mathematical Society Transactions Series 2*, vol. 60, pp. 227–283, 1967.
- [4] A. M. Krall, "The adjoint of a differential operator with integral boundary conditions," *Proceedings of the American Mathematical Society*, vol. 16, pp. 738–742, 1965.
- [5] A. M. Krall, "A nonhomogeneous eigenfunction expansion," *Transactions of the American Mathematical Society*, vol. 117, pp. 352–361, 1965.
- [6] A. M. Krall, "Second order ordinary differential operators with general boundary conditions," *Duke Mathematical Journal*, vol. 32, pp. 617–625, 1965.
- [7] A. M. Krall, "Nonhomogeneous differential operators," *The Michigan Mathematical Journal*, vol. 12, pp. 247–255, 1965.
- [8] A. M. Krall, "On non-self-adjoint ordinary differential operators of the second order," *Doklady Akademii Nauk SSSR*, vol. 165, pp. 1235–1237, 1965.
- [9] M. G. Gasymov, "Expansion in terms of the solutions of a scattering theory problem for the non-selfadjoint Schrödinger equation," *Soviet Mathematics—Doklady*, vol. 9, pp. 390–393, 1968.
- [10] M. G. Gasymov and F. G. Maksudov, "The principal part of the resolvent of nonselfadjoint operators in the neighborhood of spectral singularities," *Functional Analysis and Its Applications*, vol. 6, no. 3, pp. 185–192, 1972.
- [11] M. Adivar and E. Bairamov, "Spectral properties of non-selfadjoint difference operators," *Journal of Mathematical Analysis and Applications*, vol. 261, no. 2, pp. 461–478, 2001.
- [12] M. Adivar and E. Bairamov, "Difference equations of second order with spectral singularities," *Journal of Mathematical Analysis and Applications*, vol. 277, no. 2, pp. 714–721, 2003.
- [13] E. Bairamov, Ö. Çakar, and A. M. Krall, "An eigenfunction expansion for a quadratic pencil of a Schrödinger operator with spectral singularities," *Journal of Differential Equations*, vol. 151, no. 2, pp. 268–289, 1999.
- [14] E. Bairamov, Ö. Çakar, and A. M. Krall, "Non-selfadjoint difference operators and Jacobi matrices with spectral singularities," *Mathematische Nachrichten*, vol. 229, pp. 5–14, 2001.
- [15] E. Bairamov, Ö. Çakar, and A. M. Krall, "Spectral properties, including spectral singularities, of a quadratic pencil of Schrödinger operators on the whole real axis," *Quaestiones Mathematicae*, vol. 26, no. 1, pp. 15–30, 2003.
- [16] E. Bairamov, Ö. Çakar, and C. Yanik, "Spectral singularities of the Klein-Gordon s -wave equation," *Indian Journal of Pure and Applied Mathematics*, vol. 32, no. 6, pp. 851–857, 2001.
- [17] E. Bairamov and A. O. Çelebi, "Spectrum and spectral expansion for the non-selfadjoint discrete Dirac operators," *The Quarterly Journal of Mathematics. Oxford. Second Series*, vol. 50, no. 200, pp. 371–384, 1999.
- [18] E. Bairamov and Ö. Karaman, "Spectral singularities of Klein-Gordon s -wave equations with an integral boundary condition," *Acta Mathematica Hungarica*, vol. 97, no. 1-2, pp. 121–131, 2002.
- [19] A. M. Krall, E. Bairamov, and Ö. Çakar, "Spectrum and spectral singularities of a quadratic pencil of a Schrödinger operator with a general boundary condition," *Journal of Differential Equations*, vol. 151, no. 2, pp. 252–267, 1999.
- [20] A. M. Krall, E. Bairamov, and Ö. Çakar, "Spectral analysis of non-selfadjoint discrete Schrödinger operators with spectral singularities," *Mathematische Nachrichten*, vol. 231, pp. 89–104, 2001.
- [21] V. A. Marchenko, "Expansion in eigenfunctions of non-selfadjoint singular second order differential operators," *American Mathematical Society Transactions Series 2*, vol. 25, pp. 77–130, 1963.

- [22] V. A. Marchenko, *Sturm-Liouville Operators and Applications*, vol. 22, Operator Theory: Advances and Applications, Birkhäuser, Basel, Switzerland, 1986.
- [23] M. Jaulent and C. Jean, "The inverse s -wave scattering problem for a class of potentials depending on energy," *Communications in Mathematical Physics*, vol. 28, pp. 177–220, 1972.
- [24] E. Bairamov and N. Yokus, "Spectral singularities of Sturm-Liouville problems with eigenvalue-dependent boundary conditions," *Abstract and Applied Analysis*, p. Art. ID 289596, 8, 2009.
- [25] M. V. Keldysh, "On the completeness of the eigenfunctions of some classes of non-selfadjoint linear operators," *Soviet Mathematics—Doklady*, vol. 77, no. 4, pp. 11–14, 1951.
- [26] M. V. Keldysh, "On the completeness of the eigenfunctions of some classes of non-selfadjoint linear operators," *Russian Mathematical Surveys*, vol. 26, no. 4, pp. 15–41, 1971.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

