

# ON EQUIVALENCE OF COEFFICIENT CONDITIONS

**L. LEINDLER**

Bolyai Institute, University of Szeged,  
Aradi vértanúk tere 1,  
H-6720 Szeged, Hungary  
EMail: [leindler@math.u-szeged.hu](mailto:leindler@math.u-szeged.hu)

*Received:* 24 January, 2007

*Accepted:* 12 March, 2007

*Communicated by:* **J.M. Rassias**

*2000 AMS Sub. Class.:* 26D15, 40D15.

*Key words:* Numerical series, Equiconvergence, Weyl multiplier.

*Abstract:* Two equivalence theorems and two corollaries are proved pertaining to the equiconvergence of numerical series.



**Equivalence of Coefficient  
Conditions**

L. Leindler

vol. 8, iss. 1, art. 8, 2007

[Title Page](#)

[Contents](#)



Page 1 of 12

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b>Results</b>	<b>6</b>
<b>3</b>	<b>Proofs</b>	<b>8</b>



---

## Equivalence of Coefficient Conditions

L. Leindler

vol. 8, iss. 1, art. 8, 2007

---

Title Page

Contents



Page 2 of 12


Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756



Title Page

Contents



Page 3 of 12

Go Back

Full Screen

Close

## 1. Introduction

Some papers, see [1], [2] and [5], have dealt with the equivalence of coefficient conditions, e.g. in [3] it was proved that two conditions which guarantee that a factor-sequence should be a Weyl multiplier for a certain property of a given orthogonal series is equivalent to one assumption. An example of these results is the following general theorem proved in [2].

**Theorem 1.1.** *Let  $0 < p < q$ ,  $\{\lambda_n\}$  and  $\{c_n\}$  be sequences of nonnegative numbers, furthermore let  $\Lambda_n := \sum_{k=1}^n \lambda_k$ . The inequality*

$$(1.1) \quad S_{11} := \sum_{n=1}^{\infty} \lambda_n \left( \sum_{k=n}^{\infty} c_k^q \right)^{\frac{p}{q}} < \infty$$

*holds if and only if there exists a nondecreasing sequence  $\{\mu_n\}$  of positive numbers satisfying the following conditions*

$$(1.2) \quad S_{12} := \sum_{n=1}^{\infty} c_n^q \mu_n < \infty$$

*and*

$$(1.3) \quad S_{13} := \sum_{n=1}^{\infty} \lambda_n \left( \frac{\Lambda_n}{\mu_n} \right)^{\frac{p}{q-p}} < \infty.$$

In the special case  $p = 1$ ,  $q = 2$ , with  $a_n$  and  $\lambda_n^{-1}$  in the place of  $c_n$  and  $\lambda_n$ , the author in [3] showed that if

$$(1.4) \quad \mu_n := \Lambda_n C_n^{-1}, \quad \text{where} \quad C_n := \left( \sum_{k=n}^{\infty} c_k^2 \right)^{\frac{1}{2}},$$



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 4 of 12

[Go Back](#)

[Full Screen](#)

[Close](#)

then (1.1) implies (1.2) and (1.3).

In this special case it is easy to see that if  $\varepsilon_n \rightarrow 0$ , then with  $\mu_n^* := \varepsilon_n \mu_n$  in place of  $\mu_n$ , the condition (1.2) is also satisfied, but it can be that

$$(1.5) \quad \sum_{n=1}^{\infty} \lambda_n \frac{\Lambda_n}{\mu_n^*} = \infty$$

will occur.

This raises the question: Do (1.2) and (1.5) with the sequence  $\{\mu_n^*\}$  also imply (1.1) for arbitrary  $\{c_n\}$ ? In [3] we showed that the answer is negative. In other words, this verified the necessity of condition (1.3).

This shows that condition (1.2), in general, does not imply the inequality (1.3).

Thus, we can ask, promptly in connection with the general case considered in Theorem 1.1: What is the "optimal sequence  $\{\mu_n\}$ " when (1.2) implies (1.3) and conversely?

We shall show that the optimal sequence is

$$(1.6) \quad \mu_n := \Lambda_n C_n^{p-q}, \quad \text{where} \quad C_n := \left( \sum_{k=n}^{\infty} c_k^q \right)^{\frac{1}{q}},$$

and with this  $\{\mu_n\}$  (1.2) holds if and only if (1.3) also holds, that is, the assumptions (1.2) and (1.3) are equivalent.

Since the following symmetrical analogue of Theorem 1.1 was also verified in [2], therefore we shall set the same question pertaining to the series appearing in it.

**Theorem 1.2.** *If  $p, q, \{\lambda_n\}$  and  $\{c_n\}$  are as in Theorem 1.1, furthermore  $\tilde{\Lambda}_n := \sum_{k=n}^{\infty} \lambda_k$ , then*

$$(1.7) \quad S_{17} := \sum_{m=1}^{\infty} \lambda_m \left( \sum_{k=1}^m c_k^q \right)^{\frac{p}{q}} < \infty$$

holds if and only if there exists a nondecreasing sequence  $\{\mu_n\}$  of positive numbers satisfying conditions (1.2) and

$$(1.8) \quad S_{18} := \sum_{n=1}^{\infty} \lambda_n \left( \frac{\tilde{\Lambda}_n}{\mu_n} \right)^{\frac{p}{q-p}} < \infty.$$

In order to verify our assertions made above, first we shall prove two theorems regarding the equiconvergence of two special series.



**Equivalence of Coefficient  
Conditions**

L. Leindler

vol. 8, iss. 1, art. 8, 2007

Title Page

Contents



Page 5 of 12


Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756



## 2. Results

We prove the following assertions.

**Theorem 2.1.** *Let  $0 < \alpha < 1$ ,  $\{a_n\}$  and  $\{\lambda_n\}$  be sequences of nonnegative numbers, furthermore let  $\Lambda_n := \sum_{k=1}^n \lambda_k$ ,  $A_n := \sum_{k=n}^{\infty} a_k$  and  $\mu_n := \Lambda_n A_n^{\alpha-1}$ . Then the sum*

$$(2.1) \quad S_{21} := \sum_{n=1}^{\infty} a_n \mu_n < \infty$$

*if and only if*

$$(2.2) \quad S_{22} := \sum_{n=1}^{\infty} \lambda_n A_n^{\alpha} < \infty.$$

**Theorem 2.2.** *If  $\alpha$ ,  $\{a_n\}$  and  $\{\lambda_n\}$  are as in Theorem 2.1, furthermore  $\tilde{\Lambda}_n := \sum_{k=n}^{\infty} \lambda_k$ ,  $\tilde{A}_n := \sum_{k=1}^n a_k$ ,  $\tilde{A}_0 := 0$ , and  $\tilde{\mu}_n := \tilde{\Lambda}_n \tilde{A}_n^{\alpha-1}$ , then*

$$(2.3) \quad S_{23} := \sum_{n=1}^{\infty} a_n \tilde{\mu}_n < \infty$$

*if and only if*

$$(2.4) \quad S_{24} := \sum_{n=1}^{\infty} \lambda_n \tilde{A}_n^{\alpha} < \infty.$$

**Corollary 2.3.** *If  $p, q, \{\lambda_n\}, \{c_n\}$  and  $\Lambda_n$  are as in Theorem 1.1, and  $\mu_n$  is defined in (1.6), then the sums in (1.1), (1.2) and (1.3) are equiconvergent.*



Title Page

Contents



Page 7 of 12

Go Back

Full Screen

Close

**Corollary 2.4.** If  $p, q, \{\lambda_n\}, \{c_n\}$  and  $\tilde{\Lambda}_n$  are as in Theorem 1.2, furthermore


$$\mu_n := \tilde{\Lambda}_n \tilde{C}_n^{p-q}, \quad \text{where} \quad \tilde{C}_n := \left( \sum_{k=1}^n c_k^q \right)^{\frac{1}{q}},$$

then the sums in (1.2), (1.7) and (1.8) are equiconvergent.

*Remark 1.* Corollary 2.3 shows that if (1.1) implies a certain property of a fixed orthogonal series  $\sum_{n=1}^{\infty} c_n \varphi_n(x)$ , then there is no exact universal Weyl multiplier concerning this property, namely the multiplier sequence  $\{\mu_n\}$  depends on  $\{c_n\}$ .

*Remark 2.* The interested reader can check that the proofs of the implications (1.1) $\Rightarrow$ (1.2) and (1.7) $\Rightarrow$ (1.2) given by our corollaries are shorter than those in [2].

*Remark 3.* As far as we know, Y. Okuyama and T. Tsuchikara [4] were the first to study conditions of the type (1.7).



Title Page

Contents



Page 8 of 12

Go Back

Full Screen

Close

### 3. Proofs

*Proof of Theorem 2.1.* First we show that (2.1) implies (2.2). Since  $A_n \rightarrow 0$ , then

$$(3.1) \quad \begin{aligned} \sum_{n=1}^{\infty} \lambda_n A_n^\alpha &= \sum_{n=1}^{\infty} \lambda_n \sum_{m=n}^{\infty} (A_m^\alpha - A_{m+1}^\alpha) \\ &= \sum_{m=1}^{\infty} (A_m^\alpha - A_{m+1}^\alpha) \sum_{n=1}^m \lambda_n. \end{aligned}$$

An easy consideration yields that if  $0 \leq a < b$ ,  $0 < \alpha < 1$  and

$$(3.2) \quad \frac{b^\alpha - a^\alpha}{b - a} = \alpha \xi^{\alpha-1},$$

then

$$(3.3) \quad \xi \geq \alpha^{1/(1-\alpha)} b =: \xi_0,$$

namely if  $a = 0$  then  $\xi = \xi_0$ .

Using the relations (3.2) and (3.3) we obtain that

$$(3.4) \quad A_m^\alpha - A_{m+1}^\alpha = a_m \alpha \xi^{\alpha-1} \leq a_m A_m^{\alpha-1}.$$

This and (3.1) yield that


$$\sum_{n=1}^{\infty} \lambda_n A_n^\alpha \leq \sum_{m=1}^{\infty} a_m A_m^{\alpha-1} \Lambda_m = \sum_{m=1}^{\infty} a_m \mu_m.$$

Herewith the implication (2.1) $\Rightarrow$ (2.2) is proved.



Title Page

Contents



Page 9 of 12

Go Back

Full Screen

Close

The proof of (2.2) $\Rightarrow$ (2.1) is similar. We use the first part of (3.4),  $\alpha < 1$  and  $\xi < A_m$ . Thus

$$\begin{aligned}\sum_{m=1}^{\infty} a_m \mu_m &= \sum_{m=1}^{\infty} a_m A_m^{\alpha-1} \Lambda_m \\ &= \sum_{m=1}^{\infty} (A_m - A_{m+1}) A_m^{\alpha-1} \Lambda_m \\ &\leq \alpha^{-1} \sum_{m=1}^{\infty} (A_m^{\alpha} - A_{m+1}^{\alpha}) \sum_{n=1}^m \lambda_n \\ &= \alpha^{-1} \sum_{n=1}^{\infty} \lambda_n \sum_{m=n}^{\infty} (A_m^{\alpha} - A_{m+1}^{\alpha}) \\ &= \alpha^{-1} \sum_{n=1}^{\infty} \lambda_n A_n^{\alpha},\end{aligned}$$

that is, (2.2) $\Rightarrow$ (2.1) is verified.

The proof of Theorem 2.1 is complete. ■

*Proof of Theorem 2.2.* The proof is almost the same as that of Theorem 2.1. By (3.2) and (3.3) we get that

$$(3.5) \quad \tilde{A}_m^{\alpha} - \tilde{A}_{m-1}^{\alpha} = a_m \alpha \xi^{\alpha-1} \leq a_m \tilde{A}_m^{\alpha-1}.$$

Utilizing this at the final step we have

$$\sum_{n=1}^{\infty} \lambda_n \tilde{A}_n^{\alpha} = \sum_{n=1}^{\infty} \lambda_n \sum_{k=1}^n (\tilde{A}_k^{\alpha} - \tilde{A}_{k-1}^{\alpha})$$

$$= \sum_{k=1}^{\infty} (\tilde{A}_k^\alpha - \tilde{A}_{k-1}^\alpha) \sum_{n=k}^{\infty} \lambda_n \leq \sum_{k=1}^{\infty} a_k \tilde{A}_k^{\alpha-1} \tilde{\Lambda}_k \equiv \sum_{k=1}^{\infty} a_k \tilde{\mu}_k;$$

this proves the implication (2.3) $\Rightarrow$ (2.4).

To verify (2.4) $\Rightarrow$ (2.3) we use

$$\alpha^{-1}(\tilde{A}_m^\alpha - \tilde{A}_{m-1}^\alpha) \geq a_m \tilde{A}_m^{\alpha-1},$$

which follows from the first part of (3.5) by  $\alpha < 1$  and  $\xi < \tilde{A}_m$ .

Thus we get that

$$\begin{aligned} \sum_{n=1}^{\infty} a_n \tilde{\mu}_n &= \sum_{n=1}^{\infty} a_n \tilde{A}_n^{\alpha-1} \tilde{\Lambda}_n \\ &\leq \alpha^{-1} \sum_{n=1}^{\infty} (\tilde{A}_n^\alpha - \tilde{A}_{n-1}^\alpha) \sum_{k=n}^{\infty} \lambda_k \\ &= \alpha^{-1} \sum_{k=1}^{\infty} \lambda_k \sum_{n=1}^k (\tilde{A}_n^\alpha - \tilde{A}_{n-1}^\alpha) \\ &= \alpha^{-1} \sum_{k=1}^{\infty} \lambda_k \tilde{A}_k^\alpha. \end{aligned}$$

Summing up, the proof of Theorem 2.2 is complete. ■

*Proof of Corollary 2.3.* We shall use the results of Theorem 2.1 with  $\alpha = \frac{p}{q}$  and  $a_n = c_n^q$ . Then  $\mu_n = \Lambda_n \left( \sum_{k=n}^{\infty} c_k^q \right)^{\frac{p-q}{q}}$ , and

$$S_{22} \equiv S_{11} \equiv S_{13} \quad \text{as well as} \quad S_{21} \equiv S_{12},$$



**Equivalence of Coefficient  
Conditions**

L. Leindler

vol. 8, iss. 1, art. 8, 2007

Title Page

Contents



Page 10 of 12


Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-575b



moreover, by Theorem 2.1,  $S_{21}$  and  $S_{22}$  are equiconvergent, herewith Corollary 2.3 is proved. ■

*Proof of Corollary 2.4.* Now we utilize Theorem 2.2 with  $\alpha = \frac{p}{q}$  and  $a_n = c_n^q$ . Then

$$\tilde{\mu}_n = \tilde{\Lambda}_n \left( \sum_{k=1}^n c_k^q \right)^{\frac{p-q}{q}}, \text{ furthermore}$$

$$S_{24} \equiv S_{17} \equiv S_{18} \quad \text{and} \quad S_{23} \equiv S_{12},$$

hold. Since, by Theorem 2.2,  $S_{23}$  and  $S_{24}$  are equiconvergent, thus Corollary 2.4 is verified. ■

Title Page

Contents



Page 11 of 12

Go Back

Full Screen

Close

## References

- [1] L. LEINDLER, On relations of coefficient-conditions, *Acta Math. Hungar.*, **39** (1982), 409–420.
- [2] L. LEINDLER, On equivalence of coefficient conditions and application, *Math. Inequal. Appl.*, **1** (1998), 41–51.
- [3] L. LEINDLER, An addendum of an equivalence theorem, *Math. Inequal. Appl.*, **10**(1) (2007), 49–55.
- [4] Y. OKUYAMA AND T. TSUCHIKURA, On the absolute Riesz summability of orthogonal series, *Analysis Math.*, **7** (1981), 199–208.
- [5] V. TOTIK AND I. VINCZE, On relations of coefficient conditions, *Acta Sci. Math. (Szeged)*, **50** (1986), 93–98.



---

Equivalence of Coefficient  
Conditions

L. Leindler

vol. 8, iss. 1, art. 8, 2007

---

Title Page

Contents



Page 12 of 12

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756