



SUBORDINATION THEOREM FOR A FAMILY OF ANALYTIC FUNCTIONS ASSOCIATED WITH THE CONVOLUTION STRUCTURE

J. K. PRAJAPAT

Department of Mathematics
Bhartiya Institute of Engineering & Technology
Near Sanwali Circle, Bikaner By-Pass Road
Sikar-332001, Rajasthan, INDIA.
EMail: jkp_0007@rediffmail.com

Received: 04 May, 2007

Accepted: 01 September, 2008

Communicated by: **S.S. Dragomir**

2000 AMS Sub. Class.: 30C45.

Key words: Analytic function, Hadamard product(or convolution), Dziok-Srivastava linear operator, Subordination factor sequence, Characterization properties.

Abstract: We use the familiar convolution structure of analytic functions to introduce new class of analytic functions of complex order. The results investigated in the present paper include, the characterization and subordination properties for this class of analytic functions. Several interesting consequences of our results are also pointed out.

Acknowledgements: The author expresses his sincerest thanks to the worthy referee for valuable comments. He is also thankful to Emeritus Professor Dr. R.K. Raina for his useful suggestions.


Subordination Theorem for a Family of Analytic Functions

J. K. Prajapat

vol. 9, iss. 4, art. 102, 2008

[Title Page](#)

[Contents](#)



Page 1 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction and Preliminaries	3
2	Characterization Properties	7
3	Subordination Theorem	10



**Subordination Theorem for a
Family of Analytic Functions**

J. K. Prajapat

vol. 9, iss. 4, art. 102, 2008

Title Page

Contents



Page 2 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 3 of 16

Go Back

Full Screen

Close

1. Introduction and Preliminaries

Let \mathcal{A} denote the class of functions of the form

$$(1.1) \quad f(z) = z + \sum_{k=2}^{\infty} a_k z^k,$$

which are analytic and univalent in the open unit disk $\mathbb{U} = \{z; z \in \mathbb{C} : |z| < 1\}$. If $f \in \mathcal{A}$ is given by (1.1) and $g \in \mathcal{A}$ is given by

$$(1.2) \quad g(z) = z + \sum_{k=2}^{\infty} b_k z^k,$$

then the Hadamard product (or convolution) $f * g$ of f and g is defined (as usual) by

$$(1.3) \quad (f * g)(z) := z + \sum_{k=2}^{\infty} a_k b_k z^k.$$

In this article we study the class $\mathcal{S}_\gamma(g; \alpha)$ introduced in the following:

Definition 1.1. For a given function $g(z) \in \mathcal{A}$ defined by (1.2), where $b_k \geq 0$ ($k \geq 2$). We say that $f(z) \in \mathcal{A}$ is in $\mathcal{S}_\gamma(g; \alpha)$, provided that $(f * g)(z) \neq 0$, and

$$(1.4) \quad \operatorname{Re} \left\{ 1 + \frac{1}{\gamma} \left(\frac{z(f * g)'(z)}{(f * g)(z)} - 1 \right) \right\} > \alpha \quad (z \in \mathbb{U}; \gamma \in \mathbb{C} \setminus \{0\}; 0 \leq \alpha < 1).$$

Note that

$$\mathcal{S}_1 \left(\frac{z}{1-z}; \alpha \right) = \mathcal{S}^*(\alpha) \quad \text{and} \quad \mathcal{S}_1 \left(\frac{z}{(1-z)^2}; \alpha \right) = \mathcal{K}(\alpha),$$

are, respectively, the familiar classes of starlike and convex functions of order α in \mathbb{U} (see, for example, [11]). Also

$$\mathcal{S}_\gamma \left(\frac{z}{1-z}; 0 \right) = \mathcal{S}_\gamma^* \quad \text{and} \quad \mathcal{S}_\gamma \left(\frac{z}{(1-z)^2}; 0 \right) = \mathcal{K}_\gamma,$$

where the classes \mathcal{S}_γ^* and \mathcal{K}_γ stem essentially from the classes of starlike and convex functions of complex order, which were considered earlier by Nasr and Aouf [9] and Wiatrowski [12], respectively (see also [7] and [8]).

Remark 1. When

$$(1.5) \quad g(z) = z + \sum_{k=2}^{\infty} \frac{(\alpha_1)_{k-1} \cdots (\alpha_q)_{k-1}}{(\beta_1)_{k-1} \cdots (\beta_s)_{k-1} (k-1)!} z^k$$

$$(\alpha_j \in \mathbb{C} (j = 1, 2, \dots, q), \quad \beta_j \in \mathbb{C} \setminus \{0, -1, -2, \dots\} \quad (j = 1, 2, \dots, s)),$$

with the parameters

$$\alpha_1, \dots, \alpha_q \quad \text{and} \quad \beta_1, \dots, \beta_s,$$

being so chosen that the coefficients b_k in (1.2) satisfy the following condition:

$$(1.6) \quad b_k = \frac{(\alpha_1)_{k-1} \cdots (\alpha_q)_{k-1}}{(\beta_1)_{k-1} \cdots (\beta_s)_{k-1} (k-1)!} \geq 0,$$

then the class $\mathcal{S}_\gamma(g; \alpha)$ is transformed into a (presumably) new class $\mathcal{S}_\gamma^*(q, s, \alpha)$ defined by

$$(1.7) \quad \mathcal{S}_\gamma^*(q, s, \alpha) := \left\{ f : f \in \mathcal{A} \text{ and } \operatorname{Re} \left(1 + \frac{1}{\gamma} \left(\frac{z(H_s^q[\alpha_1]f)'(z)}{(H_s^q[\alpha_1]f)(z)} - 1 \right) \right) > \alpha \right\}$$

$$(z \in \mathbb{U}; q \leq s + 1; q, s \in \mathbb{N}_0; \gamma \in \mathbb{C} \setminus \{0\}).$$





[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 5 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

The operator

$$(H_s^q[\alpha_1]f)(z) := H_s^q(\alpha_1, \dots, \alpha_q; \beta_1, \dots, \beta_s)f(z),$$

involved in (1.7) is the Dziok-Srivastava linear operator (see for details, [3]) which contains such well known operators as the Hohlov linear operator, Carlson-Shaffer linear operator, Ruscheweyh derivative operator, the Barnardi-Libera-Livingston operator, and the Srivastava-Owa fractional derivative operator. One may refer to the papers [3] to [5] for further details and references for these operators. The Dziok-Srivastava linear operator defined in [3] was further extended by Dziok and Raina [1] (see also [2]).

In our present investigation, we require the following definitions and a related result due to Welf [13].

Definition 1.2 (Subordination Principal). For two functions f and g analytic in \mathbb{U} , we say that the function $f(z)$ is subordinated to $g(z)$ in \mathbb{U} and write $f(z) \prec g(z)$ ($z \in \mathbb{U}$), if there exists a Schwarz function $w(z)$ analytic in \mathbb{U} with $w(0) = 0$, and $|w(z)| < 1$, such that $f(z) = g(w(z))$, $z \in \mathbb{U}$. In particular, if the function $g(z)$ is univalent in \mathbb{U} , the above subordination is equivalent to $f(0) = g(0)$ and $f(\mathbb{U}) \subset g(\mathbb{U})$.

Definition 1.3 (Subordinating Factor Sequence). A sequence $\{b_k\}_{k=1}^\infty$ of complex numbers is called a subordinating factor sequence if, whenever

$$f(z) = \sum_{k=1}^\infty a_k z^k \quad (a_1 = 1),$$

is analytic, univalent and convex in \mathbb{U} , we have the subordination given by

$$(1.8) \quad \sum_{k=1}^\infty a_k b_k z^k \prec f(z) \quad (z \in \mathbb{U}).$$

Lemma 1.4 (Wilf, [13]). *The sequence $\{b_k\}_{k=1}^{\infty}$ is a subordinating factor sequence if and only if*

$$(1.9) \quad \operatorname{Re} \left\{ 1 + 2 \sum_{k=1}^{\infty} b_k z^k \right\} > 0 \quad (z \in \mathbb{U}).$$



Subordination Theorem for a Family of Analytic Functions

J. K. Prajapat

vol. 9, iss. 4, art. 102, 2008

Title Page

Contents



Page 6 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



2. Characterization Properties

In this section we establish two results (Theorem 2.1 and Theorem 2.3) which give the sufficiency conditions for a function $f(z)$ defined by (1.1) and belong to the class $f(z) \in \mathcal{S}_\gamma(g; \alpha)$.

Theorem 2.1. Let $f(z) \in \mathcal{A}$ such that

$$(2.1) \quad \left| \frac{z(f * g)'(z)}{(f * g)(z)} - 1 \right| < 1 - \beta \quad (\beta < 1; z \in \mathbb{U}),$$

then $f(z) \in \mathcal{S}_\gamma(g; \alpha)$, provided that

$$(2.2) \quad |\gamma| \geq \frac{1 - \beta}{1 - \alpha}, \quad (0 \leq \alpha < 1).$$

Proof. In view of (2.1), we write

$$\frac{z(f * g)'(z)}{(f * g)(z)} = 1 + (1 - \beta)w(z) \quad \text{where } |w(z)| < 1 \text{ for } z \in \mathbb{U}.$$

Now

$$\begin{aligned} \operatorname{Re} \left\{ 1 + \frac{1}{\gamma} \left(\frac{z(f * g)'(z)}{(f * g)(z)} - 1 \right) \right\} &= \operatorname{Re} \left\{ 1 + \frac{1}{\gamma} (1 - \beta)w(z) \right\} \\ &= 1 + (1 - \beta) \operatorname{Re} \left\{ \frac{w(z)}{\gamma} \right\} \\ &\geq 1 - (1 - \beta) \left| \frac{w(z)}{\gamma} \right| \\ &> 1 - (1 - \beta) \cdot \frac{1}{|\gamma|} \geq \alpha, \end{aligned}$$

provided that $|\gamma| \geq \frac{1 - \beta}{1 - \alpha}$. This completes the proof. \square



Title Page

Contents



Page 8 of 16

Go Back

Full Screen

Close

If we set

$$\beta = 1 - (1 - \alpha)|\gamma| \quad (0 \leq \alpha < 1; \gamma \in \mathbb{C} \setminus \{0\}),$$

in Theorem 2.1, we obtain

Corollary 2.2. *If $f(z) \in \mathcal{A}$ such that*

$$(2.3) \quad \left| \frac{z(f * g)'(z)}{(f * g)(z)} - 1 \right| < (1 - \alpha)|\gamma| \quad (z \in \mathbb{U}, 0 \leq \alpha < 1; \gamma \in \mathbb{C} \setminus \{0\}),$$

then $f(z) \in \mathcal{S}_\gamma(g; \alpha)$.

Theorem 2.3. *Let $f(z) \in \mathcal{A}$ satisfy the following inequality*

$$(2.4) \quad \sum_{k=2}^{\infty} b_k [(k - 1) + (1 - \alpha)|\gamma|] |a_k| \leq (1 - \alpha)|\gamma|$$

$$(z \in \mathbb{U}; b_k \geq 0 (k \geq 2); \gamma \in \mathbb{C} \setminus \{0\}; 0 \leq \alpha < 1),$$

then $f(z) \in \mathcal{S}_\gamma(g; \alpha)$.

Proof. Suppose the inequality (2.4) holds true. Then in view of Corollary 2.2, we have

$$\begin{aligned} & |z(f * g)'(z) - (f * g)(z)| - (1 - \alpha)|\gamma| |(f * g)(z)| \\ &= \left| \sum_{k=2}^{\infty} b_k (k - 1) a_k z^k \right| - (1 - \alpha)|\gamma| \left| z + \sum_{k=2}^{\infty} b_k a_k z^k \right| \\ &\leq \left\{ \sum_{k=2}^{\infty} b_k (k - 1) |a_k| - (1 - \alpha)|\gamma| + (1 - \alpha)|\gamma| \sum_{k=2}^{\infty} b_k |a_k| \right\} |z| \end{aligned}$$

$$\leq \left\{ \sum_{k=2}^{\infty} b_k [(k-1) + (1-\alpha)|\gamma|] |a_k| - (1-\alpha)|\gamma| \right\} \leq 0.$$

This completes the proof. □

On specializing the parameters, Theorem 2.1 would yield the following results:

Corollary 2.4. *Let $f(z) \in \mathcal{A}$ satisfy the following inequality*

$$(2.5) \quad \sum_{k=2}^{\infty} (k + |\gamma| - 1) |a_k| \leq |\gamma| \quad (z \in \mathbb{U}, \gamma \in \mathbb{C} \setminus \{0\}),$$

then $f(z) \in \mathcal{S}_{\gamma}^*$.

Corollary 2.5. *Let $f(z) \in \mathcal{A}$ satisfy the following inequality*

$$(2.6) \quad \sum_{k=2}^{\infty} k(k + |\gamma| - 1) |a_k| \leq |\gamma| \quad (z \in \mathbb{U}, \gamma \in \mathbb{C} \setminus \{0\}),$$

then $f(z) \in \mathcal{K}_{\gamma}$.

Corollary 2.6. *Let $f(z) \in \mathcal{A}$ satisfy the following inequality*

$$(2.7) \quad \sum_{k=2}^{\infty} \frac{[(k-1) + (1-\alpha)|\gamma|] (\alpha_1)_{k-1} \cdots (\alpha_q)_{k-1}}{(\beta_1)_{k-1} \cdots (\beta_s)_{k-1} (k-1)!} |a_k| \leq (1-\alpha)|\gamma|$$

$$(z \in \mathbb{U}; q \leq s+1; q, s \in \mathbb{N}_0; \gamma \in \mathbb{C} \setminus \{0\}; 0 \leq \alpha < 1),$$

then $f(z) \in \mathcal{S}_{\gamma}^*(q, s, \alpha)$.





Title Page

Contents



Page 10 of 16

Go Back

Full Screen

Close

3. Subordination Theorem

Theorem 3.1. Let the function $f(z) \in \mathcal{A}$ satisfy the inequality (2.4), and \mathcal{K} denote the familiar class of functions $h(z) \in \mathcal{A}$ which are univalent and convex in \mathbb{U} . Then for every $\psi \in \mathcal{K}$, we have

$$(3.1) \quad \frac{[1 + (1 - \alpha)|\gamma|]b_2}{2[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]}(f * \psi)(z) \prec \psi(z)$$

$(z \in \mathbb{U}; b_k \geq b_2 > 0 (k \geq 2); \gamma \in \mathbb{C} \setminus \{0\}; 0 \leq \alpha < 1),$

and

$$(3.2) \quad \operatorname{Re}\{f(z)\} > -\frac{[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]}{[1 + (1 - \alpha)|\gamma|]b_2} \quad (z \in \mathbb{U}).$$

The following constant factor

$$\frac{[1 + (1 - \alpha)|\gamma|]b_2}{2[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]}$$

in the subordination result (3.1) is the best dominant.

Proof. Let $f(z)$ satisfy the inequality (2.4) and let $\psi(z) = \sum_{k=0}^{\infty} c_k z^{k+1} \in \mathcal{K}$, then

$$(3.3) \quad \frac{[1 + (1 - \alpha)|\gamma|]b_2}{2[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]}(f * \psi)(z)$$

$$= \frac{[1 + (1 - \alpha)|\gamma|]b_2}{2[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]} \left(z + \sum_{k=2}^{\infty} a_k c_k z^k \right).$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 11 of 16

Go Back

Full Screen

Close

By invoking Definition 1.3, the subordination (3.1) of our theorem will hold true if the sequence

$$(3.4) \quad \left\{ \frac{[1 + (1 - \alpha)|\gamma|]b_2}{2[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]} a_k \right\}_{k=1}^{\infty},$$

is a subordination factor sequence. By virtue of Lemma 1.4, this is equivalent to the inequality

$$(3.5) \quad \operatorname{Re} \left\{ 1 + \sum_{k=1}^{\infty} \frac{[1 + (1 - \alpha)|\gamma|]b_2}{[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]} a_k z^k \right\} > 0 \quad (z \in \mathbb{U}).$$

Since $b_k \geq b_2 > 0$ for $k \geq 2$, we have

$$\begin{aligned} & \operatorname{Re} \left\{ 1 + \sum_{k=1}^{\infty} \frac{[1 + (1 - \alpha)|\gamma|]b_2}{[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]} a_k z^k \right\} \\ &= \operatorname{Re} \left\{ 1 + \frac{[1 + (1 - \alpha)|\gamma|]b_2}{[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]} z + \frac{1}{[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]} \sum_{k=2}^{\infty} [1 + (1 - \alpha)|\gamma|] b_2 a_k z^k \right\} \\ &\geq 1 - \frac{[1 + (1 - \alpha)|\gamma|]b_2}{[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]} r - \frac{1}{[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]} \sum_{k=2}^{\infty} [(k - 1) + (1 - \alpha)|\gamma|] b_k |a_k| r^k \\ &> 1 - \frac{[1 + (1 - \alpha)|\gamma|]b_2}{[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]} r - \frac{(1 - \alpha)|\gamma|}{[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]} r > 0 \quad (|z| = r < 1). \end{aligned}$$

This establishes the inequality (3.5), and consequently the subordination relation (3.1) of Theorem 3.1 is proved. The assertion (3.2) follows readily from (3.1) when the function $\psi(z)$ is selected as

$$(3.6) \quad \psi(z) = \frac{z}{1 - z} = z + \sum_{k=2}^{\infty} z^k \in \mathcal{K}.$$



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 12 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

The sharpness of the multiplying factor in (3.1) can be established by considering a function $h(z)$ defined by

$$h(z) = z - \frac{(1 - \alpha)|\gamma|}{[1 + (1 - \alpha)|\gamma|]} z^2 \quad (z \in \mathbb{U}; \gamma \in \mathbb{C} \setminus \{0\}; 0 \leq \alpha < 1),$$

which belongs to the class $\mathcal{S}_\gamma(g; \alpha)$. Using (3.1), we infer that

$$\frac{[1 + (1 - \alpha)|\gamma|]b_2}{2[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]} h(z) \prec \frac{z}{1 - z}.$$

It can easily be verified that

$$(3.7) \quad \min_{|z| \leq 1} \left[\frac{[1 + (1 - \alpha)|\gamma|]b_2}{2[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]} h(z) \right] = -\frac{1}{2},$$

which shows that the constant

$$\frac{[1 + (1 - \alpha)|\gamma|]b_2}{2[b_2 + (1 - \alpha)(b_2 + 1)|\gamma|]}$$

is the best estimate. □

Before concluding this paper, we consider some useful consequences of the subordination Theorem 3.1.

Corollary 3.2. *Let the function $f(z)$ defined by (1.1) satisfy the inequality (2.5). Then for every $\psi \in \mathcal{K}$, we have*

$$(3.8) \quad \frac{(1 + |\gamma|)}{2(1 + 2|\gamma|)} (f * \psi)(z) \prec \psi(z) \quad (z \in \mathbb{U}),$$



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 13 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

and

$$(3.9) \quad \operatorname{Re}\{f(z)\} > -\frac{(1+2|\gamma|)}{(1+|\gamma|)} \quad (z \in \mathbb{U}).$$

The constant factor

$$\frac{(1+|\gamma|)}{2(1+2|\gamma|)},$$

in the subordination result (3.8) is the best dominant.

Corollary 3.3. Let the function $f(z)$ defined by (1.1) satisfy the inequality (2.6). Then for every $\psi \in \mathcal{K}$, we have

$$(3.10) \quad \frac{(1+|\gamma|)}{(2+3|\gamma|)}(f * \psi)(z) \prec \psi(z) \quad (z \in \mathbb{U}),$$

and

$$(3.11) \quad \operatorname{Re}\{f(z)\} > -\frac{2(2+3|\gamma|)}{(1+|\gamma|)} \quad (z \in \mathbb{U}).$$

The constant factor

$$\frac{(1+|\gamma|)}{(2+3|\gamma|)},$$

in the subordination result (3.10) is the best dominant.

Corollary 3.4. Let the function $f(z)$ defined by (1.1) satisfy the inequality (2.7). Then for every $\psi \in \mathcal{K}$, we have

$$(3.12) \quad \frac{[1+(1-\alpha)|\gamma|]c_2}{2[c_2+(1-\alpha)(c_2+1)|\gamma|]}(f * \psi)(z) \prec \psi(z) \quad (z \in \mathbb{U}),$$



Title Page

Contents



Page 14 of 16

Go Back

Full Screen

Close

and

$$(3.13) \quad \operatorname{Re}\{f(z)\} > -\frac{[c_2 + (1 - \alpha)(c_2 + 1)|\gamma|]}{[1 + (1 - \alpha)|\gamma|]c_2} \quad (z \in \mathbb{U}).$$

The constant factor

$$\frac{[1 + (1 - \alpha)|\gamma|]c_2}{2[c_2 + (1 - \alpha)(c_2 + 1)|\gamma|]},$$

in the subordination result (3.12) is the best dominant, where c_2 is given by

$$c_2 = \frac{\alpha_1 \cdots \alpha_q}{\beta_1 \cdots \beta_s}.$$

Remark 2. On setting $\gamma = 1$ in Corollaries 3.2 and 3.3, we obtain results that correspond to those of Frasin [6, p. 5, Corollary 2.4; p. 6, Corollary 2.7] (see also, Singh [10, p. 434, Corollary 2.2]).

References

- [1] J. DZIOK AND R.K. RAINA, Families of analytic functions associated with the Wright generalized hypergeometric functions, *Demonstratio Math.*, **37** (2004), 533–542.
- [2] J. DZIOK, R.K. RAINA AND H.M. SRIVASTAVA, Some classes of analytic functions associated with operators on Hilbert space involving Wright's generalized hypergeometric functions, *Proc. Jangjeon Math. Soc.*, **7** (2004), 43–55.
- [3] J. DZIOK AND H.M. SRIVASTAVA, Classes of analytic functions associated with the generalized hypergeometric functions, *Appl. Math. Comput.*, **103** (1999), 1–13.
- [4] J. DZIOK AND H.M. SRIVASTAVA, Some subclasses of analytic functions associated with the fixed argument of coefficients associated with the generalized hypergeometric functions, *Adv. Stud. Contemp. Math.*, **5** (2002), 115–125.
- [5] J. DZIOK AND H.M. SRIVASTAVA, Certain subclasses of analytic functions associated with the generalized hypergeometric functions, *Int. Trans. Spec. Funct.*, **14** (2003), 7–18.
- [6] B.A. FRASIN, Subordination results for a class of analytic functions, *J. Inequal. Pure and Appl. Math.*, **7**(4) (2006), Art. 134. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=754>].
- [7] G. MURUGUSUNDARAMOORTHY AND H.M. SRIVASTAVA, Neighborhoods of certain classes of analytic functions of complex order, *J. Inequal. Pure and Appl. Math.*, **5**(2) (2004), Art. 24. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=374>].



Subordination Theorem for a
Family of Analytic Functions

J. K. Prajapat

vol. 9, iss. 4, art. 102, 2008

Title Page

Contents



Page 15 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



- [8] M.A. NASR AND M.K. AOUF, On convex functions of complex order, *Mansoura Sci. Bull. Egypt*, **9** (1982), 565–582.
- [9] M.A. NASR AND M.K. AOUF, Starlike function of complex order, *J. Natur. Sci. Math.*, **25**(1) (1985), 1–12.
- [10] S. SINGH, A subordination theorems for starlike functions, *Int. J. Math. Math. Sci.*, **24**(7) (2000), 433–435.
- [11] H.M. SRIVASTAVA AND S. OWA, *Current Topics in Analytic Functions Theory*, World Scientific Publishing Company, Singapore, New Jersey, London and Hong Kong, 1992.
- [12] P. WIATROWSKI, Subordinating factor sequence for convex, *Zeszyty Nauk. Uniw. Lodz. Nauki Mat. Przyrod. Ser. II*, **39** (1971), 75–85.
- [13] H.S. WILF, Subordinating factor sequence for convex maps of the unit circle, *Proc. Amer. Math. Soc.*, **12** (1961), 689–693.

Subordination Theorem for a
Family of Analytic Functions

J. K. Prajapat

vol. 9, iss. 4, art. 102, 2008

Title Page

Contents



Page 16 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756