

Journal of Inequalities in Pure and Applied Mathematics

SOME PROBLEMS AND SOLUTIONS INVOLVING MATHIEU'S SERIES AND ITS GENERALIZATIONS

H.M. SRIVASTAVA AND ŽIVORAD TOMOVSKI

Department of Mathematics and Statistics
University of Victoria
Victoria, British Columbia V8W 3P4
Canada
E-Mail: harimsri@math.uvic.ca

Institute of Mathematics
St. Cyril and Methodius University
MK-1000 Skopje
Macedonia
E-Mail: tomovski@iunona.pmf.ukim.edu.mk
URL: <http://www.pmf.ukim.edu.mk/mathematics/faculty/tomovski/>

©2000 Victoria University
ISSN (electronic): 1443-5756
146-03



volume 5, issue 2, article 45,
2004.

*Received 14 October, 2003;
accepted 01 May, 2004.*

Communicated by: P. Cerone

Abstract

Contents




Home Page

Go Back

Close

Quit





Abstract

The authors investigate several recently posed problems involving the familiar Mathieu series and its various generalizations. For certain families of generalized Mathieu series, they derive a number of integral representations and investigate several one-sided inequalities which are obtainable from some of these general integral representations or from sundry other considerations. Relevant connections of the results and open problems (which are presented or considered in this paper) with those in earlier works are also indicated. Finally, a conjectured generalization of one of the Mathieu series inequalities proven here is posed as an open problem.

2000 Mathematics Subject Classification: Primary 26D15, 33C10, 33C20, 33C60; Secondary 33E20, 40A30.

Key words: Mathieu's series, Integral representations, Bessel functions, Hypergeometric functions, One-sided inequalities, Fourier transforms, Riemann and Hurwitz Zeta functions, Eulerian integral, Polygamma functions, Laplace integral representation, Euler-Maclaurin summation formula, Riemann-Liouville fractional integral, Lommel function of the first kind.

The present investigation was supported, in part, by the *Natural Sciences and Engineering Research Council of Canada* under Grant OGP0007353.

Dedicated to Professor Blagoj Szdo Popov on the Occasion of his Eightieth Birthday

Contents

1	Introduction, Definitions, and Preliminaries	3
2	A Class of Integral Representations	7
3	Bounds Derivable from the Integral Representation (2.8)....	16
4	Inequalities Associated with Generalized Mathieu Series ...	21
	References	

Some Problems and Solutions Involving Mathieu's Series and Its Generalizations

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 2 of 27

1. Introduction, Definitions, and Preliminaries

The following familiar infinite series:

$$(1.1) \quad S(r) := \sum_{n=1}^{\infty} \frac{2n}{(n^2 + r^2)^2} \quad (r \in \mathbb{R}^+)$$

is named after Émile Leonard Mathieu (1835-1890), who investigated it in his 1890 work [13] on elasticity of solid bodies.

For the Mathieu κ series $S(r)$ defined by (1.1), Alzer *et al.* [2] showed that the best constants κ_1 and κ_2 in the following two-sided inequality:


$$(1.2) \quad \frac{1}{\kappa_1 + r^2} < S(r) < \frac{1}{\kappa_2 + r^2} \quad (r \neq 0)$$

are given by

$$\kappa_1 = \frac{1}{2\zeta(3)} \quad \text{and} \quad \kappa_2 = \frac{1}{6},$$

where $\zeta(s)$ denotes the Riemann Zeta function defined by (see, for details, [20, Chapter 2])

$$(1.3) \quad \zeta(s) := \begin{cases} \sum_{n=1}^{\infty} \frac{1}{n^s} = \frac{1}{1-2^{-s}} \sum_{n=1}^{\infty} \frac{1}{(2n-1)^s} & (\Re(s) > 1) \\ (1-2^{1-s})^{-1} \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^s} & (\Re(s) > 0; s \neq 1). \end{cases}$$



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 3 of 27

A remarkably useful integral representation for $S(r)$ in the elegant form:

$$(1.4) \quad S(r) = \frac{1}{r} \int_0^\infty \frac{x \sin(rx)}{e^x - 1} dx$$

was given by Emersleben [6]. In fact, by applying (1.4) in conjunction with the generating function:

$$(1.5) \quad \frac{z}{e^z - 1} = \sum_{n=0}^{\infty} B_n \frac{z^n}{n!} \quad (|z| < 2\pi)$$


for the Bernoulli numbers

$$B_n \quad (n \in \mathbb{N}_0 := \{0, 1, 2, \dots\}),$$

Elbert [5] derived the following asymptotic expansion for $S(r)$:

$$(1.6) \quad S(r) \sim \sum_{k=0}^{\infty} (-1)^k \frac{B_{2k}}{r^{2k+2}} = \frac{1}{r^2} - \frac{1}{6r^4} - \frac{1}{30r^6} - \dots \quad (r \rightarrow \infty).$$

More recently, Guo [10] made use of the integral representation (1.4) in order to obtain a number of interesting results including (for example) bounds for $S(r)$. For various subsequent developments using (1.4), the interested reader may be referred to the works by (among others) Qi *et al.* ([16] to [19]). (See also an *independent* derivation of the asymptotic expansion (1.6) by Wang and Wang [24]).



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 4 of 27

Several interesting problems and solutions dealing with integral representations and bounds for the following *mild* generalization of the Mathieu series (1.1):

$$(1.7) \quad \mathbb{S}_\mu(r) := \sum_{n=1}^{\infty} \frac{2n}{(n^2 + r^2)^\mu} \quad (r \in \mathbb{R}^+; \mu > 1)$$

can be found in the recent works by Diananda [4], Guo [10], Tomovski and Trenčevski [23], and Cerone and Lenard [3]. Motivated essentially by the works of Cerone and Lenard [3] (and Qi [17]), we propose to investigate the corresponding problems involving a family of generalized Mathieu series, which is defined here by

$$(1.8) \quad \mathcal{S}_\mu^{(\alpha, \beta)}(r; \mathbf{a}) = \mathcal{S}_\mu^{(\alpha, \beta)}(r; \{a_k\}_{k=1}^{\infty}) := \sum_{n=1}^{\infty} \frac{2a_n^\beta}{(a_n^\alpha + r^2)^\mu}$$


$$(r, \alpha, \beta, \mu \in \mathbb{R}^+),$$

where (and throughout this paper) it is tacitly assumed that the positive sequence

$$\mathbf{a} := \{a_k\}_{k=1}^{\infty} = \{a_1, a_2, a_3, \dots, a_k, \dots\} \quad \left(\lim_{k \rightarrow \infty} a_k = \infty \right)$$

is so chosen (and then the *positive* parameters α, β , and μ are so constrained) that the infinite series in the definition (1.8) converges, that is, that the following auxiliary series:

$$\sum_{n=1}^{\infty} \frac{1}{a_n^{\mu\alpha - \beta}}$$



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 5 of 27

is convergent. We remark in passing that, in a very recent research report (which appeared after the submission of this paper to *JIPAM*), Pogány [14] considered a substantially more general form of the definition (1.8). As a matter of fact, Pogány's investigation [14] was based largely upon such *main* mathematical tools as the Laplace integral representation of general Dirichlet series and the familiar Euler-Maclaurin summation formula (*cf.*, *e.g.*, [20, p. 36 *et seq.*]).


Clearly, by comparing the definitions (1.1), (1.7), and (1.8), we obtain

$$(1.9) \quad \mathbb{S}_2(r) = S(r) \quad \text{and} \quad \mathbb{S}_\mu(r) = \mathcal{S}_\mu^{(2,1)}(r; \{k\}_{k=1}^\infty).$$

Furthermore, the special cases

$$\mathcal{S}_2^{(2,1)}(r; \{a_k\}_{k=1}^\infty), \quad \mathcal{S}_\mu^{(2,1)}(r; \{k^\gamma\}_{k=1}^\infty), \quad \text{and} \quad \mathcal{S}_\mu^{(\alpha, \alpha/2)}(r; \{k\}_{k=1}^\infty)$$

were investigated by Qi [17], Tomovski [22], and Cerone and Lenard [3].



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 6 of 27

2. A Class of Integral Representations

First of all, we find from the definition (1.8) that

$$\mathcal{S}_{\mu}^{(\alpha, \beta)}(r; \{a_k\}_{k=1}^{\infty}) = 2 \sum_{m=0}^{\infty} \binom{\mu + m - 1}{m} (-r^2)^m \sum_{n=1}^{\infty} \frac{1}{a_n^{(\mu+m)\alpha - \beta}},$$

so that

$$(2.1) \quad \begin{aligned} \mathcal{S}_{\mu}^{(\alpha, \beta)}(r; \{k^{\gamma}\}_{k=1}^{\infty}) \\ = 2 \sum_{m=0}^{\infty} \binom{\mu + m - 1}{m} (-r^2)^m \zeta(\gamma[(\mu + m)\alpha - \beta]) \\ (r, \alpha, \beta, \gamma \in \mathbb{R}^+; \gamma(\mu\alpha - \beta) > 1) \end{aligned}$$


in terms of the Riemann Zeta function defined by (1.3).

Now, by making use of the familiar integral representation (cf., e.g., [20, p. 96, Equation 2.3 (4)]):

$$(2.2) \quad \zeta(s) = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{x^{s-1}}{e^x - 1} dx \quad (\Re(s) > 1)$$

in (2.1), we obtain

$$(2.3) \quad \begin{aligned} \mathcal{S}_{\mu}^{(\alpha, \beta)}(r; \{k^{\gamma}\}_{k=1}^{\infty}) = \frac{2}{\Gamma(\mu)} \int_0^{\infty} \frac{x^{\gamma(\mu\alpha - \beta) - 1}}{e^x - 1} \\ \cdot {}_1\Psi_1[(\mu, 1); (\gamma(\mu\alpha - \beta), \gamma\alpha); -r^2 x^{\gamma\alpha}] dx, \\ (r, \alpha, \beta, \gamma \in \mathbb{R}^+; \gamma(\mu\alpha - \beta) > 1), \end{aligned}$$



Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 7 of 27

where ${}_p\Psi_q$ denotes the Fox-Wright generalization of the hypergeometric ${}_pF_q$ function with p numerator and q denominator parameters, defined by [21, p. 50, Equation 1.5 (21)]

$$(2.4) \quad {}_p\Psi_q [(\alpha_1, A_1), \dots, (\alpha_p, A_p); (\beta_1, B_1), \dots, (\beta_q, B_q); z] \\ := \sum_{m=0}^{\infty} \frac{\prod_{j=1}^p \Gamma(\alpha_j + A_j m)}{\prod_{j=1}^q \Gamma(\beta_j + B_j m)} \cdot \frac{z^m}{m!},$$

$$\left(A_j \in \mathbb{R}^+ \ (j = 1, \dots, p); B_j \in \mathbb{R}^+ \ (j = 1, \dots, q); 1 + \sum_{j=1}^q B_j - \sum_{j=1}^p A_j > 0 \right),$$

so that, obviously,


$$(2.5) \quad {}_p\Psi_q [(\alpha_1, 1), \dots, (\alpha_p, 1); (\beta_1, 1), \dots, (\beta_q, 1); z] \\ = \frac{\Gamma(\alpha_1) \cdots \Gamma(\alpha_p)}{\Gamma(\beta_1) \cdots \Gamma(\beta_q)} {}_pF_q(\alpha_1, \dots, \alpha_p; \beta_1, \dots, \beta_q; z).$$

In its special case when

$$\gamma\alpha = q \quad (q \in \mathbb{N} := \{1, 2, 3, \dots\}),$$

we can apply the Gauss-Legendre multiplication formula [21, p. 23, Equation 1.1 (27)]:

$$(2.6) \quad \Gamma(mz) = (2\pi)^{\frac{1}{2}(1-m)} m^{mz-\frac{1}{2}} \prod_{j=1}^m \Gamma\left(z + \frac{j-1}{m}\right) \\ \left(z \in \mathbb{C} \setminus \left\{0, -\frac{1}{m}, -\frac{2}{m}, \dots\right\}; m \in \mathbb{N} \right)$$



Some Problems and Solutions Involving Mathieu's Series and Its Generalizations

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 8 of 27

on the right-hand side of our integral representation (2.3). We thus find that

$$(2.7) \quad \mathcal{S}_\mu^{(\alpha, \beta)} \left(r; \{k^{q/\alpha}\}_{k=1}^\infty \right) = \frac{2}{\Gamma \left(q \left[\mu - \frac{\beta}{\alpha} \right] \right)} \int_0^\infty \frac{x^{q \left[\mu - \frac{\beta}{\alpha} \right] - 1}}{e^x - 1} \\ \cdot {}_1F_q \left(\mu; \Delta \left(q; q \left[\mu - \frac{\beta}{\alpha} \right] \right); -r^2 \left(\frac{x}{q} \right)^q \right) dx \\ \left(r, \alpha, \beta \in \mathbb{R}^+; \mu - \frac{\beta}{\alpha} > q^{-1}; q \in \mathbb{N} \right),$$

where, for convenience, $\Delta(q; \lambda)$ abbreviates the array of q parameters

$$\frac{\lambda}{q}, \frac{\lambda + 1}{q}, \dots, \frac{\lambda + q - 1}{q} \quad (q \in \mathbb{N}).$$

For $q = 2$, (2.7) can easily be simplified to the form:

$$(2.8) \quad \mathcal{S}_\mu^{(\alpha, \beta)} \left(r; \{k^{2/\alpha}\}_{k=1}^\infty \right) = \frac{2}{\Gamma \left(2 \left[\mu - \frac{\beta}{\alpha} \right] \right)} \int_0^\infty \frac{x^{2 \left[\mu - \frac{\beta}{\alpha} \right] - 1}}{e^x - 1} \\ \cdot {}_1F_2 \left(\mu; \mu - \frac{\beta}{\alpha}, \mu - \frac{\beta}{\alpha} + \frac{1}{2}; -\frac{r^2 x^2}{4} \right) dx \\ \left(r, \alpha, \beta \in \mathbb{R}^+; \mu - \frac{\beta}{\alpha} > \frac{1}{2} \right).$$

A further special case of (2.8) can be deduced in terms of the Bessel function $J_\nu(z)$ of order ν :



Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 9 of 27

$$(2.9) \quad J_\nu(z) := \sum_{m=0}^{\infty} \frac{(-1)^m \left(\frac{1}{2}z\right)^{\nu+2m}}{m! \Gamma(\nu+m+1)}$$

$$= \frac{\left(\frac{1}{2}z\right)^\nu}{\Gamma(\nu+1)} {}_0F_1\left(-; \nu+1; -\frac{z^2}{4}\right).$$

Thus, by setting $\beta = \frac{1}{2}\alpha$ and $\mu \mapsto \mu + 1$ in (2.8), and applying (2.9) as well as (2.6) with $m = 2$, we obtain the following known result [3, p. 3, Theorem 2.1]:

$$(2.10) \quad \mathcal{S}_{\mu+1}^{(\alpha, \alpha/2)}\left(r; \{k^{2/\alpha}\}_{k=1}^{\infty}\right)$$

$$= \mathcal{S}_{\mu+1}^{(2,1)}\left(r; \{k\}_{k=1}^{\infty}\right) = \mathbb{S}_{\mu+1}(r)$$

$$= \frac{\sqrt{\pi}}{(2r)^{\mu-\frac{1}{2}} \Gamma(\mu+1)} \int_0^\infty \frac{x^{\mu+\frac{1}{2}}}{e^x - 1} J_{\mu-\frac{1}{2}}(rx) dx$$


$$(r, \mu \in \mathbb{R}^+).$$

In a similar manner, a limit case of (2.8) when $\beta \rightarrow 0$ would formally yield the formula:

$$(2.11) \quad \mathcal{S}_\mu^{(\alpha, 0)}\left(r; \{k^{2/\alpha}\}_{k=1}^{\infty}\right) = \sum_{n=1}^{\infty} \frac{2}{(n^2 + r^2)^\mu}$$

$$= \frac{2\sqrt{\pi}}{(2r)^{\mu-\frac{1}{2}} \Gamma(\mu)} \int_0^\infty \frac{x^{\mu-\frac{1}{2}}}{e^x - 1} J_{\mu-\frac{1}{2}}(rx) dx$$

$$\left(r \in \mathbb{R}^+; \mu > \frac{1}{2}\right),$$



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 10 of 27

which is, in fact, equivalent to the following 1906 result of Willem Kapteyn (1849-1927) [25, p. 386, Equation 13.2 (9)]:

$$(2.12) \quad \int_0^\infty \frac{x^\nu}{e^{\pi x} - 1} J_\nu(\lambda x) dt = \frac{(2\lambda)^\nu}{\sqrt{\pi}} \Gamma\left(\nu + \frac{1}{2}\right) \sum_{n=1}^\infty \frac{1}{(n^2 \pi^2 + \lambda^2)^{\nu + \frac{1}{2}}}$$

$$(\Re(\nu) > 0; |\Im(\lambda)| < \pi).$$

Furthermore, a rather simple consequence of (2.11) or (2.12) in the form:

$$(2.13) \quad \sum_{n=-\infty}^\infty \frac{1}{(n^2 + c^2)^s} = c^{-2s} + \frac{2\sqrt{\pi}}{(2c)^{s-\frac{1}{2}} \Gamma(s)} \int_0^\infty \frac{x^{s-\frac{1}{2}}}{e^x - 1} J_{s-\frac{1}{2}}(cx) dx$$

$$\left(\Re(s) > \frac{1}{2}; |c| < 1 \right).$$

appears erroneously in the works by (for example) Hansen [11, p. 122, Entry (6.3.59)] and Prudnikov *et al.* [15, p. 685, Entry 5.1.25.1]. And, by making use of the Trigamma function $\psi'(z)$ defined, in general, by [20, p. 22, Equation 1.2 (52)]


$$(2.14) \quad \psi^{(m)}(z) := \frac{d^{m+1}}{dz^{m+1}} \{\log \Gamma(z)\} = \frac{d^m}{dz^m} \{\psi(z)\}$$

$$(m \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}; z \in \mathbb{C} \setminus \mathbb{Z}_0^-; \mathbb{Z}_0^- := \{0, -1, -2, \dots\})$$

or, equivalently, by

$$(2.15) \quad \psi^{(m)}(z) := (-1)^{m+1} m! \sum_{k=0}^\infty \frac{1}{(k+z)^{m+1}}$$

$$=: (-1)^{m+1} m! \zeta(m+1, z)$$



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 11 of 27

$$(m \in \mathbb{N}; z \in \mathbb{C} \setminus \mathbb{Z}_0^-)$$

in terms of the Hurwitz (or generalized) Zeta function $\zeta(s, a)$ [20, p. 88, Equation 2.2 (1) *et seq.*], both Hansen [11, p. 111, Entry (6.1.137)] and Prudnikov *et al.* [15, p. 687, Entry 5.1.25.28] have recorded the following *explicit* evaluation of the classical Mathieu series:

$$(2.16) \quad S(r) := \sum_{k=1}^{\infty} \frac{2k}{(k^2 + r^2)^2} = \frac{\psi'(-ir) - \psi'(ir)}{2ir} \quad (i := \sqrt{-1}).$$

We remark in passing that, in light of one of the familiar relationships:


$$(2.17) \quad J_{\mp \frac{1}{2}}(z) = \sqrt{\frac{2}{\pi z}} \cdot \begin{cases} \cos z \\ \sin z \end{cases},$$

a special case of (2.10) when $\mu = 1$ would immediately yield the well-exploited integral representation (1.4).

Next, in the theory of Bessel functions, it is fairly well known that (*cf.*, *e.g.*, [7, p. 49, Equation 7.7.3 (16)])

$$(2.18) \quad \int_0^{\infty} e^{-st} t^{\lambda-1} J_{\nu}(\rho t) dt \\ = \left(\frac{\rho}{2s}\right)^{\nu} s^{-\lambda} \frac{\Gamma(\nu + \lambda)}{\Gamma(\nu + 1)} {}_2F_1 \left[\begin{matrix} \frac{1}{2}(\nu + \lambda), \frac{1}{2}(\nu + \lambda + 1); \\ \nu + 1; \end{matrix} -\frac{\rho^2}{s^2} \right]$$

$$(\Re(s) > |\Im(\rho)|; \Re(\nu + \lambda) > 0).$$



Some Problems and Solutions Involving Mathieu's Series and Its Generalizations

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 12 of 27

Since

$$(2.19) \quad {}_1F_0(\lambda; -; z) = (1 - z)^{-\lambda} \quad (|z| < 1; \lambda \in \mathbb{C}),$$

the integral formula (2.18) would simplify considerably when $\lambda = \nu + 1$ and when $\lambda = \nu + 2$, giving us [see also Equations (2.10) and (2.11) above]

$$(2.20) \quad \int_0^\infty e^{-st} t^\nu J_\nu(\rho t) dt = \frac{(2\rho)^\nu}{\sqrt{\pi}} \cdot \frac{\Gamma(\nu + \frac{1}{2})}{(s^2 + \rho^2)^{\nu + \frac{1}{2}}}$$

$$\left(\Re(s) > |\Im(\rho)|; \Re(\nu) > -\frac{1}{2} \right)$$

and


$$(2.21) \quad \int_0^\infty e^{-st} t^{\nu+1} J_\nu(\rho t) dt = \frac{2s(2\rho)^\nu}{\sqrt{\pi}} \cdot \frac{\Gamma(\nu + \frac{3}{2})}{(s^2 + \rho^2)^{\nu + \frac{3}{2}}}$$

$$\left(\Re(s) > |\Im(\rho)|; \Re(\nu) > -1 \right),$$

respectively. While each of the special cases (2.20) and (2.21), too, together with the parent formula (2.18), are readily accessible in many different places in various mathematical books and tables (cf., e.g., [26, p. 72]), (2.20) appears slightly erroneously in [7, p. 49, Equation 7.7.3 (17)]. The integral formula (2.21) would follow also when we differentiate both sides of (2.20) partially with respect to the parameter s .

Now we turn once again to our definition (1.8) which, for $\alpha = 2$, yields

$$(2.22) \quad \mathcal{S}_\mu^{(2,\beta)}(r; \{a_k\}_{k=1}^\infty) = \sum_{n=1}^\infty \frac{2a_n^\beta}{(a_n^2 + r^2)^\mu} \quad (r, \beta, \mu \in \mathbb{R}^+).$$



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 13 of 27

Making use of the integral formulas (2.20) and (2.21), we find from (2.16) that

$$(2.23) \quad \mathcal{S}_\mu^{(2,\beta)}(r; \{a_k\}_{k=1}^\infty) \\ = \frac{2\sqrt{\pi}}{(2r)^{\mu-\frac{1}{2}} \Gamma(\mu)} \int_0^\infty \left(\sum_{n=1}^\infty a_n^\beta e^{-a_n x} \right) x^{\mu-\frac{1}{2}} J_{\mu-\frac{1}{2}}(rx) dx \\ (r, \beta, \mu \in \mathbb{R}^+)$$

and

$$(2.24) \quad \mathcal{S}_\mu^{(2,\beta)}(r; \{a_k\}_{k=1}^\infty) \\ = \frac{\sqrt{\pi}}{(2r)^{\mu-\frac{3}{2}} \Gamma(\mu)} \int_0^\infty \left(\sum_{n=1}^\infty a_n^{\beta-1} e^{-a_n x} \right) x^{\mu-\frac{1}{2}} J_{\mu-\frac{3}{2}}(rx) dx \\ (r, \beta, \mu \in \mathbb{R}^+),$$

respectively.


A special case of the integral representation (2.24) when

$$\beta = 1 \quad \text{and} \quad \mu \mapsto \mu + 1$$

was given by Cerone and Lenard [3, p. 9, Equation (4.5)].

Finally, in view of the Eulerian integral formula:

$$(2.25) \quad \int_0^\infty e^{-st} t^{\lambda-1} dt = \frac{\Gamma(\lambda)}{s^\lambda} \quad (\Re(s) > 0; \Re(\lambda) > 0),$$



Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 14 of 27

we find from the definition (1.8) that

$$(2.26) \quad \mathcal{S}_{\mu}^{(\alpha, \beta)}(r; \{a_k\}_{k=1}^{\infty}) = \frac{2}{\Gamma(\mu)} \int_0^{\infty} x^{\mu-1} e^{-r^2 x} \varphi(x) dx$$

$$(r, \alpha, \beta, \mu \in \mathbb{R}^+),$$


where, for convenience,

$$(2.27) \quad \varphi(x) := \sum_{n=1}^{\infty} a_n^{\beta} \exp(-a_n^{\alpha} x).$$

In terms of the generalized Mathieu series $\mathbb{S}_{\mu}(r)$ defined by (1.7), a special case of the integral representation (2.26) when

$$\alpha = 2, \quad \beta = 1, \quad \text{and} \quad a_k = k \quad (k \in \mathbb{N}),$$

was given by Tomovski and Trenčevski [23, p. 6, Equation (2.3)].



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 15 of 27

3. Bounds Derivable from the Integral Representation (2.8)

For the generalized hypergeometric ${}_pF_q$ function of p numerator and q denominator parameters, which is defined by (2.4) and (2.5), we first recall here the following equivalent form of a familiar Riemann-Liouville fractional integral formula (cf., e.g., [8, p. 200, Entry 13.1 (95)]:


$$(3.1) \quad {}_{p+1}F_{q+1}(\rho, \alpha_1, \dots, \alpha_p; \rho + \sigma, \beta_1, \dots, \beta_q; z) \\ = \frac{\Gamma(\rho + \sigma)}{\Gamma(\rho)\Gamma(\sigma)} \int_0^1 t^{\rho-1} (1-t)^{\sigma-1} {}_pF_q(\alpha_1, \dots, \alpha_p; \beta_1, \dots, \beta_q; zt) dt \\ (p \leq q + 1; \min\{\Re(\rho), \Re(\sigma)\} > 0; |z| < 1 \text{ when } p = q + 1),$$

which, for

$$p = q - 1 = 0 \quad \left(\beta_1 = \mu - \frac{\beta}{\alpha} \right), \quad \rho = \mu, \quad \sigma = \frac{1}{2} - \frac{\beta}{\alpha}, \quad \text{and} \quad z = -\frac{r^2 x^2}{4},$$

immediately yields

$$(3.2) \quad {}_1F_2\left(\mu; \mu - \frac{\beta}{\alpha}, \mu - \frac{\beta}{\alpha} + \frac{1}{2}; -\frac{r^2 x^2}{4}\right) \\ = \frac{\Gamma\left(\mu - \frac{\beta}{\alpha}\right) \Gamma\left(\mu - \frac{\beta}{\alpha} + \frac{1}{2}\right)}{\Gamma(\mu) \Gamma\left(\frac{1}{2} - \frac{\beta}{\alpha}\right)} \left(\frac{2}{rx}\right)^{\mu - (\beta/\alpha) - 1} \\ \cdot \int_0^1 (\sqrt{t})^{\mu + (\beta/\alpha) - 1} (1-t)^{-(\beta/\alpha) - \frac{1}{2}} J_{\mu - (\beta/\alpha) - 1}(rx\sqrt{t}) dt$$



Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 16 of 27

$$\left(r, x, \mu \in \mathbb{R}^+; \frac{\beta}{\alpha} < \frac{1}{2} \right).$$

In terms of the Lommel function $s_{\mu,\nu}(z)$ of the first kind, defined by [7, p. 40, Equation 7.5.5 (69)]

$$(3.3) \quad s_{\mu,\nu}(z) = \frac{z^{\mu+1}}{(\mu - \nu + 1)(\mu + \nu + 1)} \cdot {}_1F_2 \left(1; \frac{1}{2}\mu - \frac{1}{2}\nu + \frac{3}{2}; \frac{1}{2}\mu + \frac{1}{2}\nu + \frac{3}{2}; -\frac{z^2}{4} \right),$$


the special case $\mu = 1$ of (3.2) can be found recorded as a Riemann-Liouville fractional integral formula by Erdélyi *et al.* [8, p. 194, Entry 13.1 (64)] (see also [8, p. 195, Entry 13.1 (65)]).

Now we turn to a recent investigation by Landau [12] in which several best possible uniform bounds for the Bessel functions were obtained by using monotonicity arguments. Following also the work of Cerone and Lenard [3, Section 3], we choose to recall here two of Landau's inequalities given below. The first inequality:

$$(3.4) \quad |J_\nu(x)| \leq \frac{b_L}{\nu^{1/3}}$$

holds true uniformly in the argument x and is the best possible in the exponent $\frac{1}{3}$, with the constant b_L given by

$$(3.5) \quad b_L = 2^{1/3} \sup_x \{ \text{Ai}(x) \} \cong 0.674885 \dots,$$



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 17 of 27

where $\text{Ai}(z)$ denotes the Airy function satisfying the differential equation:

$$(3.6) \quad \frac{d^2 w}{dz^2} - zw = 0 \quad (w = \text{Ai}(z)).$$

The second inequality:

$$(3.7) \quad |J_\nu(x)| \leq \frac{c_L}{x^{1/3}}$$

holds true uniformly in the order $\nu \in \mathbb{R}^+$ and is the best possible in the exponent $\frac{1}{3}$, with the constant c_L given by

$$(3.8) \quad c_L = \sup_x \{x^{1/3} J_0(x)\} \cong 0.78574687\dots$$


By appealing appropriately to the bounds in (3.4) and (3.7), we find from (3.2) that

$$(3.9) \quad \left| {}_1F_2 \left(\mu; \mu - \frac{\beta}{\alpha}, \mu - \frac{\beta}{\alpha} + \frac{1}{2}; -\frac{r^2 x^2}{4} \right) \right|$$

$$\leq b_L \left(\frac{2}{rx} \right)^{\mu - (\beta/\alpha) - 1} \left(\mu - \frac{\beta}{\alpha} - 1 \right)^{-\frac{1}{3}}$$

$$\cdot \frac{\Gamma(\mu - \frac{\beta}{\alpha}) \Gamma(\mu - \frac{\beta}{\alpha} + \frac{1}{2}) \Gamma(\frac{\mu}{2} + \frac{\beta}{2\alpha} + \frac{1}{2})}{\Gamma(\mu) \Gamma(\frac{\mu}{2} - \frac{\beta}{2\alpha} + 1)}$$

$$\left(r, x \in \mathbb{R}^+; \mu - \frac{\beta}{\alpha} > 1; \mu + \frac{\beta}{\alpha} > -1; \frac{\beta}{\alpha} < \frac{1}{2} \right)$$



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 18 of 27

and

$$\begin{aligned}
 (3.10) \quad & \left| {}_1F_2 \left(\mu; \mu - \frac{\beta}{\alpha}, \mu - \frac{\beta}{\alpha} + \frac{1}{2}; -\frac{r^2 x^2}{4} \right) \right| \\
 & \leq \frac{c_L}{(rx)^{1/3}} \left(\frac{2}{rx} \right)^{\mu - (\beta/\alpha) - 1} \\
 & \quad \cdot \frac{\Gamma(\mu - \frac{\beta}{\alpha}) \Gamma(\mu - \frac{\beta}{\alpha} + \frac{1}{2}) \Gamma(\frac{\mu}{2} + \frac{\beta}{2\alpha} + \frac{1}{3})}{\Gamma(\mu) \Gamma(\frac{\mu}{2} - \frac{\beta}{2\alpha} + \frac{5}{6})} \\
 & \quad \left(r, x \in \mathbb{R}^+; \mu - \frac{\beta}{\alpha} > 1; \mu + \frac{\beta}{\alpha} > -\frac{2}{3}; \frac{\beta}{\alpha} < \frac{1}{2} \right),
 \end{aligned}$$

where b_L and c_L are given by (3.5) and (3.8), respectively.

Finally, we apply the inequalities (3.9) and (3.10) in our integral representation (2.8). We thus obtain the following bounds for the generalized Mathieu series occurring in (2.8):

$$\begin{aligned}
 (3.11) \quad & S_{\mu}^{(\alpha, \beta)} \left(r; \{k^{2/\alpha}\}_{k=1}^{\infty} \right) \\
 & \leq \frac{b_L \sqrt{\pi}}{(2r)^{\mu - (\beta/\alpha) - 1}} \left(\mu - \frac{\beta}{\alpha} - 1 \right)^{-\frac{1}{3}} \\
 & \quad \cdot \frac{\Gamma(\mu - \frac{\beta}{\alpha} + 1) \Gamma(\frac{\mu}{2} + \frac{\beta}{2\alpha} + \frac{1}{2})}{\Gamma(\mu) \Gamma(\frac{\mu}{2} - \frac{\beta}{2\alpha} + 1)} \zeta \left(\mu - \frac{\beta}{\alpha} + 1 \right) \\
 & \quad \left(r, x, \alpha, \beta \in \mathbb{R}^+; \frac{\beta}{\alpha} < \frac{1}{2}; \mu - \frac{\beta}{\alpha} > 1 \right)
 \end{aligned}$$



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 19 of 27

and


$$(3.12) \quad \mathcal{S}_\mu^{(\alpha, \beta)} \left(r; \{k^{2/\alpha}\}_{k=1}^\infty \right) \leq \frac{c_L \sqrt{\pi}}{2^{\mu - (\beta/\alpha) - 1} r^{\mu - (\beta/\alpha) - \frac{2}{3}}} \cdot \frac{\Gamma\left(\mu - \frac{\beta}{\alpha} + \frac{2}{3}\right) \Gamma\left(\frac{\mu}{2} + \frac{\beta}{2\alpha} + \frac{1}{2}\right)}{\Gamma(\mu) \Gamma\left(\frac{\mu}{2} - \frac{\beta}{2\alpha} + 1\right)} \zeta\left(\mu - \frac{\beta}{\alpha} + \frac{2}{3}\right) \left(r, x, \alpha, \beta \in \mathbb{R}^+; \frac{\beta}{\alpha} < \frac{1}{2}; \mu - \frac{\beta}{\alpha} > 1 \right),$$

where we have employed the integral representation (2.2) for the Riemann Zeta function $\zeta(s)$, b_L and c_L being given (as before) by (3.5) and (3.8), respectively.

In their special case when

$$\beta \longrightarrow \frac{1}{2}\alpha \quad \text{and} \quad \mu \longmapsto \mu + 1,$$

the bounds in (3.11) and (3.12) would correspond naturally to those given earlier by Cerone and Lenard [3, p. 7, Theorem 3.1]. The *second* bound asserted by Cerone and Lenard [3, p. 7, Equation (3.12)] should, in fact, be corrected to include $\Gamma(\mu + 1)$ in the *denominator* on the right-hand side.



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 20 of 27

4. Inequalities Associated with Generalized Mathieu Series

We first prove the following inequality which was recently posed as an *open* problem by Qi [17, p. 7, Open Problem 2]:

$$(4.1) \quad \left(\int_0^\infty \frac{x \sin(rx)}{e^x - 1} dx \right)^2 > 2r^2 \int_0^\infty x^2 e^{-r^2 x} f(x) dx$$
$$\left(r \in \mathbb{R}^+; f(x) := \sum_{n=1}^\infty n e^{-n^2 x} \right),$$

which, in view of the integral representation (1.4), is equivalent to the inequality:


$$(4.2) \quad [S(r)]^2 > 2 \int_0^\infty x^2 e^{-r^2 x} f(x) dx,$$

where $f(x)$ is defined as in (4.1).

Proof. Since the infinite series:

$$\sum_{n=1}^\infty n e^{-(n^2+r^2)x}$$

is uniformly convergent when $x \in \mathbb{R}^+$, for the right-hand side of the inequality



Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 21 of 27

(4.2), we have

$$\begin{aligned} 2 \int_0^{\infty} x^2 e^{-r^2 x} f(x) dx &= 2 \int_0^{\infty} x^2 \left(\sum_{n=1}^{\infty} n e^{-(n^2+r^2)x} \right) dx \\ &= 2 \sum_{n=1}^{\infty} n \int_0^{\infty} x^2 e^{-(n^2+r^2)x} dx \\ &= 4 \sum_{n=1}^{\infty} \frac{n}{(n^2+r^2)^3} =: 2\mathbb{S}_3(r), \end{aligned}$$

where we have used the Eulerian integral formula (2.25). Hence it is sufficient to prove the following inequality:

$$(4.3) \quad [S(r)]^2 > 2\mathbb{S}_3(r),$$


which was, in fact, conjectured by Alzer and Brenner [2] and proven by Wilkins [27] by remarkably applying series and integral representations for the Trigamma function $\psi'(z)$ defined by (2.14) for $m = 1$. \square

We conclude our present investigation by remarking that it seems to be very likely that the inequality (4.1) can be generalized to the following form:

Open Problem. Prove or disprove that

$$(4.4) \quad \left(\int_0^{\infty} \frac{x \sin(rx)}{e^x - 1} dx \right)^{\mu} > r^{\mu} \Gamma(\mu + 1) \int_0^{\infty} x^{\mu} e^{-r^2 x} f(x) dx$$

$$\left(r, \mu \in \mathbb{R}^+; f(x) := \sum_{n=1}^{\infty} n e^{-n^2 x} \right)$$



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 22 of 27

or, equivalently, that


$$(4.5) \quad [S(r)]^\mu > \frac{\{\Gamma(\mu+1)\}^2}{2} \mathbb{S}_{\mu+1}(r) \\ (r, \mu \in \mathbb{R}^+),$$

since

$$(4.6) \quad \int_0^\infty x^\mu e^{-r^2 x} f(x) dx = \Gamma(\mu+1) \sum_{n=1}^\infty \frac{n}{(n^2+r^2)^{\mu+1}} \\ =: \frac{\Gamma(\mu+1)}{2} \mathbb{S}_{\mu+1}(r),$$

by virtue of the Eulerian integral formula (2.25) once again.

The open problem (4.1), which we have completely solved here, corresponds to the *special* case $\mu = 2$ of the Open Problem (4.4) posed in this paper.



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back


Close

Quit

Page 23 of 27

References

- [1] H. ALZER AND J. BRENNER, An inequality (Problem 97-1), *SIAM Rev.*, **39** (1997), 123.
- [2] H. ALZER, J.L. BRENNER, AND O.G. RUEHR, On Mathieu's inequality, *J. Math. Anal. Appl.*, **218** (1998), 607–610.
- [3] P. CERONE AND C.T. LENARD, On integral forms of generalised Mathieu series, *RGMIA Res. Rep. Coll.*, **6** (2) (2003), Art. 19, 1–11; see also *J. Inequal. Pure Appl. Math.*, **4** (5) (2003), Art. 100, 1–11 (electronic). ONLINE [<http://jipam.vu.edu.au/article.php?sid=341>].
- [4] P.H. DIANANDA, Some inequalities related to an inequality of Mathieu, *Math. Ann.*, **250** (1980), 95–98.
- [5] Á. ELBERT, Asymptotic expansion and continued fraction for Mathieu's series, *Period. Math. Hungar.*, **13** (1982), 1–8.
- [6] O. EMERSLEBEN, Über die Reihe $\sum_{k=1}^{\infty} k / (k^2 + c^2)^2$, *Math. Ann.*, **125** (1952), 165–171.
- [7] A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER, AND F.G. TRICOMI, *Higher Transcendental Functions*, Vol. **II**, McGraw-Hill Book Company, New York, Toronto, and London, 1953.
- [8] A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER, AND F.G. TRICOMI, *Tables of Integral Transforms*, Vol. **II**, McGraw-Hill Book Company, New York, Toronto, and London, 1954.



Some Problems and Solutions Involving Mathieu's Series and Its Generalizations

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents




Go Back

Close

Quit

Page 24 of 27

- [9] I. GAVREA, Some remarks on Mathieu's series, in *Mathematical Analysis and Approximation Theory* (Fifth Romanian-German Seminar on Approximation Theory and Its Applications; Sibiu, Romania, June 12-15, 2002) (A. Lupaş, H. Gonska, and L. Lupaş, Editors), pp. 113–117, Burg Verlag, Sibiu, Romania, 2002.
- [10] B.-N. GUO, Note on Mathieu's inequality, *RGMIA Res. Rep. Coll.*, **3** (3) (2000), Article 5, 1–3. ONLINE [<http://rgmia.vu.edu.au/v3n3.html>].
- [11] E.R. HANSEN, *A Table of Series and Products*, Prentice-Hall, Englewood Cliffs, New Jersey, 1975.
- [12] L. LANDAU, Monotonicity and bounds on Bessel functions, in *Proceedings of the Symposium on Mathematical Physics and Quantum Field Theory* (Berkeley, California; June 11-13, 1999) (H. Warchall, Editor), pp. 147-154; *Electron. J. Differential Equations Conf.*, **4**, Southwest Texas State University, San Marcos, Texas, 2000.
- [13] É.L. MATHIEU, *Traité de Physique Mathématique*. VI-VII: *Théory de l'Élasticité des Corps Solides* (Part 2), Gauthier-Villars, Paris, 1890.
- [14] T.K. POGÁNY, Integral representation of Mathieu (a, λ) -series, *RGMIA Res. Rep. Coll.*, **7** (1) (2004), Article 9, 1–5. ONLINE [<http://rgmia.vu.edu.au/v7n1.html>].
- [15] A.P. PRUDNIKOV, YU. A. BRYČKOV, AND O.I. MARIČEV, *Integrals and Series (Elementary Functions)*, “Nauka”, Moscow, 1981 (Russian); En-



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 25 of 27

glish translation: *Integrals and Series*, Vol. 1: *Elementary Functions*, Gordon and Breach Science Publishers, New York, 1986.

- [16] F. QI, Inequalities for Mathieu's series, *RGMI Res. Rep. Coll.*, **4** (2) (2001), Article 3, 1–7. ONLINE [<http://rgmia.vu.edu.au/v4n2.html>].
- [17] F. QI, Integral expression and inequalities of Mathieu type series, *RGMI Res. Rep. Coll.*, **6** (2) (2003), Article 10, 1–8. ONLINE [<http://rgmia.vu.edu.au/v6n2.html>].
- [18] F. QI, An integral expression and some inequalities of Mathieu type series, *Rostock. Math. Kolloq.*, **58** (2004), 37–46.
- [19] F. QI AND C.-P. CHEN, Notes on double inequalities of Mathieu's series. Preprint 2003.
- [20] H.M. SRIVASTAVA AND J. CHOI, *Series Associated with the Zeta and Related Functions*, Kluwer Academic Publishers, Dordrecht, Boston, and London, 2001.
- [21] H.M. SRIVASTAVA AND H.L. MANOCHA, *A Treatise on Generating Functions*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane, and Toronto, 1984.
- [22] Ž. TOMOVSKI, New double inequalities for Mathieu type series, *RGMI Res. Rep. Coll.*, **6** (2) (2003), Article 17, 1–4. ONLINE [<http://rgmia.vu.edu.au/v6n2.html>].



Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents




Go Back

Close

Quit

Page 26 of 27

- [23] Ž. TOMOVSKI AND K. TRENČEVSKI, On an open problem of Bai-Ni Guo and Feng Qi, *J. Inequal. Pure Appl. Math.*, **4** (2) (2003), Article 29, 1–7 (electronic). ONLINE [<http://jipam.vu.edu.au/article.php?sid=267>].
- [24] C.-L. WANG AND X.-H. WANG, A refinement of the Mathieu inequality, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.*, No. **716-734** (1981), 22–24.
- [25] G.N. WATSON, *A Treatise on the Theory of Bessel Functions*, Second Edition, Cambridge University Press, Cambridge, London, and New York, 1944.
- [26] A.D. WHEELON, *Tables of Summable Series and Integrals Involving Bessel Functions*, Holden-Day, San Francisco, London, and Amsterdam, 1968.
- [27] J.E. WILKINS, Jr., An inequality (Solution of Problem 97-1 posed by H. Alzer and J. Brenner), *SIAM Rev.*, **40** (1998), 126–128.



**Some Problems and Solutions
Involving Mathieu's Series and
Its Generalizations**

H.M. Srivastava and
Živorad Tomovski

Title Page

Contents



Go Back

Close

Quit

Page 27 of 27