



## ON SOME ADVANCED INTEGRAL INEQUALITIES AND THEIR APPLICATIONS

XUEQIN ZHAO AND FANWEI MENG  
xqzhao1972@126.com

DEPARTMENT OF MATHEMATICS  
QUFU NORMAL UNIVERSITY  
QUFU 273165  
PEOPLE'S REPUBLIC OF CHINA  
fwmeng@qfnu.edu.cn

*Received 08 November, 2004; accepted 03 April, 2005*

*Communicated by S.S. Dragomir*

---

ABSTRACT. In this paper, we obtain a generalization of advanced integral inequality and by means of examples we show the usefulness of our results.

---

*Key words and phrases:* Advanced integral inequality; Integral equation.

2000 *Mathematics Subject Classification.* 26D15, 26D10.

### 1. INTRODUCTION

Integral inequalities play an important role in the qualitative analysis of the solutions to differential and integral equations. Many retarded inequalities have been discovered (see [2], [3], [5], [7]). However, we almost neglect the importance of advanced inequalities. After all, it does great benefit to solve the bound of certain integral equations, which help us to fulfill a diversity of desired goals. In this paper we establish two advanced integral inequalities and an application of our results is also given.

### 2. PRELIMINARIES AND LEMMAS

In this paper, we assume throughout that  $\mathbb{R}_+ = [0, \infty)$ , is a subset of the set of real numbers  $\mathbb{R}$ . The following lemmas play an important role in this paper.

**Lemma 2.1.** *Let  $\varphi \in C(\mathbb{R}_+, \mathbb{R}_+)$  be an increasing function with  $\varphi(\infty) = \infty$ . Let  $\psi \in C(\mathbb{R}_+, \mathbb{R}_+)$  be a nondecreasing function and let  $c$  be a nonnegative constant. Let  $\alpha \in C^1(\mathbb{R}_+, \mathbb{R}_+)$  be nondecreasing with  $\alpha(t) \geq t$  on  $\mathbb{R}_+$ . If  $u, f \in C(\mathbb{R}_+, \mathbb{R}_+)$  and*

$$(2.1) \quad \varphi(u(t)) \leq c + \int_{\alpha(t)}^{\infty} f(s)\psi(u(s))ds, \quad t \in \mathbb{R}_+,$$

then for  $0 \leq T \leq t < \infty$ ,

$$(2.2) \quad u(t) \leq \varphi^{-1} \left\{ G^{-1} \left[ G(c) + \int_{\alpha(t)}^{\infty} f(s) ds \right] \right\},$$

where  $G(z) = \int_{z_0}^z \frac{ds}{\psi[\varphi^{-1}(s)]}$ ,  $z \geq z_0 > 0$ ,  $\varphi^{-1}, G^{-1}$  are respectively the inverse of  $\varphi$  and  $G$ ,  $T \in \mathbb{R}_+$  is chosen so that

$$(2.3a) \quad G(c) + \int_{\alpha(t)}^{\infty} f(s) ds \in \text{Dom}(G^{-1}), \quad t \in [T, \infty).$$

$$(2.3b) \quad G^{-1} \left[ G(c) + \int_{\alpha(t)}^{\infty} f(s) ds \right] \in \text{Dom}(\varphi^{-1}), \quad t \in [T, \infty).$$

*Proof.* Define the nonincreasing positive function  $z(t)$  and make

$$(2.4) \quad z(t) = c + \varepsilon + \int_{\alpha(t)}^{\infty} f(s) \psi(u(s)) ds, \quad t \in \mathbb{R}_+,$$

where  $\varepsilon$  is an arbitrary small positive number. From inequality (2.1), we have

$$(2.5) \quad u(t) \leq \varphi^{-1}[z(t)].$$

Differentiating (2.4) and using (2.5) and the monotonicity of  $\varphi^{-1}, \psi$ , we deduce that

$$\begin{aligned} z'(t) &= -f(\alpha(t)) \psi[u(\alpha(t))] \alpha'(t) \\ &\geq -f(\alpha(t)) \psi[\varphi^{-1}(z(\alpha(t)))] \alpha'(t) \\ &\geq -f(\alpha(t)) \psi[\varphi^{-1}(z(t))] \alpha'(t). \end{aligned}$$

For

$$\psi[\varphi^{-1}(z(t))] \geq \psi[\varphi^{-1}(z(\infty))] = \psi[\varphi^{-1}(c + \varepsilon)] > 0,$$

from the definition of  $G$ , the above relation gives

$$\frac{d}{dt} G(z(t)) = \frac{z'(t)}{\psi[\varphi^{-1}(z(t))]} \geq -f(\alpha(t)) \alpha'(t).$$

Setting  $t = s$ , and integrating it from  $t$  to  $\infty$  and letting  $\varepsilon \rightarrow 0$  yields

$$G(z(t)) \leq G(c) + \int_{\alpha(t)}^{\infty} f(s) ds, \quad t \in \mathbb{R}_+.$$

From (2.3), (2.5) and the above relation, we obtain the inequality (2.2).  $\square$

In fact, we can regard Lemma 2.1 as a generalized form of an Ou-Iang type inequality with advanced argument.

**Lemma 2.2.** *Let  $u, f$  and  $g$  be nonnegative continuous functions defined on  $\mathbb{R}_+$ , and let  $\varphi \in C(\mathbb{R}_+, \mathbb{R}_+)$  be an increasing function with  $\varphi(\infty) = \infty$  and let  $c$  be a nonnegative constant. Moreover, let  $w_1, w_2 \in C(\mathbb{R}_+, \mathbb{R}_+)$  be nondecreasing functions with  $w_i(u) > 0$  ( $i = 1, 2$ ) on  $(0, \infty)$ ,  $\alpha \in C^1(\mathbb{R}_+, \mathbb{R}_+)$  be nondecreasing with  $\alpha(t) \geq t$  on  $\mathbb{R}_+$ . If*

$$(2.6) \quad \varphi(u(t)) \leq c + \int_{\alpha(t)}^{\infty} f(s) w_1(u(s)) ds + \int_t^{\infty} g(s) w_2(u(s)) ds, \quad t \in \mathbb{R}_+,$$

then for  $0 \leq T \leq t < \infty$ ,

(i) For the case  $w_2(u) \leq w_1(u)$ ,

$$(2.7) \quad u(t) \leq \varphi^{-1} \left\{ G_1^{-1} \left[ G_1(c) + \int_{\alpha(t)}^{\infty} f(s) ds + \int_t^{\infty} g(s) ds \right] \right\}.$$

(ii) For the case  $w_1(u) \leq w_2(u)$ ,

$$(2.8) \quad u(t) \leq \varphi^{-1} \left\{ G_2^{-1} \left[ G_2(c) + \int_{\alpha(t)}^{\infty} f(s) ds + \int_t^{\infty} g(s) ds \right] \right\},$$

where

$$G_i(r) = \int_{r_0}^r \frac{ds}{w_i(\varphi^{-1}(s))}, \quad r \geq r_0 > 0, \quad (i = 1, 2)$$

and  $\varphi^{-1}$ ,  $G_i^{-1}$  ( $i = 1, 2$ ) are respectively the inverse of  $\varphi$ ,  $G_i$ ,  $T \in \mathbb{R}_+$  is chosen so that

$$(2.9) \quad G_i(c) + \int_{\alpha(t)}^{\infty} f(s) ds + \int_t^{\infty} g(s) ds \in \text{Dom}(G_i^{-1}), \quad (i = 1, 2), \quad t \in [T, \infty).$$

*Proof.* Define the nonincreasing positive function  $z(t)$  and make

$$(2.10) \quad z(t) = c + \varepsilon + \int_{\alpha(t)}^{\infty} f(s) w_1(u(s)) ds + \int_t^{\infty} g(s) w_2(u(s)) ds, \quad 0 \leq T \leq t < \infty,$$

where  $\varepsilon$  is an arbitrary small positive number. From inequality (2.6), we have

$$(2.11) \quad u(t) \leq \varphi^{-1}[z(t)], \quad t \in \mathbb{R}_+.$$

Differentiating (2.10) and using (2.11) and the monotonicity of  $\varphi^{-1}$ ,  $w_1$ ,  $w_2$ , we deduce that

$$\begin{aligned} z'(t) &= -f(\alpha(t)) w_1[u(\alpha(t))] \alpha'(t) - g(t) w_2[u(t)], \\ &\geq -f(\alpha(t)) w_1[\varphi^{-1}(z(\alpha(t)))] \alpha'(t) - g(t) w_2[\varphi^{-1}(z(t))], \\ &\geq -f(\alpha(t)) w_1[\varphi^{-1}(z(t))] \alpha'(t) - g(t) w_2[\varphi^{-1}(z(t))]. \end{aligned}$$

(i) When  $w_2(u) \leq w_1(u)$

$$z'(t) \geq -f(\alpha(t)) w_1[\varphi^{-1}(z(t))] \alpha'(t) - g(t) w_1[\varphi^{-1}(z(t))], \quad t \in \mathbb{R}_+.$$

For

$$w_1[\varphi^{-1}(z(t))] \geq w_1[\varphi^{-1}(z(\infty))] = w_1[\varphi^{-1}(c + \varepsilon)] > 0,$$

from the definition of  $G_1(r)$ , the above relation gives

$$\frac{d}{dt} G_1(z(t)) = \frac{z'(t)}{w_1[\varphi^{-1}(z(t))]} \geq -f(\alpha(t)) \alpha'(t) - g(t), \quad t \in \mathbb{R}_+.$$

Setting  $t = s$  and integrating it from  $t$  to  $\infty$  and let  $\varepsilon \rightarrow 0$  yields

$$G_1(z(t)) \leq G_1(c) + \int_{\alpha(t)}^{\infty} f(s) ds + \int_t^{\infty} g(s) ds, \quad t \in \mathbb{R}_+,$$

so,

$$z(t) \leq G_1^{-1} \left[ G_1(c) + \int_{\alpha(t)}^{\infty} f(s) ds + \int_t^{\infty} g(s) ds \right], \quad 0 \leq T \leq t < \infty.$$

Using (2.11), we have

$$u(t) \leq \varphi^{-1} \left\{ G_1^{-1} \left[ G_1(c) + \int_{\alpha(t)}^{\infty} f(s) ds + \int_t^{\infty} g(s) ds \right] \right\}, \quad 0 \leq T \leq t < \infty.$$

(ii) When  $w_1(u) \leq w_2(u)$ , the proof can be completed similarly.  $\square$

### 3. MAIN RESULTS

In this section, we obtain our main results as follows:

**Theorem 3.1.** *Let  $u$ ,  $f$  and  $g$  be nonnegative continuous functions defined on  $\mathbb{R}_+$  and let  $c$  be a nonnegative constant. Moreover, let  $\varphi \in C(\mathbb{R}_+, \mathbb{R}_+)$  be an increasing function with  $\varphi(\infty) = \infty$ ,  $\psi \in C(\mathbb{R}_+, \mathbb{R}_+)$  be a nondecreasing function with  $\psi(u) > 0$  on  $(0, \infty)$  and  $\alpha \in C^1(\mathbb{R}_+, \mathbb{R}_+)$  be nondecreasing with  $\alpha(t) \geq t$  on  $\mathbb{R}_+$ . If*

$$(3.1) \quad \varphi(u(t)) \leq c + \int_{\alpha(t)}^{\infty} [f(s)u(s)\psi(u(s)) + g(s)u(s)]ds, \quad t \in \mathbb{R}_+$$

then for  $0 \leq T \leq t < \infty$ ,

$$(3.2) \quad u(t) \leq \varphi^{-1} \left\{ \Omega^{-1} \left[ G^{-1} \left( G[\Omega(c) + \int_{\alpha(t)}^{\infty} g(s)ds] + \int_{\alpha(t)}^{\infty} f(s)ds \right) \right] \right\},$$

where

$$\Omega(r) = \int_{r_0}^r \frac{ds}{\varphi^{-1}(s)}, \quad r \geq r_0 > 0, \quad G(z) = \int_{z_0}^z \frac{ds}{\psi\{\varphi^{-1}[\Omega^{-1}(s)]\}}, \quad z \geq z_0 > 0,$$

$\Omega^{-1}, \varphi^{-1}, G^{-1}$  are respectively the inverse of  $\Omega, \varphi, G$  and  $T \in \mathbb{R}_+$  is chosen so that

$$G \left[ \Omega(c) + \int_{\alpha(t)}^{\infty} g(s)ds \right] + \int_{\alpha(t)}^{\infty} f(s)ds \in \text{Dom}(G^{-1})$$

and

$$G^{-1} \left\{ G \left[ \Omega(c) + \int_{\alpha(t)}^{\infty} g(s)ds \right] + \int_{\alpha(t)}^{\infty} f(s)ds \right\} \in \text{Dom}(\Omega^{-1})$$

for  $t \in [T, \infty)$ .

*Proof.* Let us first assume that  $c > 0$ . Define the nonincreasing positive function  $z(t)$  by the right-hand side of (3.1). Then  $z(\infty) = c$ ,  $u(t) \leq \varphi^{-1}[z(t)]$  and

$$\begin{aligned} z'(t) &= - [f(\alpha(t))u(\alpha(t))\psi[u(\alpha(t))] - g(\alpha(t))u(\alpha(t))] \alpha'(t) \\ &\geq - [f(\alpha(t))\varphi^{-1}(z(\alpha(t)))\psi[\varphi^{-1}(z(\alpha(t)))] - g(\alpha(t))\varphi^{-1}(z(\alpha(t)))] \alpha'(t) \\ &\geq - [f(\alpha(t))\varphi^{-1}(z(t))\psi[\varphi^{-1}(z(\alpha(t)))] - g(\alpha(t))\varphi^{-1}(z(t))] \alpha'(t). \end{aligned}$$

Since  $\varphi^{-1}(z(t)) \geq \varphi^{-1}(c) > 0$ ,

$$\frac{z'(t)}{\varphi^{-1}(z(t))} \geq - \{f(\alpha(t))\psi[\varphi^{-1}(z(\alpha(t)))] + g(\alpha(t))\} \alpha'(t).$$

Setting  $t = s$  and integrating it from  $t$  to  $\infty$  yields

$$\Omega(z(t)) \leq \Omega(c) + \int_{\alpha(t)}^{\infty} g(s)ds + \int_{\alpha(t)}^{\infty} f(s)\psi[\varphi^{-1}(z(s))]ds.$$

Let  $T \leq T_1$  be an arbitrary number. We denote  $p(t) = \Omega(c) + \int_{\alpha(t)}^{\infty} g(s)ds$ . From the above relation, we deduce that

$$\Omega(z(t)) \leq p(T_1) + \int_{\alpha(t)}^{\infty} f(s)\psi[\varphi^{-1}(z(s))]ds, \quad T_1 \leq t < \infty.$$

Now an application of Lemma 2.1 gives

$$z(t) \leq \Omega^{-1} \left\{ G^{-1} \left[ G(p(T_1)) + \int_{\alpha(t)}^{\infty} f(s)ds \right] \right\}, \quad T_1 \leq t < \infty,$$

so,

$$u(t) \leq \varphi^{-1} \left\{ \Omega^{-1} \left[ G^{-1} \left( G(p(T_1)) + \int_{\alpha(t)}^{\infty} f(s) ds \right) \right] \right\}, \quad T_1 \leq t < \infty.$$

Taking  $t = T_1$  in the above inequality, since  $T_1$  is arbitrary, we can prove the desired inequality (3.2).

If  $c = 0$  we carry out the above procedure with  $\varepsilon > 0$  instead of  $c$  and subsequently let  $\varepsilon \rightarrow 0$ .  $\square$

**Corollary 3.2.** *Let  $u, f$  and  $g$  be nonnegative continuous functions defined on  $\mathbb{R}_+$  and let  $c$  be a nonnegative constant. Moreover, let  $\psi \in C(\mathbb{R}_+, \mathbb{R}_+)$  be a nondecreasing function with  $\psi(u) > 0$  on  $(0, \infty)$  and  $\alpha \in C^1(\mathbb{R}_+, \mathbb{R}_+)$  be nondecreasing with  $\alpha(t) \geq t$  on  $\mathbb{R}_+$ . If*

$$u^2(t) \leq c^2 + \int_{\alpha(t)}^{\infty} [f(s)u(s)\psi(u(s)) + g(s)u(s)] ds, \quad t \in \mathbb{R}_+,$$

then for  $0 \leq T \leq t < \infty$ ,

$$u(t) \leq \Omega^{-1} \left[ \Omega \left( c + \frac{1}{2} \int_{\alpha(t)}^{\infty} g(s) ds \right) + \frac{1}{2} \int_{\alpha(t)}^{\infty} f(s) ds \right],$$

where

$$\Omega(r) = \int_1^r \frac{ds}{\psi(s)}, \quad r > 0,$$

$\Omega^{-1}$  is the inverse of  $\Omega$ , and  $T \in \mathbb{R}_+$  is chosen so that

$$\Omega \left( c + \frac{1}{2} \int_{\alpha(t)}^{\infty} g(s) ds \right) + \frac{1}{2} \int_{\alpha(t)}^{\infty} f(s) ds \in \text{Dom}(\Omega^{-1})$$

for all  $t \in [T, \infty)$ .

**Corollary 3.3.** *Let  $u, f$  and  $g$  be nonnegative continuous functions defined on  $\mathbb{R}_+$  and let  $c$  be a nonnegative constant. Moreover, let  $\alpha \in C^1(\mathbb{R}_+, \mathbb{R}_+)$  be nondecreasing with  $\alpha(t) \geq t$  on  $\mathbb{R}_+$ . If*

$$u^2(t) \leq c^2 + \int_{\alpha(t)}^{\infty} [f(s)u^2(s) + g(s)u(s)] ds, \quad t \geq 0,$$

then

$$u(t) \leq \left( c + \frac{1}{2} \int_{\alpha(t)}^{\infty} g(s) ds \right) \exp \left[ \frac{1}{2} \int_{\alpha(t)}^{\infty} f(s) ds \right], \quad t \geq 0.$$

**Corollary 3.4.** *Let  $u, f$  and  $g$  be nonnegative continuous functions defined on  $\mathbb{R}_+$  and let  $c$  be a nonnegative constant. Moreover, let  $p, q$  be positive constants with  $p \geq q, p \neq 1$ . Let  $\alpha \in C^1(\mathbb{R}_+, \mathbb{R}_+)$  be nondecreasing with  $\alpha(t) \geq t$  on  $\mathbb{R}_+$ . If*

$$u^p(t) \leq c + \int_{\alpha(t)}^{\infty} [f(s)u^q(s) + g(s)u(s)] ds, \quad t \in \mathbb{R}_+,$$

then for  $t \in \mathbb{R}_+$ ,

$$u(t) \leq \begin{cases} \left( c^{(1-\frac{1}{p})} + \frac{p-1}{p} \int_{\alpha(t)}^{\infty} g(s) ds \right)^{\frac{p}{p-1}} \exp \left[ \frac{1}{p} \int_{\alpha(t)}^{\infty} f(s) ds \right], & \text{when } p = q, \\ \left[ \left( c^{(1-\frac{1}{p})} + \frac{p-1}{p} \int_{\alpha(t)}^{\infty} g(s) ds \right)^{\frac{p-q}{p-1}} + \frac{p-q}{p} \int_{\alpha(t)}^{\infty} f(s) ds \right]^{\frac{1}{p-q}}, & \text{when } p > q. \end{cases}$$

**Theorem 3.5.** Let  $u, f$  and  $g$  be nonnegative continuous functions defined on  $\mathbb{R}_+$ , and let  $\varphi \in C(\mathbb{R}_+, \mathbb{R}_+)$  be an increasing function with  $\varphi(\infty) = \infty$  and let  $c$  be a nonnegative constant. Moreover, let  $w_1, w_2 \in C(\mathbb{R}_+, \mathbb{R}_+)$  be nondecreasing functions with  $w_i(u) > 0$  ( $i = 1, 2$ ) on  $(0, \infty)$  and  $\alpha \in C^1(\mathbb{R}_+, \mathbb{R}_+)$  be nondecreasing with  $\alpha(t) \geq t$  on  $\mathbb{R}_+$ . If

$$(3.3) \quad \varphi(u(t)) \leq c + \int_{\alpha(t)}^{\infty} f(s)u(s)w_1(u(s))ds + \int_t^{\infty} g(s)u(s)w_2(u(s))ds,$$

then for  $0 \leq T \leq t < \infty$ ,

(i) For the case  $w_2(u) \leq w_1(u)$ ,

$$(3.4) \quad u(t) \leq \varphi^{-1} \left\{ \Omega^{-1} \left[ G_1^{-1} \left( G_1(\Omega(c)) + \int_{\alpha(t)}^{\infty} f(s)ds + \int_t^{\infty} g(s)ds \right) \right] \right\},$$

(ii) For the case  $w_1(u) \leq w_2(u)$ ,

$$(3.5) \quad u(t) \leq \varphi^{-1} \left\{ \Omega^{-1} \left[ G_2^{-1} \left( G_2(\Omega(c)) + \int_{\alpha(t)}^{\infty} f(s)ds + \int_t^{\infty} g(s)ds \right) \right] \right\},$$

where

$$\Omega(r) = \int_{r_0}^r \frac{ds}{\varphi^{-1}(s)}, \quad r \geq r_0 > 0,$$

$$G_i(z) = \int_{z_0}^z \frac{ds}{w_i\{\varphi^{-1}[\Omega^{-1}(s)]\}}, \quad z \geq z_0 > 0 \quad (i = 1, 2),$$

$\Omega^{-1}, \varphi^{-1}, G^{-1}$  are respectively the inverse of  $\Omega, \varphi, G$ , and  $T \in \mathbb{R}_+$  is chosen so that

$$G_i \left( \Omega(c) + \int_{\alpha(t)}^{\infty} f(s)ds + \int_t^{\infty} g(s)ds \right) \in \text{Dom}(G_i^{-1}),$$

$$G_i^{-1} \left[ G_i \left( \Omega(c) + \int_{\alpha(t)}^{\infty} f(s)ds + \int_t^{\infty} g(s)ds \right) \right] \in \text{Dom}(\Omega^{-1}),$$

for all  $t \in [T, \infty)$ .

*Proof.* Let  $c > 0$ , define the nonincreasing positive function  $z(t)$  and make

$$(3.6) \quad z(t) = c + \int_{\alpha(t)}^{\infty} f(s)u(s)w_1(u(s))ds + \int_t^{\infty} g(s)u(s)w_2(u(s))ds.$$

From inequality (3.3), we have

$$(3.7) \quad u(t) \leq \varphi^{-1}[z(t)].$$

Differentiating (3.6) and using (3.7) and the monotonicity of  $\varphi^{-1}, w_1, w_2$ , we deduce that

$$\begin{aligned} z'(t) &= -f(\alpha(t))u(\alpha(t))w_1[u(\alpha(t))] \alpha'(t) - g(t)u(t)w_2[u(t)], \\ &\geq -f(\alpha(t))\varphi^{-1}(z(\alpha(t)))w_1[\varphi^{-1}(z(\alpha(t)))] \alpha'(t) - g(t)\varphi^{-1}(z(t))w_2[\varphi^{-1}(z(t))], \\ &\geq -f(\alpha(t))\varphi^{-1}(z(t))w_1[\varphi^{-1}(z(t))] \alpha'(t) - g(t)\varphi^{-1}(z(t))w_2[\varphi^{-1}(z(t))]. \end{aligned}$$

(i) When  $w_2(u) \leq w_1(u)$

$$\frac{z'(t)}{\varphi^{-1}(z(t))} \geq -f(\alpha(t))w_1[\varphi^{-1}(z(t))] \alpha'(t) - g(t)w_1[\varphi^{-1}(z(t))].$$

For

$$w_1[\varphi^{-1}(z(t))] \geq w_1[\varphi^{-1}(z(\infty))] = w_1[\varphi^{-1}(c + \varepsilon)] > 0,$$

setting  $t = s$  and integrating from  $t$  to  $\infty$  yields

$$\Omega(z(t)) \leq \Omega(c) + \int_{\alpha(t)}^{\infty} f(s)w_1[\varphi^{-1}(z(t))] ds + \int_t^{\infty} g(s)w_1[\varphi^{-1}(z(t))] ds.$$

From Lemma 2.2, we obtain

$$z(t) \leq \Omega^{-1} \left\{ G_1^{-1} \left[ G_1(\Omega(c)) + \int_{\alpha(t)}^{\infty} f(s)ds + \int_t^{\infty} g(s)ds \right] \right\}, \quad 0 \leq T \leq t < \infty.$$

Using  $u(t) \leq \varphi^{-1}[z(t)]$ , we get the inequality in (3.4)

If  $c = 0$ , we can carry out the above procedure with  $\varepsilon > 0$  instead of  $c$  and subsequently let  $\varepsilon \rightarrow 0$ .

(ii) When  $w_1(u) \leq w_2(u)$ , the proof can be completed similarly.  $\square$

#### 4. AN APPLICATION

We consider an integral equation

$$(4.1) \quad x^p(t) = a(t) + \int_t^{\infty} F[s, x(s), x(\phi(s))]ds.$$

Assume that:

$$(4.2) \quad |F(x, y, u)| \leq f(x)|u|^q + g(x)|u|$$

and

$$(4.3) \quad |a(t)| \leq c, \quad c > 0 \quad p \geq q > 0, \quad p \neq 1,$$

where  $f, g$  are nonnegative continuous real-valued functions, and  $\phi \in C^1(\mathbb{R}_+, \mathbb{R}_+)$  is nondecreasing with  $\phi(t) \geq t$  on  $\mathbb{R}_+$ . From (4.1), (4.2) and (4.3) we have

$$|x(t)|^p \leq c + \int_t^{\infty} f(s)|x(\phi(s))|^q + g(s)|x(\phi(s))|ds.$$

Making the change of variables from the above inequality and taking

$$M = \sup_{t \in \mathbb{R}_+} \frac{1}{\phi'(t)},$$

we have

$$|x(t)|^p \leq c + M \int_{\phi(t)}^{\infty} \bar{f}(s)|x(s)|^q + \bar{g}(s)|x(s)|ds,$$

in which  $\bar{f}(s) = f(\phi^{-1}(s))$ ,  $\bar{g}(s) = g(\phi^{-1}(s))$ . From Corollary 3.4, we obtain

$$|x(t)| \leq \begin{cases} \left( c^{(1-\frac{1}{p})} + \frac{M(p-1)}{p} \int_{\phi(t)}^{\infty} \bar{g}(s)ds \right)^{\frac{p}{p-1}} \exp \left[ \frac{M}{p} \int_{\phi(t)}^{\infty} \bar{f}(s)ds \right], & \text{when } p = q \\ \left[ \left( c^{(1-\frac{1}{p})} + \frac{M(p-1)}{p} \int_{\phi(t)}^{\infty} \bar{g}(s)ds \right)^{\frac{p-q}{p-1}} + \frac{M(p-q)}{p} \int_{\phi(t)}^{\infty} \bar{f}(s)ds \right]^{\frac{1}{p-q}}, & \text{when } p > q. \end{cases}$$

If the integrals of  $f(s), g(s)$  are bounded, then we have the bound of the solution of (4.1).

## REFERENCES

- [1] I. BIHARI, A generalization of a lemma of Bellman and its application to uniqueness problems of differential equations, *Acta. Math. Acad. Sci. Hungar.*, **7** (1956), 71–94.
- [2] O. LIPOVAN, A retarded Gronwall-like inequality and its applications, *J. Math. Anal. Appl.*, **252** (2000), 389–401.
- [3] O. LIPOVAN, A retarded Integral inequality and its applications, *J. Math. Anal. Appl.*, **285** (2003), 436–443.
- [4] M. MEDVED, Nonlinear singular integral inequalities for functions in two and  $n$  independent variables, *J. Inequalities and Appl.*, **5** (2000), 287–308.
- [5] B.G. PACHPATTE, On some new inequalities related to certain inequalities in the theory of differential equations, *J. Math. Anal. Appl.*, **189** (1995), 128–144.
- [6] B.G. PACHPATTE, Explicit bounds on certain integral inequalities, *J. Math. Anal. Appl.*, **267** (2002), 48–61.
- [7] B.G. PACHPATTE, On some new inequalities related to a certain inequality arising in the theory of differential equations, *J. Math. Anal. Appl.*, **251** (2000), 736–751.