

A.M. Bikchentaev

THE CONTINUITY OF MULTIPLICATION FOR TWO  
TOPOLOGIES ASSOCIATED WITH A SEMIFINITE  
TRACE ON VON NEUMANN ALGEBRA

(submitted by D. Mushtari)

ABSTRACT. Let  $\mathcal{M}$  be a semifinite von Neumann algebra in a Hilbert space  $\mathcal{H}$  and  $\tau$  be a normal faithful semifinite trace on  $\mathcal{M}$ . Let  $\mathcal{M}^{\text{pr}}$  denote the set of all projections in  $\mathcal{M}$ ,  $e$  denote the unit of  $\mathcal{M}$ , and  $\|\cdot\|$  denote the  $C^*$ -norm on  $\mathcal{M}$ .

The set of all  $\tau$ -measurable operators  $\widetilde{\mathcal{M}}$  with sum and product defined as the respective closures of the usual sum and product, is a  $*$ -algebra. The sets  $U(\varepsilon, \delta) = \{x \in \widetilde{\mathcal{M}} : \|xp\| \leq \varepsilon \text{ and } \tau(e - p) \leq \delta \text{ for some } p \in \mathcal{M}^{\text{pr}}, \varepsilon > 0, \delta > 0\}$ , form a base at 0 for a metrizable vector topology  $t_\tau$  on  $\widetilde{\mathcal{M}}$ , called *the measure topology*. Equipped with this topology,  $\widetilde{\mathcal{M}}$  is a complete topological  $*$ -algebra. We will write  $x_i \xrightarrow{\tau} x$  in case a net  $\{x_i\}_{i \in I} \subset \widetilde{\mathcal{M}}$  converges to  $x \in \widetilde{\mathcal{M}}$  for the measure topology on  $\widetilde{\mathcal{M}}$ . By definition, a net  $\{x_i\}_{i \in I} \subset \widetilde{\mathcal{M}}$  converges  $\tau$ -locally to  $x \in \widetilde{\mathcal{M}}$  (notation:  $x_i \xrightarrow{\tau^l} x$ ) if  $x_i p \xrightarrow{\tau} xp$  for all  $p \in \mathcal{M}^{\text{pr}}$ ,  $\tau(p) < \infty$ ; and a net  $\{x_i\}_{i \in I} \subset \widetilde{\mathcal{M}}$  converges weak  $\tau$ -locally to  $x \in \widetilde{\mathcal{M}}$  (notation:  $x_i \xrightarrow{w\tau^l} x$ ) if  $px_i p \xrightarrow{\tau} p x p$  for all  $p \in \mathcal{M}^{\text{pr}}$ ,  $\tau(p) < \infty$ .

**Theorem 1.** Let  $x_i, x \in \widetilde{\mathcal{M}}$ .

1. If  $x_i \xrightarrow{\tau^l} x$ , then  $x_i y \xrightarrow{\tau^l} xy$  and  $y x_i \xrightarrow{\tau^l} yx$  for every fixed  $y \in \widetilde{\mathcal{M}}$ .

---

2000 Mathematical Subject Classification. 46L51, 46L53, 46E30, 46H05.

Key words and phrases. Hilbert space, von Neumann algebra, noncommutative integration, measurable operator, semifinite trace, convergence with respect to measure, compact operator, topological algebra.

Partially supported by Russian Foundation for Basic Research (Grant 01-01-00129) and the scientific program "Universities of Russia – Basic Research" (Grant UR. 04.01.061) .

**2.** If  $x_i \xrightarrow{w\tau^l} x$ , then  $x_i y \xrightarrow{w\tau^l} xy$  and  $y x_i \xrightarrow{w\tau^l} yx$  for every fixed  $y \in \widetilde{\mathcal{M}}$ .

**Theorem 2.** If  $\{x_i\}_{i \in I} \subset \widetilde{\mathcal{M}}$  is bounded in measure and if  $x_i \xrightarrow{\tau^l} x \in \widetilde{\mathcal{M}}$ , then  $x_i y \xrightarrow{\tau} xy$  for all  $\tau$ -compact  $y \in \widetilde{\mathcal{M}}$ .

**Theorem 3.** Let  $x, y, x_i, y_i \in \widetilde{\mathcal{M}}$  and let a set  $\{x_i\}_{i \in I}$  be bounded in measure. If  $x_i \xrightarrow{\tau^l} x$  and  $y_i \xrightarrow{\tau^l} y$ , then  $x_i y_i \xrightarrow{\tau^l} xy$ .

If  $\mathcal{M}$  is abelian, then the weak  $\tau$ -local and  $\tau$ -local convergencies on  $\widetilde{\mathcal{M}}$  coincides with the familiar convergence locally in measure. If  $\tau(e) = \infty$ , then the boundedness condition cannot be omitted in Theorem 2.

If  $\mathcal{M}$  is  $\mathcal{B}(\mathcal{H})$  with standard trace, then Theorem 2 for sequences is a "Basic lemma" of the theory of projection methods: *If  $y$  is compact and  $x_n \rightarrow x$  strongly, then  $x_n y \rightarrow xy$  uniformly, i.e.  $\|x_n y - xy\| \rightarrow 0$  as  $n \rightarrow \infty$ .* Theorem 3 means that the mapping

$$(x, y) \mapsto xy : (\mathcal{B}(\mathcal{H})_1 \times \mathcal{B}(\mathcal{H})) \rightarrow \mathcal{B}(\mathcal{H})$$

is strong-operator continuous ( $\mathcal{B}(\mathcal{H})_1$  denotes the unit ball of  $\mathcal{B}(\mathcal{H})$ ).

*The author is greatly indebted to O. E. Tikhonov for drawing author's attention to the problem of the  $\tau$ -local continuity of operator functions.*

## 1. INTRODUCTION

Let  $\mathcal{M}$  be a semifinite von Neumann algebra of operators in a Hilbert space  $\mathcal{H}$  and  $\tau$  be a distinguished normal faithful semifinite trace on  $\mathcal{M}$ . Let  $\mathcal{M}^{\text{pr}}$  denote the lattice of all projections in  $\mathcal{M}$ ,  $e$  denote the identity, and  $\mathcal{M}_1$  denote the unit ball of  $\mathcal{M}$  in the  $C^*$ -norm  $\|\cdot\|$  on  $\mathcal{M}$ . The closed, densely defined linear operator  $x$  in  $\mathcal{H}$  with domain  $\mathcal{D}(x)$  is said to be *affiliated* with  $\mathcal{M}$  if and only if  $u^* x u = x$  for all unitary operators  $u$  in the commutant  $\mathcal{M}'$  of  $\mathcal{M}$ . If  $x$  is affiliated with  $\mathcal{M}$  then  $x$  is said to be  $\tau$ -*measurable* if and only if, for every  $\varepsilon > 0$  there exists a projection  $p \in \mathcal{M}^{\text{pr}}$  for which  $p(\mathcal{H}) \subseteq \mathcal{D}(x)$  and  $\tau(e - p) < \varepsilon$ . We denote by  $\widetilde{\mathcal{M}}$  the set of all  $\tau$ -measurable operators. With sum and product defined as the respective closures of the usual sum and product,  $\widetilde{\mathcal{M}}$  is a  $*$ -algebra. The sets

$$U(\varepsilon, \delta) = \{x \in \widetilde{\mathcal{M}} : \|xp\| \leq \varepsilon \text{ and } \tau(e - p) \leq \delta \text{ for some } p \in \mathcal{M}^{\text{pr}}\},$$

where  $\varepsilon > 0$ ,  $\delta > 0$ , form a base at 0 for a metrizable vector topology  $t_\tau$  on  $\widetilde{\mathcal{M}}$ , called *the measure topology* ([8]; [11, p. 18]). Equipped with this topology,  $\widetilde{\mathcal{M}}$  is a complete topological  $*$ -algebra in which  $\mathcal{M}$  is dense. We will write  $x_i \xrightarrow{\tau} x$  in case a net  $\{x_i\}_{i \in I} \subset \widetilde{\mathcal{M}}$  converges to  $x \in \widetilde{\mathcal{M}}$  for the measure topology on  $\widetilde{\mathcal{M}}$ .

A subset  $X$  of  $\widetilde{\mathcal{M}}$  is *bounded in measure*, if it is bounded with respect to this topology on the vector space of  $\widetilde{\mathcal{M}}$ , that is in case for every neighborhood  $U$  of 0 there is an  $\alpha > 0$  such that  $\alpha X \subset U$  [8, p. 106].

If  $\mathcal{M}$  is  $\mathcal{B}(\mathcal{H})$ , the von Neumann algebra of all bounded linear operators in  $\mathcal{H}$  is equipped with the usual standard trace, then  $\widetilde{\mathcal{M}}$  coincides with  $\mathcal{M}$  and in this case the measure topology coincides with the  $\|\cdot\|$ -topology. If  $\mathcal{M}$  is abelian, then  $\mathcal{M}$  may be identified with  $L^\infty(\Omega, \mu)$  and  $\tau(f) = \int_\Omega f \, d\mu$  where  $(\Omega, \mu)$  is a localizable measure space. In this case,  $\widetilde{\mathcal{M}}$  is the space  $S_0(\Omega)$  consisting of those measurable complex-valued functions on  $\Omega$  which are bounded except on a set of finite measure and the measure topology on  $\widetilde{\mathcal{M}}$  may be identified simply with the familiar topology of convergence in measure.

If  $x$  is any self-adjoint operator in  $\mathcal{H}$  and if

$$x = \int_{\mathbb{R}} \lambda \, de_\lambda^x$$

is its spectral representation, we will write  $\chi_T(x)$  for the spectral projection of  $x$  corresponding to the Borel subset  $T \subset \mathbb{R}$ . In particular  $e_\lambda^x = \chi_{(-\infty, \lambda]}(x)$ . If  $x$  is closed, densely defined linear operator affiliated with  $\mathcal{M}$  and  $|x| = \sqrt{x^*x}$ , then the spectral resolution  $\chi_\bullet(|x|)$  is contained in  $\mathcal{M}$  and  $x \in \widetilde{\mathcal{M}}$  if and only if there exists  $\lambda \in \mathbb{R}$  such that  $\tau(\chi_{(\lambda, \infty)}(|x|)) < \infty$ .

For  $p, q \in \mathcal{M}^{\text{pr}}$  we write  $p \sim q$  (*the Murray – von Neumann equivalence*), if  $u^*u = p$  and  $uu^* = q$  for some  $u \in \mathcal{M}$ .

A linear set  $\mathcal{D}$  in  $\mathcal{H}$  is said to be *associated* with  $\mathcal{M}$  if  $u(\mathcal{D}) \subset \mathcal{D}$  for every unitary operator  $u$  in  $\mathcal{M}'$ . If  $\mathcal{D}$  is a closed linear manifold then  $\mathcal{D}$  is associated with  $\mathcal{M}$  if and only if the projection onto  $\mathcal{D}$  lies in  $\mathcal{M}$  [9, p. 403]. For every  $x \in \widetilde{\mathcal{M}}$  the projection onto the closure of the range of  $x$  lies in  $\mathcal{M}$ . It is equal to the left support projection

$$s_l(x) = \wedge \{q \in \mathcal{M}^{\text{pr}} : qx = x\}$$

and  $s_l(x) \sim s_l(x^*)$ .

The two-sided ideal of  $\tau$ -compact operators

$$\widetilde{\mathcal{M}}_0 = \{x \in \widetilde{\mathcal{M}} : \tau(\chi_{(\lambda, \infty)}(|x|)) < \infty \text{ for all } \lambda > 0\}$$

is closed in measure topology [12]. If  $\mathcal{M}$  is  $\mathcal{B}(\mathcal{H})$  with standard trace, then  $\widetilde{\mathcal{M}}_0$  is precisely the ideal of compact operators. Let

$$\mathcal{M}_0^{\text{pr}} = \widetilde{\mathcal{M}}_0 \cap \mathcal{M}^{\text{pr}} = \{p \in \mathcal{M}^{\text{pr}} : \tau(p) < \infty\}.$$

**Definition 1** (cf. [3, p. 114]). A net  $\{x_i\}_{i \in I} \subset \widetilde{\mathcal{M}}$  is said to converge  $\tau$ -locally to  $x \in \widetilde{\mathcal{M}}$  (notation:  $x_i \xrightarrow{\tau l} x$ ) if  $x_i p \xrightarrow{\tau} xp$  for all  $p \in \mathcal{M}_0^{\text{pr}}$ .

**Definition 2** (cf. [3, p. 114]; [5, p. 746]). A net  $\{x_i\}_{i \in I} \subset \widetilde{\mathcal{M}}$  is said to converge weak  $\tau$ -locally to  $x \in \widetilde{\mathcal{M}}$  (notation:  $x_i \xrightarrow{w\tau l} x$ ) if  $px_i p \xrightarrow{\tau} p x p$  for all  $p \in \mathcal{M}_0^{\text{pr}}$ .

It is clear that

$$x_i \xrightarrow{\tau} x \implies x_i \xrightarrow{\tau l} x \implies x_i \xrightarrow{w\tau l} x \text{ for } x_i, x \in \widetilde{\mathcal{M}}.$$

If  $\mathcal{M}$  is  $\mathcal{B}(\mathcal{H})$  with standard trace, then  $\tau$ -local (respectively, weak  $\tau$ -local) convergence coincides with strong-operator (respectively, weak-operator) convergence. If  $\tau(e) < \infty$ , then  $\widetilde{\mathcal{M}}$  consists of all densely defined closed linear operators affiliated with  $\mathcal{M}$  and weak  $\tau$ -local convergence is precisely the convergence in measure topology on  $\widetilde{\mathcal{M}}$ . Moreover, the measure topology is a minimal one in the class of all topologies which are Hausdorff, metrizable, and compatible with the ring structure of  $\widetilde{\mathcal{M}}$  [1, Theorem 2].

## 2. MAIN RESULTS

Further we assume that  $\tau(e) = \infty$ .

**Theorem 1.** Let  $x_i, x \in \widetilde{\mathcal{M}}$ .

1. If  $x_i \xrightarrow{\tau l} x$ , then  $x_i y \xrightarrow{\tau l} xy$  and  $y x_i \xrightarrow{\tau l} yx$  for every fixed  $y \in \widetilde{\mathcal{M}}$ .
2. If  $x_i \xrightarrow{w\tau l} x$ , then  $x_i y \xrightarrow{w\tau l} xy$  and  $y x_i \xrightarrow{w\tau l} yx$  for every fixed  $y \in \widetilde{\mathcal{M}}$ .

**Proof.** Let  $x_i, x, y \in \widetilde{\mathcal{M}}$  and let  $p \in \mathcal{M}_0^{\text{pr}}$ . Since  $s_l(y p) \sim s_l(p y^*) \leq p$ , one has  $s_l(y p) \in \mathcal{M}_0^{\text{pr}}$ .

1. Suppose that  $x_i \xrightarrow{\tau l} x$ . One has

$$y x_i \xrightarrow{\tau l} yx \text{ and } x_i y p = x_i s_l(y p) y p \xrightarrow{\tau} x s_l(y p) y p = x y p,$$

since the multiplication operations  $z \mapsto yz$  ( $\widetilde{\mathcal{M}} \rightarrow \widetilde{\mathcal{M}}$ ) and  $z \mapsto z y p$  ( $\widetilde{\mathcal{M}} \rightarrow \widetilde{\mathcal{M}}$ ) are continuous in the measure topology.

2. One has  $r = p \vee q \in \mathcal{M}_0^{\text{pr}}$  for  $p, q \in \mathcal{M}_0^{\text{pr}}$ , since  $p \vee q - p \sim q - p \wedge q$  [8, p. 105]. By [3, p. 114]  $x_i \xrightarrow{w\tau l} x$  if and only if  $p x_i q \xrightarrow{\tau} p x q$  for all  $p, q \in \mathcal{M}_0^{\text{pr}}$ . Indeed, from  $r x_i r \xrightarrow{\tau} r x r$  it follows that

$$p x_i q = p \cdot r x_i r \cdot q \xrightarrow{\tau} p \cdot r x r \cdot q = p x q.$$

Therefore,

$$p x_i y p = p x_i s_l(y p) y p \xrightarrow{\tau} p x s_l(y p) y p = p x y p.$$

Now the convergence  $yx_i \xrightarrow{w\tau^l} yx$  follows from the fact that the mapping  $z \mapsto z^* (\widetilde{\mathcal{M}} \rightarrow \widetilde{\mathcal{M}})$  is weak  $\tau$ -local continuous and by taking adjoints.

**Theorem 2.** *If  $\{x_i\}_{i \in I} \subset \widetilde{\mathcal{M}}$  is bounded in measure and if  $x_i \xrightarrow{\tau^l} x \in \widetilde{\mathcal{M}}$ , then  $x_i y \xrightarrow{\tau} xy$  for all  $y \in \widetilde{\mathcal{M}}_0$ .*

**Proof. Step 1.** Without loss of generality we may assume that  $y \in \widetilde{\mathcal{M}}_0$  is self-adjoint and non-negative. Indeed, let  $y \in \widetilde{\mathcal{M}}_0$  and  $y^* = u|y^*|$  be the polar decomposition of  $y^*$ . Then  $y = |y^*|u^*$  and from  $x_i|y^*| \xrightarrow{\tau} x|y^*|$  it follows that  $x_i y \xrightarrow{\tau} xy$ , since the multiplication operation  $z \mapsto zu^* (\widetilde{\mathcal{M}} \rightarrow \widetilde{\mathcal{M}})$  is continuous in the measure topology.

**Step 2.** Fix non-negative  $y \in \widetilde{\mathcal{M}}_0$  and  $\varepsilon, \delta > 0$ . A subset  $X$  of  $\widetilde{\mathcal{M}}$  is bounded in measure if and only if for every  $d > 0$  there exists a constant  $c < \infty$  such that  $X \subset U(c, d)$  [8, p. 106]. Let  $n \in \mathbb{N}$  and

$$y_{1,n} = \int_{[0, n^{-1})} \lambda \, d e_\lambda^y, \quad y_{2,n} = \int_{[n^{-1}, n)} \lambda \, d e_\lambda^y, \quad y_{3,n} = \int_{[n, \infty)} \lambda \, d e_\lambda^y.$$

Then  $y = y_{1,n} + y_{2,n} + y_{3,n}$  and for  $z_i = x_i - x$  one has

$$x_i y - xy = z_i y_{1,n} + z_i y_{2,n} + z_i y_{3,n}, \quad i \in I. \quad (1)$$

The set  $\{z_i\}_{i \in I}$  is bounded in measure. There exists a constant  $c > 0$  such that

$$\{z_i\}_{i \in I} \subset U(c, \delta). \quad (2)$$

Let

$$n_1 = \min\{k \in \mathbb{N} : 2\varepsilon k \geq c\}.$$

Since  $\|y_{1,n}\| < n^{-1}$ , one has  $y_{1,n} \xrightarrow{\tau} 0$  as  $n \rightarrow \infty$ . Since

$$\tau(s_l(y_{3,n})) = \tau(\chi_{[n, \infty)}(y)) \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

one has  $y_{3,n} \xrightarrow{\tau} 0$  as  $n \rightarrow \infty$ . Therefore  $y_{1,n} + y_{3,n} \xrightarrow{\tau} 0$  as  $n \rightarrow \infty$ . Then there exists  $m \in \mathbb{N}$  such that

$$y_{1,n} + y_{3,n} \subset U(n_1^{-1}, \delta) \text{ for all } n \geq m. \quad (3)$$

Recall that

$$U(\varepsilon_1, \delta_1)U(\varepsilon_2, \delta_2) \subset U(\varepsilon_1\varepsilon_2, \delta_1 + \delta_2) \text{ for all } \varepsilon_1, \delta_1, \varepsilon_2, \delta_2 > 0 \quad (4)$$

by [8, p. 107], [11, p. 18]. Now by (2) and (3) one has

$$z_i y_{1,n} + z_i y_{3,n} \in U(2\varepsilon, 2\delta) \text{ for all } i \in I, \, n \geq m. \quad (5)$$

**Step 3.** Let  $m$  be as above,  $\lambda_k > 0$  and  $p_k \in \mathcal{M}_0^{\text{pr}}$  ( $k = 1, \dots, j$ ),  $p_k p_l = 0$  for  $k \neq l$ , such that

$$y_{2,m}^2 = \int_{[m^{-1}, m)} \lambda^2 \, d e_\lambda^y \leq \sum_{k=1}^j \lambda_k^2 p_k$$

(one can choose  $p_k$  as spectral projections of  $y$ ). There exists  $z \in \mathcal{M}_1$  such that

$$y_{2,m} = \left( \sum_{k=1}^j \lambda_k p_k \right) \cdot z$$

[4, Chap. 1, Sect. 1, Lemma 2]. Since  $z_i \xrightarrow{\tau^l} 0$ , one has  $z_i p_k \xrightarrow{\tau} 0$  for all  $k = 1, \dots, j$ . Now

$$z_i y_{2,m} = \sum_{k=1}^j \lambda_k z_i p_k z \xrightarrow{\tau} 0,$$

because the multiplication operation  $t \mapsto tz$  ( $\widetilde{\mathcal{M}} \rightarrow \widetilde{\mathcal{M}}$ ) is continuous in the measure topology. Therefore, there exists  $i_0 \in I$  such that

$$z_i y_{2,m} \in U(\varepsilon, \delta) \text{ for all } i \in I, i \geq i_0. \quad (6)$$

**Step 4.** Recall that

$$U(\varepsilon_1, \delta_1) + U(\varepsilon_2, \delta_2) \subset U(\varepsilon_1 + \varepsilon_2, \delta_1 + \delta_2) \text{ for all } \varepsilon_1, \delta_1, \varepsilon_2, \delta_2 > 0 \quad (7)$$

by [8, p. 107], [11, p. 18]. The assertion of Theorem 2 follows from (1), (5) and (6), since

$$x_i y - x y \in U(3\varepsilon, 3\delta) \text{ for all } i \in I, i \geq i_0.$$

**Theorem 3.** Let  $x, y, x_i, y_i \in \widetilde{\mathcal{M}}$  and let a set  $\{x_i\}_{i \in I}$  be bounded in measure. If  $x_i \xrightarrow{\tau^l} x$  and  $y_i \xrightarrow{\tau^l} y$ , then  $x_i y_i \xrightarrow{\tau^l} xy$ .

**Proof.** For every  $p \in \mathcal{M}_0^{\text{pr}}$  one has

$$x_i y_i p - x y p = x_i (y_i p - y p) + (x_i - x) y p, \quad i \in I. \quad (8)$$

Fix  $\varepsilon, \delta > 0$ . By assumption of the theorem, there exists a constant  $c > 0$  such that

$$\{x_i\}_{i \in I} \subset U(c, \delta). \quad (9)$$

Since  $y_i p - y p \xrightarrow{\tau} 0$ , there exists  $i_1 \in I$  such that

$$y_i p - y p \in U(2\varepsilon c^{-1}, \delta) \text{ for all } i \in I, i \geq i_1. \quad (10)$$

Now by (9), (10) and (4) one has

$$x_i (y_i p - y p) \in U(2\varepsilon, 2\delta) \text{ for all } i \in I, i \geq i_1. \quad (11)$$

Since  $x_i - x \xrightarrow{\tau_l} 0$  and  $yp \in \widetilde{\mathcal{M}}_0$ , it follows by Theorem 2 that there exists  $i_2 \in I$  such that

$$(x_i - x)yp \in U(\varepsilon, \delta) \text{ for all } i \in I, i \geq i_2. \quad (12)$$

There exists  $i_0 \in I$  such that  $i_0 \geq i_1$  and  $i_0 \geq i_2$ . Now by (8), (11), (12) and (7) one has

$$x_i y_i p - xyp \in U(3\varepsilon, 3\delta) \text{ for all } i \in I, i \geq i_0.$$

This proves the theorem.

**Example 1.** If  $\mathcal{M}$  is abelian, then the weak  $\tau$ -local and  $\tau$ -local convergencies on  $\widetilde{\mathcal{M}}$  coincides with the familiar convergence locally in measure (i.e., in other words, convergence in measure on every set of finite measure). The boundedness condition for  $\{x_i\}_{i \in I}$  cannot be omitted in Theorem 2. Indeed, let  $\Omega = (0, \infty)$  equipped with the Lebesgue measure  $\mu$ . Define the functions

$$y(t) = \min\{1, t^{-1}\}; \quad x_n(t) = t \chi_{[n, 2n]}(t) \quad (t \in (0, \infty), n \in \mathbb{N}).$$

Then

- i)  $x_n \xrightarrow{\tau_l} 0$  as  $n \rightarrow \infty$ ;
- ii)  $\{x_n\}_{n=1}^\infty$  is not bounded in measure;
- iii)  $y \in \mathcal{M}_0 \cap \mathcal{M}_1$ ;
- iv) since  $(x_n y)(t) = \chi_{[n, 2n]}(t)$  for every  $t \in (0, \infty)$ ,  $n \in \mathbb{N}$ ,  $x_n y$  does not converge in measure topology.

**Example 2.** If  $\mathcal{M}$  is  $\mathcal{B}(\mathcal{H})$  with standard trace, then Theorem 2 for sequences is a "Basic lemma" of the projection methods [2, pp. 18–19] (the boundedness condition for  $\{x_n\}_{n=1}^\infty$  follows from *the principle of uniform boundedness*):

*If  $y$  is compact and  $x_n \rightarrow x$  strongly, then  $x_n y \rightarrow xy$  uniformly, i.e.  $\|x_n y - xy\| \rightarrow 0$  as  $n \rightarrow \infty$ .*

Theorem 3 means that the mapping

$$(x, y) \mapsto xy : (\mathcal{B}(\mathcal{H})_1 \times \mathcal{B}(\mathcal{H})) \rightarrow \mathcal{B}(\mathcal{H})$$

is strong-operator continuous [7, pp. 115–117].

**Remark.** The second part of Theorem 1 was already used in [6] and [10].

## REFERENCES

- [1] Bikchentaev A.M. *On minimality of the topology of convergence in measure on finite von Neumann algebras*, Kolmogorov and contemporary mathematics. Abstracts of international conference in commemoration of the centennial of A.N.Kolmogorov (Moscow, June 16–21, 2003). Moscow. Univ., 2003. 915 p., 139–140.
- [2] Böttcher A. ... [et al.]; edited by P. Lancaster, *Lectures on operator theory and its applications*. Fields Institute monographs. Providence, Rhode Island, Amer. Math. Soc. 1996. 340 p.
- [3] Ciach L.J., *Some remarks on the convergence in measure and on a dominated sequence of operators measurable with respect to a semifinite von Neumann algebra*, Coll. Math. **LV** (1988), 109–121.
- [4] Dixmier J. *Les algebres d'operateurs dans l'espace Hilbertien (algebres de von Neumann)*. Paris, Gauthier-Villars, 1969. 367 p.
- [5] Dodds P.G., Dodds T.K., and de Pagter B. *Non-commutative Köthe duality*, Trans. Amer. Math. Soc. **339** (1993), 717–750.
- [6] Dodds P.G., Dodds T.K., Sukochev F.A. and Tikhonov O.E. *A non-commutative Yosida-Hewitt theorem and applications to convex sets of measurable operators closed locally in measure*, in preparation.
- [7] Kadison R.V. and Ringrose J.R. *Fundamentals of the theory of operator algebras*. V. I. New York – London – Paris: Academic Press, 1983. 398 p.
- [8] Nelson E. *Notes on non-commutative integration*, J. Funct. Anal. **15** (1974), 103–116.
- [9] Segal I. *A non-commutative extension of abstract integration*, Ann. Math. **57** (1953), 401–457.
- [10] Skvortsova G.Sh. *Weak sequential completeness of factor spaces of the space of integrable operators*, Russian Mathematics (Iz. VUZ) **45** (2002), 68–71.
- [11] Terp M.  *$L^p$  spaces associated with von Neumann algebras*, Notes, Copenhagen Univ. (1981). 100 p.
- [12] Yeadon F.J. *Non-commutative  $L^p$ -spaces*, Math. Proc. Cambridge Phil. Soc. **77** (1975), 91–102.

RESEARCH INSTITUTE OF MATHEMATICS AND MECHANICS, KAZAN STATE UNIVERSITY, UNIVERSITETSKAYA STR. 17, KAZAN:420008, RUSSIA

*E-mail address:* Airat.Bikchentaev@ksu.ru

Received January 8, 2004