

On the index of certain standard congruence subgroups

Thomas Morgan

ABSTRACT. For an epimorphism of the free group on two generators onto a finite group G , one can associate a finite index subgroup of the automorphism group of the free group called the standard congruence subgroup. We calculate the index of this group when G is a non-abelian semi-direct product of cyclic groups of prime order.

CONTENTS

1. Introduction	848
2. Independence of the choice of π	849
2.1. A description of $\text{Aut}(G)$	849
2.2. Independence of the choice of π	851
3. Proof of Theorem 1.1	852
3.1. Image of primitive elements	852
3.2. Proof of Theorem 1.1	853
References	854

1. Introduction

There is a well-known surjective representation of the automorphism group of a free group $\rho_0 : \text{Aut}(F_n) \rightarrow \text{Aut}(F_n/[F_n, F_n]) \cong \text{GL}_n(\mathbb{Z})$. The kernel of this representation is called the *Torelli subgroup*, denoted $IA(F_n)$, and the subgroup of $\text{Aut}(F_n)$ whose elements have determinant 1 under ρ_0 is called the *special automorphism group*, denoted $\text{Aut}^+(F_n)$. While this linear representation of $\text{Aut}^+(F_n)$ is well studied, only a few other representations were studied until 2006, when Grunewald and Lubotzky[GL09] published a paper detailing the construction of a family of virtual linear representations of $\text{Aut}(F_n)$ indexed by finite groups G and surjective homomorphisms $\pi : F_n \rightarrow G$. This gave rise to a generalization of the Torelli subgroup different from the Johnson filtration and proved that $\text{Aut}(F_3)$ is large, which implies that it does not have Kazhdan’s property (T).

Received April 23, 2018.

2010 *Mathematics Subject Classification.* 20F28.

Key words and phrases. automorphism groups, free groups, congruence subgroups.

In constructing these representations, Grunewald and Lubotzky used the subgroup $\Gamma(G, \pi) \leq \text{Aut}(F_n)$. This subgroup is called the *standard congruence subgroup* of $\text{Aut}(F_n)$ associated to G and π . The subgroup is defined as follows: let $R := \ker(\pi)$. Then the action of F_n on R by conjugation leads to an action of G on the relation module $\bar{R} := R/[R, R]$. Define $\Gamma(G, \pi) := \{\varphi \in \text{Aut}(F_n) \mid \varphi(R) = R, \varphi \text{ induces identity on } F_n/R \cong G\}$. This is exactly the G -equivariant automorphisms under this action. It is also analogous to congruence subgroups, which are extensively studied for arithmetic groups.

Following Grunewald and Lubotzky’s paper, Appel and Ribnere [AR09] began a more systematic study of these standard congruence subgroups in the case where $n = 2$. First they restricted themselves to $\Gamma^+(G, \pi) = \Gamma(G, \pi) \cap \text{Aut}^+(F_2)$. Then they computed the index $[\text{Aut}^+(F_2) : \Gamma^+(G, \pi)]$ for G abelian or dihedral. In doing so, and with some further analysis, they gave some partial results to the congruence subgroup problem for $\text{Aut}^+(F_2)$. Appel and Ribnere also posed a conjecture stating the index when G is the non-abelian semidirect product of two cyclic groups of prime order. We prove their conjecture.

Theorem 1.1. *Let G be the non-abelian semidirect product of two cyclic groups, $G = \mathbb{Z}/p\mathbb{Z} \rtimes \mathbb{Z}/q\mathbb{Z}$, where p and q are primes with $p \equiv_q 1$. Then*

$$[\text{Aut}^+(F_2) : \Gamma^+(G, \pi)] = |G| \cdot [SL_2(\mathbb{Z}) : \Gamma_1(q)] = pq(q^2 - 1).$$

Here $\Gamma_1(q) := \left\{ \begin{bmatrix} \alpha & \beta \\ \delta & \epsilon \end{bmatrix} \in SL_2(\mathbb{Z}) \mid \delta \equiv_q 0, \epsilon \equiv_q 1 \right\}$. Note that this is true independent of the choice of π . We prove this in Section 2. The second equality follows from the study of congruence subgroups (see for example [DS05]). Then in Section 3, we prove the first equality by using the primitive elements constructed in [OZ81] to construct enough automorphisms in $\Gamma^+(G, \pi)$ to prove that $\rho_0(\Gamma^+(G, \pi)) = \Gamma_1(q)$. The equality then follows from the following proposition from [AR09].

Proposition 1.2 (Appel, Ribnere). *Let $\pi : F_2 \rightarrow G$ be an epimorphism of F_2 onto a finite group G . Then*

$$[\text{Aut}^+(F_2) : \Gamma^+(G, \pi)] = [SL_2(\mathbb{Z}) : \rho_0(\Gamma^+(G, \pi))] \cdot [G : Z(G)].$$

Here $\rho_0 : \text{Aut}(F_n) \rightarrow \text{Aut}(F_n/[F_n, F_n]) \cong GL_n(\mathbb{Z})$ is the representation introduced earlier.

2. Independence of the choice of π

2.1. A description of $\text{Aut}(G)$. Let $p, q \in \mathbb{N}$ be primes such that $p \equiv_q 1$. Then the non-abelian semi-direct product $G := \mathbb{Z}/p\mathbb{Z} \rtimes \mathbb{Z}/q\mathbb{Z}$ has the following presentation:

$$G = \langle a, b \mid a^p = b^q = 1, bab^{-1} = a^\lambda \rangle$$

for some $1 < \lambda < p$ and $\lambda^q \equiv_p 1$ ($\lambda = 1$ would be the abelian case). Indeed, let $\mathbb{Z}/p\mathbb{Z} = \langle a \rangle$ and $\mathbb{Z}/q\mathbb{Z} = \langle b \rangle$. Then the above presentation comes from the outer semi-direct product associated with the homomorphism $\varphi : \mathbb{Z}/q\mathbb{Z} \rightarrow \text{Aut}(\mathbb{Z}/p\mathbb{Z})$ such that $\varphi(b)(a) = a^\lambda$.

Let $F_2 = \langle x, y \rangle$ be the free group on two generators. To see that the index we wish to calculate is independent of the choice of $\pi : F_2 \rightarrow G$, we look at an action of $\text{Aut}^+(F_2)$ on the following set. We define

$$\mathbf{R}_2(G) := \{\ker(\pi) \mid \pi : F_2 \rightarrow G \text{ is an epimorphism}\}.$$

We can define an action of $\text{Aut}^+(F_2)$ on $\mathbf{R}_2(G)$ by

$$\varphi.R := \varphi(R) \text{ for } \varphi \in \text{Aut}^+(F_2), R \in \mathbf{R}_2(G).$$

If the action is transitive, then $[\text{Aut}^+(F_2) : \Gamma^+(G, \pi)]$ is independent of the choice of π . Indeed, let $\pi, \pi' : F_2 \rightarrow G$ be epimorphisms, and assume that there is some $\varphi \in \text{Aut}^+(F_2)$ such that $\varphi.\ker(\pi) = \ker(\pi')$. Since $\varphi.\ker(\pi) = \ker(\pi \circ \varphi^{-1})$, we have that $\Gamma^+(G, \pi') = \{\varphi \circ \psi \circ \varphi^{-1} \mid \psi \in \Gamma^+(G, \pi)\}$. Since $\Gamma^+(G, \pi)$ and $\Gamma^+(G, \pi')$ are conjugate subgroups of $\text{Aut}^+(F_2)$, we conclude that $[\text{Aut}^+(F_2) : \Gamma^+(G, \pi)] = [\text{Aut}^+(F_2) : \Gamma^+(G, \pi')]$.

In order to show that this action is transitive, it is helpful to first understand the automorphisms of the group G . To that effect, we prove the following lemma.

Lemma 2.1. *For each $0 < i < p, 0 \leq j < p$ there is a unique automorphism $\varphi_{i,j} : G \rightarrow G$ such that $\varphi(a) = a^i, \varphi(b) = a^j b$. Moreover $\text{Aut}(G) = \{\varphi_{i,j} \mid 0 < i < p, 0 \leq j < p\}$.*

Proof. Let $0 < i < p, 0 \leq j < p$. Since $\{a, b\}$ is a generating set of G , we can define $\varphi_{i,j}$ on $\{a, b\}$ as above and extend to a homomorphism in a unique way. To see this is a well-defined homomorphism, we will check that it satisfies the relations. First, we have that

$$\varphi(a)^p = (a^i)^p = a^{ip} = (a^p)^i = 1^i = 1.$$

Since $p \nmid (1 - \lambda)$ and $\lambda^q \equiv_p 1$, it follows that

$$\varphi(b)^q = (a^j b)^q = a^j a^{j\lambda} \dots a^{j\lambda^{q-1}} b^q = a^r,$$

where

$$r = \sum_{i=0}^{q-1} j\lambda^i = j \sum_{i=0}^{q-1} \lambda^i = j \left(\frac{1 - \lambda^q}{1 - \lambda} \right) \equiv_p 0.$$

Finally, we have that

$$\varphi(b)\varphi(a)\varphi(b)^{-1} = a^j b a^i b^{-1} a^{-j} = a^j a^{\lambda^i} a^{-j} = (a^i)^\lambda = \varphi(a)^\lambda.$$

Thus φ is a homomorphism.

To see this map is surjective, note that a^i is a generator of $\langle a \rangle$ for $0 < i < p$. It follows that $a \in \text{Im}(\varphi_{i,j})$. Then $a, a^j b \in \text{Im}(\varphi_{i,j}) \implies b \in \text{Im}(\varphi_{i,j})$. This shows that $\text{Im}(\varphi_{i,j})$ contains a generating set, so $\varphi_{i,j}$ is surjective. Because G is finite, this is enough to show that $\varphi_{i,j}$ is an automorphism.

Now let H denote the set $\{\varphi_{i,j} \mid 0 < i < p, 0 \leq j < p\}$. We have shown that $H \subset \text{Aut}(G)$. Thus it remains to show the reverse inclusion.

Let $\varphi \in \text{Aut}(G)$. Since $\langle a \rangle$ contains all of the elements of order p in G ,

$$\begin{aligned} \varphi(a) &= a^i \text{ for some } 0 < i < p. \\ \varphi(b) &= a^j b^k \text{ for some } 0 \leq j < p, 0 < k < q. \end{aligned}$$

On the one hand,

$$\varphi(bab^{-1}) = \varphi(a^\lambda) = a^{i\lambda}.$$

On the other hand,

$$\varphi(bab^{-1}) = \varphi(b)\varphi(a)\varphi(b)^{-1} = a^j b^k a^i b^{-k} a^{-j} = a^j a^{i\lambda^k} b^k b^{-k} a^{-j} = a^{i\lambda^k}.$$

Thus $a^{i\lambda} = a^{i\lambda^k}$. Since a is of order p and p does not divide i , it follows that $\lambda^{k-1} \equiv_p 1$. We know that λ is of order q , so $q \mid (k-1)$. But $0 < k < q$, so $k = 1$. Thus $\text{Aut}(G) \leq H$. □

2.2. Independence of the choice of π . We will now show that the action of $\text{Aut}^+(F_2)$ on $\mathbf{R}_2(G)$ defined above is transitive. This will mean that the index $[\text{Aut}^+(F_2) : \Gamma^+(G, \pi)]$ is independent of the choice of π . Thus we will be able to compute the index using the epimorphism

$$\begin{aligned} \pi_0 : F_2 &\rightarrow G \\ x &\mapsto a \\ y &\mapsto b \end{aligned}$$

Lemma 2.2. *The action of $\text{Aut}^+(F_2)$ on the set $\mathbf{R}_2(G)$ is transitive.*

Proof. Let $\pi : F_2 \rightarrow G$ be an arbitrary epimorphism of F_2 onto G . Then for some $i, j, k, \ell \in \mathbb{Z}$, we have $\pi(x) = a^i b^j$, $\pi(y) = a^k b^\ell$. Let $\beta : G \rightarrow \mathbb{Z}/q\mathbb{Z}$, $\beta(a^m b^n) = n$. Then $\beta \circ \pi$ is an epimorphism of F_2 onto $\mathbb{Z}/q\mathbb{Z}$. The following diagram commutes:

$$\begin{array}{ccccc} F_2 & \xrightarrow{\alpha} & (\mathbb{Z}/q\mathbb{Z})^2 & \xrightarrow{\delta} & (\mathbb{Z}/q\mathbb{Z}) \\ & & \searrow & \nearrow & \\ & & & \beta \circ \pi & \end{array}$$

where $\alpha(x) = (1, 0)$, $\alpha(y) = (0, 1)$, and $\delta(m, n) = mj + n\ell$. Since $\delta \circ \alpha$ is surjective, there is an element $(m, n) \in (\mathbb{Z}/q\mathbb{Z})^2$ such that $\delta(m, n) = 1$. Furthermore, since δ is not injective, there is a non-zero element (u, v) such

that $\delta(u, v) = 0$. It is clear that (m, n) and (u, v) are linearly independent, so $d := \begin{vmatrix} u & m \\ v & n \end{vmatrix} \not\equiv_q 0$. Thus we can choose a $\tilde{d} \in \mathbb{Z}$ such that $d\tilde{d} \equiv_q 1$. The vector $(\tilde{d}u, \tilde{d}v)$ is in the kernel of δ since (u, v) is, and $M := \begin{pmatrix} \tilde{d}u & m \\ \tilde{d}v & n \end{pmatrix} \in \text{Sl}_2(\mathbb{Z}/q\mathbb{Z})$.

Choose a matrix $N \in \text{Sl}_2(\mathbb{Z})$ such that $N \equiv_q M$. Let $\rho_0 : \text{Aut}(F_n) \rightarrow \text{Aut}(F_n/[F_n, F_n]) \cong \text{GL}_n(\mathbb{Z})$ be the homomorphism described in the introduction. Because ρ_0 is surjective, we may choose an automorphism $\phi \in \text{Aut}^+(F_2)$ such that $\rho_0(\phi) = N$. It follows that $\delta \circ \alpha \circ \phi(x) = 0$, $\delta \circ \alpha \circ \phi(y) = 1$. Thus $\pi \circ \phi(x) = a^g$ and $\pi \circ \phi(y) = a^hb$ for some $g, h \in \mathbb{Z}$. By Lemma 2.1, $\varphi_{g,h}^{-1} \circ \pi \circ \phi = \pi_0$. It follows that $\ker(\pi)$, $\ker(\pi_0)$ lie in the same $\text{Aut}^+(F_2)$ orbit. \square

3. Proof of Theorem 1.1

3.1. Image of primitive elements. Let p, q be as above. Now that we know the action of $\text{Aut}^+(F_2)$ on $\mathbf{R}_2(G)$ is transitive, we need to show that

$$\rho_0(\Gamma^+(G, \pi_0)) = \Gamma_1(q). \text{ Here } \Gamma_1(q) := \left\{ \begin{bmatrix} \alpha & \beta \\ \delta & \epsilon \end{bmatrix} \in \text{Sl}_2(\mathbb{Z}) \mid \delta \equiv_q 0, \epsilon \equiv_q 1 \right\}.$$

To do this, we use the description of primitive elements from [OZ81] to construct elements of $\Gamma^+(G, \pi_0)$. In Section 2 of [OZ], given $\alpha, \delta \in \mathbb{Z}^+$ such that $\gcd(\alpha, \delta) = 1$, Osbourne and Zieschang outline a geometric construction of a primitive element $v_{\alpha, \delta}$ containing α copies of x and δ copies of y which we reproduce here. Draw a directed line segment from $(0, 0)$ to (α, δ) . We use this line segment to generate a word in F_2 . Starting at $(0, 0)$, every time the segment passes a vertical integer grid line write an x , and every time the segment passes a horizontal integer grid line write a y . Call the resulting word $v'_{\alpha, \delta}$. Define $v_{\alpha, \delta} := xyv'_{\alpha, \delta}$. Lemma 2.3 of [OZ81] shows that $v_{\alpha, \delta}$ is primitive by relating it to a construction earlier in the paper which is algebraically shown to be primitive. We use these primitive elements and their geometric construction in the following lemma:

Lemma 3.1. *Let $\alpha, \delta \in \mathbb{Z}^+$ such that $\gcd(\alpha, \delta) = 1$ and $q|\delta$. Then $\pi_0(v_{\alpha, \delta}) = a^z b^\delta$ for some $z \in \mathbb{Z}$ which depends only on $\alpha \pmod{q}$.*

Proof. Note that the x 's in $v'_{\alpha, \delta}$ correspond to points $(i, i\delta/\alpha)$ for $0 < i < \alpha$. Given each x in $v'_{\alpha, \delta}$ we want to consider what it will take to move it to the front of the word. That is, we want to count the number of y 's that occur before each x . For the i th x , this is precisely the integer part of $i\delta/\alpha$. This is equal to $(i\delta - r_i)/\alpha$ where $0 \leq r_i < \alpha$ is the remainder when $i\delta$ is divided by α . In terms of the image of $v'_{\alpha, \delta}$ under π , we only care about the number of y 's mod q . Since $\gcd(\alpha, \delta) = 1$ and $q|\delta$, we have $\gcd(q, \alpha) = 1$. Thus there exists an $\tilde{\alpha} \in \mathbb{Z}$ such that $\alpha\tilde{\alpha} \equiv_q 1$. It follows that $(i\delta - r_i)/\alpha \equiv_q -\tilde{\alpha}r_i$ since $q|\delta$. Therefore, after commuting the i th a in the image of $v'_{\alpha, \delta}$ to the front of

the word it becomes $a^{\lambda^{-\tilde{\alpha}r_i}}$. Thus the image of $v'_{\alpha,\delta}$ under π is $a^s b^{\delta-1}$ where

$$s = \sum_{i=1}^{\alpha-1} \lambda^{-\tilde{\alpha}r_i} = \sum_{i=1}^{\alpha-1} (\lambda^{-\tilde{\alpha}})^{r_i}.$$

Consider the list of remainders $(r_1, r_2, \dots, r_{\alpha-1})$. By the above considerations, this list completely determines the power of a in $\pi(v'_{\alpha,\delta})$. Since $\gcd(\alpha, \delta) = 1$, $[\delta] \in \mathbb{Z}/\alpha\mathbb{Z}$ is a generator. Furthermore, by definition $[r_i] = i[\delta]$ in $\mathbb{Z}/\alpha\mathbb{Z}$. Thus up to reordering, $(r_1, r_2, \dots, r_{\alpha-1}) = (1, 2, \dots, \alpha - 1)$. As this is true for any choice of δ meeting our requirements, for fixed α the power of a in $v'_{\alpha,\delta}$ and hence $v_{\alpha,\delta}$ is independent of our choice of δ .

Now fix δ . If $\alpha_2 = \alpha_1 + q$, then we get two different lists of remainders. They are $(1, 2, \dots, \alpha_1 - 1)$ and $(1, 2, \dots, \alpha_2 - 1) = (1, 2, \dots, q, q + 1, q + 2, \dots, q + \alpha_1 - 1)$. Let $\lambda' := \lambda^{-\tilde{\alpha}_2}$. Note that $\alpha_1 \equiv_q \alpha_2 \implies \tilde{\alpha}_1 \equiv_q \tilde{\alpha}_2$. Thus $\lambda' = \lambda^{-\tilde{\alpha}_1}$. Plugging this into our formula for the power of a in $\pi(v'_{\alpha,\delta})$ and noting that $(\lambda')^q \equiv_p 1$, we get

$$\begin{aligned} \sum_{i=1}^{\alpha_2-1} (\lambda')^{r_i} &= \sum_{i=1}^q (\lambda')^{r_i} + \sum_{i=q+1}^{q+\alpha_1-1} (\lambda')^{r_i} \\ &\equiv_p \lambda' \left(\frac{1 - (\lambda')^q}{1 - \lambda'} \right) + \sum_{i=1}^{\alpha_1-1} (\lambda')^{r_i} \\ &\equiv_p \sum_{i=1}^{\alpha_1-1} (\lambda')^{r_i}. \end{aligned}$$

Thus $\pi_0(v'_{\alpha_1,\delta}) = \pi_0(v'_{\alpha_2,\delta})$. By induction, we see that $\pi_0(v'_{\alpha,\delta})$ and hence $\pi_0(v_{\alpha,\delta})$ only depends upon $\alpha \pmod q$. □

3.2. Proof of Theorem 1.1. With Proposition 1.2 and our lemmas, we are ready to prove Theorem 1.1.

Proof of Theorem 1.1. By Lemma 2.2, we may assume that $\pi = \pi_0$. Noting that $Z(G) = 1$ so $[G : Z(G)] = |G|$, by Proposition 1.2 it suffices to show that $\rho_0(\Gamma^+(G, \pi)) = \Gamma_1(q)$. Since $\rho_0(\Gamma^+(G, \pi)) \leq \rho_0(\Gamma^+(G^{\text{ab}}, \bar{\pi})) = \Gamma_1(q)$ where $f \circ \pi = \bar{\pi}$ and $f : G \rightarrow G^{\text{ab}}$ is the abelianization map, it remains to show $\rho_0(\Gamma^+(G, \pi)) \geq \Gamma_1(q)$.

Let $A = \begin{bmatrix} \alpha & \beta \\ \delta & \epsilon \end{bmatrix} \in \Gamma_1(q)$. Then let φ be the automorphism determined by

$$\begin{aligned} \varphi : F_2 &\rightarrow F_2 \\ x &\mapsto v_{\alpha,\delta} \\ y &\mapsto v_{\beta,\epsilon} \end{aligned}$$

By Theorem 1.2 of [OZ81], we have that $v_{\alpha,\delta}$ and $v_{\beta,\epsilon}$ generate F_2 . Thus $\varphi \in \text{Aut}^+(F_2)$. Let $\pi(v_{\beta,\epsilon}) = a^j b$. It must be of this form since $\epsilon \equiv_q 1$ and b

has order q .

First assume $\alpha, \delta > 0$. Since $p \nmid (\lambda - 1)$, there exists an $\ell \in \mathbb{Z}$ such that $(\lambda - 1)\ell \equiv_p j$. Then

$$\pi(x^\ell v_{\beta, \epsilon} x^{-\ell}) = a^\ell a^j b a^{-\ell} = a^\ell a^j a^{-\lambda \ell} b = a^{(1-\lambda)\ell + j} b = b$$

By Lemma 3.1, the a exponent of $\pi(v_{\alpha, \delta})$ only depends upon $\alpha \bmod q$. But $\alpha \equiv_q 1$ and $v_{1, \delta} = xy^\delta$ by direct computation. Thus

$$\pi(x^\ell v_{\alpha, \delta} x^{-\ell}) = a^\ell a a^{-\ell} = a$$

This shows that $c \circ \varphi \in \Gamma^+(G, \pi)$ where c is conjugation by x^ℓ . Since $\rho_0(c \circ \varphi) = A$ by construction, this shows $A \in \rho_0(\Gamma^+(G, \pi))$.

We now consider the case where α and δ are arbitrary. By the above argument, $\begin{bmatrix} 1 & 0 \\ q & 1 \end{bmatrix} \in \rho_0(\Gamma^+(G, \pi))$. Furthermore, for the correct choice of ℓ as above, the automorphism

$$\begin{aligned} \psi : F_2 &\rightarrow F_2 \\ x &\mapsto x \\ y &\mapsto x^{\ell+1} y x^{-\ell} \end{aligned}$$

is in $\Gamma^+(G, \pi)$, and $\rho_0(\psi) = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \in \rho_0(\Gamma^+(G, \pi))$. For general α, δ , we may write A as a product of powers of these matrices and some $A' = \begin{bmatrix} \alpha' & \beta' \\ \delta' & \epsilon' \end{bmatrix} \in \Gamma_1(q)$ with $\alpha', \delta' > 0$. This shows that $A \in \rho_0(\Gamma^+(G, \pi))$. \square

References

- [AR09] APPEL, DANIEL; RIBNERE, EVIJA. On the index of congruence subgroups of $\text{Aut}(F_n)$. *J. Algebra* **321** (2009), no. 10, 2875–2889. [MR2512632](#) (2010d:20039), [Zbl 1178.20036](#), [arXiv:0804.2578](#), doi: [10.1016/j.jalgebra.2009.01.022](#). 849
- [DS05] DIAMOND, FRED; SHURMAN, JERRY. A first course in modular forms. Graduate Texts in Mathematics, 228. *Springer-Verlag, New York*, 2005. xvi+436 pp. ISBN: 0-387-23229-X. [MR2112196](#) (2006f:11045), [Zbl 1062.11022](#), doi: [10.1007/978-0-387-27226-9](#). 849
- [GL09] GRUNEWALD, FRITZ; LUBOTZKY, ALEXANDER. Linear representations of the automorphism group of a free group. *Geom. Funct. Anal.* **18** (2009), no. 5, 1564–1608. [MR2481737](#) (2010i:20039), [Zbl 1175.20028](#), [arXiv:math/0606182](#), doi: [10.1007/s00039-009-0702-2](#). 848
- [LS01] LYNDON, ROGER C.; SCHUPP, PAUL E. Combinatorial group theory. Reprint of the 1977 edition. Classics in Mathematics. *Springer-Verlag, Berlin*, 2001. xiv+339 pp. ISBN: 3-540-41158-5. [MR1812024](#) (2001i:20064), [Zbl 0997.20037](#), doi: [10.1007/978-3-642-61896-3](#).
- [OZ81] OSBORNE, RICHARD P.; ZIESCHANG, HEINER. Primitives in the free group on two generators. *Invent. Math.* **63** (1981), no. 1, 17–24. [MR0608526](#) (82i:20042), [Zbl 0438.20017](#), doi: [10.1007/BF01389191](#). 849, 852, 853

(Thomas Morgan) DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF OKLAHOMA,
NORMAN, OK 73019, USA

tmorgan@ou.edu

This paper is available via <http://nyjm.albany.edu/j/2018/24-38.html>.