

The Musielak-Orlicz Herz spaces

Baohua Dong, Yu Li and Jingshi Xu

ABSTRACT. Inspired by the fact that the boundedness of certain sublinear operators on $L^p(\mathbb{R}^n)$ implies their boundedness on the Herz spaces, we obtain that the boundedness of certain sublinear operators on Musielak-Orlicz spaces implies their boundedness on Musielak-Orlicz Herz spaces.

CONTENTS

1. Introduction	1287
2. The proof of Theorems 1.2 and 1.3	1291
3. The proof of Theorems 1.5 and 1.6	1296
Acknowledgement	1298
References	1298

1. Introduction

Herz spaces can be traced back to the work [3] where Beurling studied certain convolution algebra by introduced the space A_q . It is the original version of the non-homogeneous Herz space. The general Herz spaces are introduced in [16] by Herz. Many researchers made their contributions to the theory of Herz spaces at that time, for example, Feichtinger ([7]), Flett ([8]), and Baernstein and Sawyer ([2]), as well as García-Cuerva and Herrero ([11, 12]). Before presenting the definition of Herz spaces, we need the necessary notations. Let $B_k := \{x \in \mathbb{R}^n : |x| \leq 2^k\}$ and $A_k := B_k \setminus B_{k-1}$ for $k \in \mathbb{Z}$. Denote $\chi_k := \chi_{A_k}$ for $k \in \mathbb{Z}$, $\tilde{\chi}_k := \chi_k$ if $k \in \mathbb{N} \setminus \{0\}$ and $\tilde{\chi}_0 := \chi_{B_0}$, where χ_{A_k} is the characteristic function of A_k . Let $\alpha \in \mathbb{R}$, $p, q \in (0, \infty]$. The homogeneous Herz space $\dot{K}_p^{\alpha, q}(\mathbb{R}^n)$ is defined by

$$\dot{K}_p^{\alpha, q}(\mathbb{R}^n) := \left\{ f \in L_{\text{loc}}^p(\mathbb{R}^n \setminus \{0\}) : \|f\|_{\dot{K}_p^{\alpha, q}(\mathbb{R}^n)} < \infty \right\},$$

Received July 30, 2023.

2020 *Mathematics Subject Classification.* Primary 42B35, 42B20.

Key words and phrases. Sublinear operator; Musielak-Orlicz Herz space.

The corresponding author Jingshi Xu is supported by the National Natural Science Foundation of China (Grant No. 12161022).

where

$$\|f\|_{\dot{K}_p^{\alpha,q}(\mathbb{R}^n)} := \left\| \left\{ 2^{k\alpha} \|f\chi_k\|_{L^p} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^q} = \left\{ \sum_{k \in \mathbb{Z}} 2^{k\alpha q} \|f\chi_k\|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}}.$$

The non-homogeneous Herz space $K_p^{\alpha,q}(\mathbb{R}^n)$ is defined by

$$K_p^{\alpha,q}(\mathbb{R}^n) := \left\{ f \in L_{\text{loc}}^p(\mathbb{R}^n) : \|f\|_{K_p^{\alpha,q}(\mathbb{R}^n)} < \infty \right\},$$

where

$$\|f\|_{K_p^{\alpha,q}(\mathbb{R}^n)} := \left\| \left\{ 2^{k\alpha} \|f\tilde{\chi}_k\|_{L^p} \right\}_{k \in \mathbb{N}} \right\|_{\ell^q} = \left\{ \sum_{k=0}^{\infty} 2^{k\alpha q} \|f\tilde{\chi}_k\|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}}.$$

The usual modifications are made when $p = \infty$ or $q = \infty$.

Herz spaces are quite useful in harmonic analysis. They can be used to characterize the multipliers on Hardy spaces ([2]). They are natural spaces to consider the problems related to multidimensional θ -summation and θ -means of Fourier transforms ([9, 10, 23]), as in $L^p(\mathbb{R}^n)$ spaces and Wiener amalgam spaces $W(L^p, \ell_\infty)(\mathbb{R}^n)$ ([37, 40]). When $p = q, \alpha = \beta/p, \beta \in \mathbb{R}$, the Herz space $\dot{K}_p^{\beta/p,p}(\mathbb{R}^n)$ is exactly the weighted Lebesgue space $L^p(\mathbb{R}^n, |x|^\beta)$. One of the motivations to study the boundedness of sublinear operators in Herz spaces is due to the enlightening works of Stein ([36]), Hofmann ([17]), and Soria and Weiss ([38]). These results show that the boundedness on the space $L^p(\mathbb{R}^n, |x|^\beta)$ can be deduced from its boundedness on $L^p(\mathbb{R}^n)$ for sublinear operators T satisfying the condition (1):

$$|Tf(x)| \lesssim \int_{\mathbb{R}^n} \frac{|f(y)|}{|x-y|^n} dy, \quad f \in L^1(\mathbb{R}^n) \text{ with compact support, } x \notin \text{supp } f.$$

Here and below the notation “ \lesssim ” means “ $\leq C$ ” where C is a constant and may change from line to line.

Because Herz spaces are generalizations of weighted Lebesgue spaces, it is natural to have the following question.

Question 1.1. Does “the boundedness of some sublinear operators on $L^p(\mathbb{R}^n)$ ” imply “the boundedness of the operators on Herz spaces $\dot{K}_p^{\alpha,q}(\mathbb{R}^n)$ and $K_p^{\alpha,q}(\mathbb{R}^n)$ ”?

In [30, 32], Li, Yang and their collaborators obtained a positive answer about the question with the help of size condition (I) :

$$\begin{cases} |Tf(x)| \lesssim |x|^{-n} \|f\|_{L^1(\mathbb{R}^n)}, & \text{if } \text{supp } f \subseteq A_k, |x| \geq 2^{k+1}, k \in \mathbb{Z}, \\ |Tf(x)| \lesssim 2^{-kn} \|f\|_{L^1(\mathbb{R}^n)}, & \text{if } \text{supp } f \subseteq A_k, |x| \leq 2^{k-2}, k \in \mathbb{Z}, \end{cases}$$

and size condition (II) :

$$|Tf(x)| \lesssim |x|^{-n} \|f\|_{L^1(\mathbb{R}^n)},$$

when $\text{supp } f \subseteq B_0$ and $|x| > 2$ or $\text{supp } f \subseteq A_k$ and $|x| \geq 2^{k+1}$ with $k \in \mathbb{N}$, and

$$|Tf(x)| \lesssim 2^{-kn} \|f\|_{L^1(\mathbb{R}^n)},$$

when $\text{supp } f \subseteq A_k$ and $|x| \leq 2^{k-2}$ with $k \geq 2$. The related theorems in [32] are the generalized versions of the results in [17, 36, 38] because the size conditions (I) and (II) are more general conditions than the condition (1).

They also generalized these results in [30, 32] to the cases of fractional integral operators T_σ ($0 < \sigma < n$) satisfying the size condition $(\text{I})_\sigma$:

$$\begin{cases} |T_\sigma(f)(x)| \lesssim |x|^{-(n-\sigma)} \|f\|_{L^1(\mathbb{R}^n)}, & \text{if } \text{supp } f \subseteq A_k, |x| \geq 2^{k+1}, k \in \mathbb{Z}, \\ |T_\sigma(f)(x)| \lesssim 2^{-k(n-\sigma)} \|f\|_{L^1(\mathbb{R}^n)}, & \text{if } \text{supp } f \subseteq A_k, |x| \leq 2^{k-2}, k \in \mathbb{Z}, \end{cases}$$

and size condition $(\text{II})_\sigma$: let T_σ be a sublinear operator satisfying

$$|T_\sigma(f)(x)| \lesssim |x|^{-(n-\sigma)} \|f\|_{L^1},$$

when $\text{supp } f \subseteq B_0$ and $|x| \geq 2$ or $\text{supp } f \subseteq A_k$ and $|x| \geq 2^{k+1}$ with $k \in \mathbb{N}$, and

$$|T_\sigma(f)(x)| \lesssim 2^{-k(n-\sigma)} \|f\|_{L^1},$$

when $\text{supp } f \subseteq A_k$ and $|x| \leq 2^{k-2}$ with $k \geq 2$. The size conditions $(\text{I})_\sigma$ and $(\text{II})_\sigma$ contain the classical fractional integral sublinear operators which satisfies

$$|T_\sigma(f)(x)| \lesssim \int_{\mathbb{R}^n} \frac{|f(y)|}{|x-y|^{n-\sigma}} dy, x \notin \text{supp } f$$

for any integrable function f with compact support.

Besides, there are a lot of very important research works on the theories of sublinear operators in Herz spaces, see [10, 15, 21, 22] and the references therein. After that, many researchers paid their attention on these topics and many function spaces related to Herz space appeared. We just list some but not all the references about these function spaces here, such as Herz spaces with variable exponents [1, 19, 20, 24, 28, 34, 35, 42], weighted Herz spaces and their variable exponents cases [26, 27, 29], Herz type function spaces and their variable exponents generalizations [4, 6, 14, 18, 31, 32, 33, 39, 41, 43, 44], and so on. For more details, please see the above references and references therein.

It is well known that $L^p(\mathbb{R}^n)$ is a particular case of Musielak-Orlicz spaces. In detail, denote by $L^0(\mathbb{R}^n)$ the set of all measurable functions $f : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$. A function $\varphi : \mathbb{R}^n \times [0, \infty) \rightarrow [0, \infty]$ is said to be a generalized φ -function if (1) $x \mapsto \varphi(x, |f(x)|)$ is measurable for every $f \in L^0(\mathbb{R}^n)$; (2) $t \mapsto \varphi(x, t)$ is a convex and left-continuous function with $\varphi(x, 0) = 0$, $\lim_{t \rightarrow 0^+} \varphi(x, t) = 0$ and $\lim_{t \rightarrow \infty} \varphi(x, t) = \infty$, for almost every $x \in \mathbb{R}^n$. Denote by $\Phi(\mathbb{R}^n)$ the set of all generalized φ -functions. Let $\varphi \in \Phi(\mathbb{R}^n)$ and $f \in L^0(\mathbb{R}^n)$, the semimodular of f is denoted by

$$\varrho_{L^\varphi}(f) := \int_{\mathbb{R}^n} \varphi(x, |f(x)|) dx.$$

Then the Musielak-Orlicz space is defined by

$$L^\varphi(\mathbb{R}^n) := \left\{ f \in L^0(\mathbb{R}^n) : \lim_{\lambda \rightarrow 0^+} \varrho_{L^\varphi}(\lambda f) = 0 \right\}.$$

The Musielak-Orlicz spaces are Banach spaces when they are equipped with the norm

$$\|f\|_{L^\varphi(\mathbb{R}^n)} := \inf \left\{ \lambda > 0 : \varrho_{L^\varphi} \left(\frac{f}{\lambda} \right) \leq 1 \right\}.$$

The Musielak-Orlicz space $L^\varphi(\mathbb{R}^n)$ is precisely the Lebesgue space $L^p(\mathbb{R}^n)$ if $\varphi(x, t) = t^p$, $t \in [0, \infty)$, $p \in [1, \infty)$. What is noteworthy is that the boundedness of Calderón-Zygmund singular integrals operators and some fractional integral sublinear operators on $L^\varphi(\mathbb{R}^n)$ can be found in [13]. Inspired by above works, in order to consider a similar question like **Question 1.1** in the new setting, we need the Musielak-Orlicz Herz spaces.

Definition 1.1. Let $\varphi \in \Phi(\mathbb{R}^n)$, $\alpha \in \mathbb{R}$, $p \in (0, \infty]$.

(1) The homogeneous Musielak-Orlicz Herz space $\dot{K}_\varphi^{\alpha, p}(\mathbb{R}^n)$ is defined by

$$\dot{K}_\varphi^{\alpha, p}(\mathbb{R}^n) = \left\{ f \in L_{\text{loc}}^\varphi(\mathbb{R}^n \setminus \{0\}) : \|f\|_{\dot{K}_\varphi^{\alpha, p}(\mathbb{R}^n)} < \infty \right\},$$

where

$$\|f\|_{\dot{K}_\varphi^{\alpha, p}(\mathbb{R}^n)} = \left\| \left\{ 2^{k\alpha} \|f\chi_k\|_{L^\varphi} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p}.$$

(2) The non-homogeneous Musielak-Orlicz Herz space $K_\varphi^{\alpha, p}(\mathbb{R}^n)$ is defined by

$$K_\varphi^{\alpha, p}(\mathbb{R}^n) = \left\{ f \in L_{\text{loc}}^\varphi(\mathbb{R}^n) : \|f\|_{K_\varphi^{\alpha, p}(\mathbb{R}^n)} < \infty \right\},$$

where

$$\|f\|_{K_\varphi^{\alpha, p}(\mathbb{R}^n)} = \left\| \left\{ 2^{k\alpha} \|f\tilde{\chi}_k\|_{L^\varphi} \right\}_{k \in \mathbb{N}} \right\|_{\ell^p}.$$

They are general frameworks of many generalizations of Herz spaces, such as Herz spaces with variable exponents, weighted Herz spaces and their variable exponents cases and so on. And we are interested in the following question.

Question 1.2. Does “the boundedness of some sublinear operators on $L^\varphi(\mathbb{R}^n)$ ” imply “the boundedness of the operators on the Musielak-Orlicz Herz spaces”?

In order to solve the question, one of the difficulties is to calculate

$$\frac{\|\chi_B\|_{L^\varphi(\mathbb{R}^n)}}{\|\chi_S\|_{L^\varphi(\mathbb{R}^n)}} \text{ and } \frac{\|\chi_S\|_{L^\varphi(\mathbb{R}^n)}}{\|\chi_B\|_{L^\varphi(\mathbb{R}^n)}},$$

where $B \subset \mathbb{R}^n$ is a ball and $S \subset B$ is a measurable subset. It is the main reason why we need some restrictions on the index φ (see Section 2 for details). Then we give positive answers to **Question 1.2** respectively by Theorems 1.2 and 1.3 for the sublinear operators satisfying the size conditions (I) and (II).

Theorem 1.2. Let $p \in (0, \infty]$ and $\varphi, \varphi^* \in \mathcal{A}_{\text{loc}} \cap \mathcal{A}_\infty$ be proper functions. Let T be a bounded sublinear operator in $L^\varphi(\mathbb{R}^n)$ satisfying the size condition (I). Then there exist $\delta_\varphi, \delta_{\varphi^*} \in (0, 1)$ such that T is bounded on $\dot{K}_\varphi^{\alpha, p}(\mathbb{R}^n)$ for all $\alpha \in (-n\delta_\varphi, n\delta_{\varphi^*})$.

Theorem 1.3. Let $p \in (0, \infty]$ and $\varphi, \varphi^* \in \mathcal{A}_{\text{loc}} \cap \mathcal{A}_\infty$ be proper functions. Let T be a bounded sublinear operator in $L^\varphi(\mathbb{R}^n)$ satisfying the size condition (II). Then there exist $\delta_\varphi, \delta_{\varphi^*} \in (0, 1)$ such that T is bounded on $K_\varphi^{\alpha, p}(\mathbb{R}^n)$ for all $\alpha \in (-n\delta_\varphi, n\delta_{\varphi^*})$.

Many classical operators in harmonic analysis satisfy the condition (1) which can imply size conditions (I) and (II), see [29, 32] and references therein. So the following corollary is due to Theorems 1.2 and 1.3.

Corollary 1.4. Let $p \in (0, \infty]$ and $\varphi, \varphi^* \in \mathcal{A}_{\text{loc}} \cap \mathcal{A}_\infty$ be proper functions. Let T be a bounded sublinear operator in $L^\varphi(\mathbb{R}^n)$ satisfying the condition (1). Then there exist $\delta_\varphi, \delta_{\varphi^*} \in (0, 1)$ such that T is bounded respectively on $\dot{K}_\varphi^{\alpha, p}(\mathbb{R}^n)$ and $K_\varphi^{\alpha, p}(\mathbb{R}^n)$ for all $\alpha \in (-n\delta_\varphi, n\delta_{\varphi^*})$.

Next two theorems indicate that **Question 1.2** are also correct respectively for the fractional integral operators under size conditions $(\text{I})_\sigma$ and $(\text{II})_\sigma$.

Theorem 1.5. Let $p \in (0, \infty]$ and $\varphi_2, \varphi_1^* \in \mathcal{A}_{\text{loc}} \cap \mathcal{A}_\infty$ be proper functions. Let $0 < \sigma < n$, φ_1 and φ_2 be σ -connected, T_σ be a bounded sublinear operator from $L^{\varphi_1}(\mathbb{R}^n)$ to $L^{\varphi_2}(\mathbb{R}^n)$ satisfying the size condition $(\text{I})_\sigma$. Then there exist $\delta_{\varphi_2}, \delta_{\varphi_1^*} \in (0, 1)$ such that T_σ is bounded from $\dot{K}_{\varphi_1}^{\alpha, p}(\mathbb{R}^n)$ to $\dot{K}_{\varphi_2}^{\alpha, p}(\mathbb{R}^n)$ for all $\alpha \in (-n\delta_{\varphi_2}, n\delta_{\varphi_1^*})$.

Theorem 1.6. Let $p \in (0, \infty]$ and $\varphi_2, \varphi_1^* \in \mathcal{A}_{\text{loc}} \cap \mathcal{A}_\infty$ be proper functions. Let $0 < \sigma < n$, φ_1 and φ_2 be σ -connected, T_σ be a bounded sublinear operator from $L^{\varphi_1}(\mathbb{R}^n)$ to $L^{\varphi_2}(\mathbb{R}^n)$ satisfying the size condition $(\text{II})_\sigma$. Then there exist $\delta_{\varphi_2}, \delta_{\varphi_1^*} \in (0, 1)$ such that T_σ is bounded from $K_{\varphi_1}^{\alpha, p}(\mathbb{R}^n)$ to $K_{\varphi_2}^{\alpha, p}(\mathbb{R}^n)$ for all $\alpha \in (-n\delta_{\varphi_2}, n\delta_{\varphi_1^*})$.

The structure of the paper is as follows. Section 2 contain some useful lemmas and the proof of Theorem 1.2. Section 3 is the proof of Theorem 1.5.

2. The proof of Theorems 1.2 and 1.3

The aim of this section is to prove the Theorems 1.2 and 1.3. As mentioned in Section 1, we need to calculate the following two fractions.

$$\frac{\|\chi_B\|_{L^\varphi(\mathbb{R}^n)}}{\|\chi_S\|_{L^\varphi(\mathbb{R}^n)}} \text{ and } \frac{\|\chi_S\|_{L^\varphi(\mathbb{R}^n)}}{\|\chi_B\|_{L^\varphi(\mathbb{R}^n)}},$$

where $B \subset \mathbb{R}^n$ is a ball and $S \subset B$ is a measurable subset. So, we need the following background knowledge.

Let \mathcal{Q} be a family of pairwise disjoint cubes in \mathbb{R}^n . Define the averaging operator $T_{\mathcal{Q}} : L_{\text{loc}}^1(\mathbb{R}^n) \rightarrow L^0(\mathbb{R}^n)$ by

$$T_{\mathcal{Q}}f := \sum_{Q \in \mathcal{Q}} \chi_Q M_Q f = \sum_{Q \in \mathcal{Q}} \chi_Q \cdot \frac{1}{|Q|} \int_Q |f(y)| dy.$$

The notation \mathcal{A} denotes the set of all $\varphi \in \Phi(\mathbb{R}^n)$ which have the property that the averaging operators T_Q are bounded uniformly in φ for all families \mathcal{Q} . In particular, denote \mathcal{A}_{loc} by the set of all $\varphi \in \Phi(\mathbb{R}^n)$ which have the property that the averaging operators T_Q are bounded uniformly in φ for all families \mathcal{Q} only contained a single cube, that is $\mathcal{Q} = \{Q\}$. So, $\mathcal{A} \subset \mathcal{A}_{\text{loc}}$. For more details, see page 117 in [5].

Let $\varphi \in \Phi(\mathbb{R}^n)$. The associated space of $L^\varphi(\mathbb{R}^n)$ is defined by

$$(L^\varphi(\mathbb{R}^n))' := \left\{ g \in L^0(\mathbb{R}^n) : \|g\|_{(L^\varphi(\mathbb{R}^n))'} < \infty \right\},$$

where

$$\|g\|_{(L^\varphi(\mathbb{R}^n))'} := \sup_{f \in L^\varphi : \|f\|_\varphi \leq 1} \int_{\mathbb{R}^n} |f(x)| |g(x)| dx.$$

A generalized φ -function is called a proper function if the set of simple functions $S(\mathbb{R}^n)$ satisfies $S(\mathbb{R}^n) \subset L^\varphi(\mathbb{R}^n) \cap (L^\varphi(\mathbb{R}^n))'$.

Let $\varphi \in \Phi(\mathbb{R}^n)$, for every $x \in \mathbb{R}^n$, denote by $\varphi^*(x, \cdot)$ the conjugate function of $\varphi(x, \cdot)$ which is defined by

$$\varphi^*(x, t) = \sup_{\mu \geq 0} (\mu t - \varphi(x, \mu))$$

for all $t \in [0, \infty)$. From Corollary 2.7.9 in [5], we know that a generalized φ -function is a proper function if and only if its conjugate function is a proper function. Besides, we have the following two lemmas.

Lemma 2.1 (Lemma 2.6.5 in [5]). Let $\varphi \in \Phi(\mathbb{R}^n)$, $f \in L^\varphi(\mathbb{R}^n)$ and $g \in L^{\varphi^*}(\mathbb{R}^n)$. Then

$$\int_{\mathbb{R}^n} |f| |g| dx \leq 2 \|f\|_{L^\varphi(\mathbb{R}^n)} \|g\|_{L^{\varphi^*}(\mathbb{R}^n)}.$$

Lemma 2.2 (Remark 4.5.8 in [5]). Let $\varphi \in \Phi(\mathbb{R}^n)$ be a proper function. The following are equivalent.

- (1) $\varphi \in \mathcal{A}_{\text{loc}}$.
- (2) $\varphi^* \in \mathcal{A}_{\text{loc}}$.
- (3) $\|\chi_Q\|_{L^\varphi(\mathbb{R}^n)} \|\chi_Q\|_{L^{\varphi^*}(\mathbb{R}^n)} \approx |Q|$, uniformly for all cubes or balls $Q \in \mathbb{R}^n$.

Let \mathcal{A}_∞ be the set of all $\varphi \in \Phi(\mathbb{R}^n)$ which have the following property: for every $\tau \in (0, 1)$ there exists $\beta \in (0, 1)$ such that if $D \subset \mathbb{R}^n$ (measurable) satisfies

$$|Q \cap D| \leq \tau |Q| \text{ for all } Q \in \mathcal{Q},$$

then

$$\left\| \sum_{Q \in \mathcal{Q}} t_Q \chi_{Q \cap D} \right\|_{L^\varphi(\mathbb{R}^n)} \leq \beta \left\| \sum_{Q \in \mathcal{Q}} t_Q \chi_Q \right\|_{L^\varphi(\mathbb{R}^n)}$$

for any sequence $\{t_Q\}_{Q \in \mathcal{Q}} \in \mathbb{R}^\mathcal{Q}$. Here $\mathbb{R}^\mathcal{Q}$ denotes the set of all real number sequences $\{t_Q\}_{Q \in \mathcal{Q}}$. The smallest constant β for $\tau = 1/2$ is called the \mathcal{A}_∞ -constant of φ . More detailed information can be found in [5] on pages 151, 159 and 160.

Lemma 2.3 (Lemma 5.4.12 in [5]). Let $\varphi \in \mathcal{A}_\infty$ and \mathcal{Q} be a family of pairwise disjoint cubes in \mathbb{R}^n . Then there exists $\delta \in (0, 1)$ and $K \geq 1$ which only depend on the \mathcal{A}_∞ -constant of φ such that

$$\left\| \sum_{Q \in \mathcal{Q}} t_Q \left| \frac{f}{M_Q f} \right|^\delta \chi_Q \right\|_{L^\varphi(\mathbb{R}^n)} \leq K \left\| \sum_{Q \in \mathcal{Q}} t_Q \chi_Q \right\|_{L^\varphi(\mathbb{R}^n)}$$

for all $\{t_Q\}_{Q \in \mathcal{Q}}$, $t_Q \geq 0$, and all $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ with $M_Q f \neq 0$, $Q \in \mathcal{Q}$.

The next main lemma solves the difficulty mentioned at the beginning of the section.

Lemma 2.4. Let $\varphi \in \mathcal{A}_{\text{loc}} \cap \mathcal{A}_\infty$ be a proper function. Then there exists $\delta \in (0, 1)$ such that for all balls B in \mathbb{R}^n and all measurable subsets $S \subset B$,

$$\frac{\|\chi_B\|_{L^\varphi(\mathbb{R}^n)}}{\|\chi_S\|_{L^\varphi(\mathbb{R}^n)}} \lesssim \frac{|B|}{|S|}, \quad \frac{\|\chi_S\|_{L^\varphi(\mathbb{R}^n)}}{\|\chi_B\|_{L^\varphi(\mathbb{R}^n)}} \lesssim \left(\frac{|S|}{|B|} \right)^\delta.$$

Proof. Let B be a ball and $S \subset B$ be a measurable subset arbitrarily, then

$$\|\chi_B\|_{L^\varphi} \frac{|S|}{|B|} = \|\chi_B\|_{L^\varphi} \frac{1}{|B|} \int_{\mathbb{R}^n} \chi_B(y) \chi_S(y) dy.$$

By Lemma 2.1 and 2.2,

$$\|\chi_B\|_{L^\varphi} \frac{|S|}{|B|} \leq \|\chi_B\|_{L^\varphi} \frac{2}{|B|} \|\chi_B\|_{L^{\varphi^*}} \|\chi_S\|_{L^\varphi} \lesssim \|\chi_S\|_{L^\varphi}.$$

Hence

$$\frac{\|\chi_B\|_{L^\varphi}}{\|\chi_S\|_{L^\varphi}} \lesssim \frac{|B|}{|S|}.$$

Next we prove the second inequality of the lemma. We take an open cube Q so that $B \subset Q \subset \sqrt{n}B$. Putting $f = \chi_S$ and $\mathcal{Q} = \{Q\}$ in Lemma 2.3 we have

$$\frac{\|\chi_S\|_{L^\varphi}}{\|\chi_Q\|_{L^\varphi}} \lesssim \left(\frac{|S|}{|Q|} \right)^\delta.$$

By virtue of $B \subset Q \subset \sqrt{n}B$,

$$\frac{\|\chi_S\|_{L^\varphi}}{\|\chi_B\|_{L^\varphi}} = \frac{\|\chi_S\|_{L^\varphi}}{\|\chi_B\|_{L^\varphi}} \cdot \frac{\|\chi_Q\|_{L^\varphi}}{\|\chi_B\|_{L^\varphi}} \leq \frac{\|\chi_S\|_{L^\varphi}}{\|\chi_Q\|_{L^\varphi}} \cdot \frac{\|\chi_{\sqrt{n}B}\|_{L^\varphi}}{\|\chi_B\|_{L^\varphi}}.$$

With the help of the first inequality in the lemma that we already proved, we have

$$\frac{\|\chi_S\|_{L^\varphi}}{\|\chi_B\|_{L^\varphi}} \lesssim \left(\frac{|S|}{|Q|} \right)^\delta \cdot \frac{|\sqrt{n}B|}{|B|} \lesssim \left(\frac{|S|}{|B|} \right)^\delta.$$

Thus we finish the proof of Lemma 2.4. \square

Now we turn to prove Theorem 1.2. We omit the proof of Theorem 1.3 because it is similar to the proof of Theorem 1.2.

Proof of Theorem 1.2. By the equation $f = \sum_{j \in \mathbb{Z}} f \chi_j$, we have

$$\begin{aligned} \|Tf\|_{\dot{K}_\varphi^{\alpha,p}} &= \left\| \left\{ 2^{k\alpha} \|T(f)\chi_k\|_{L^\varphi} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \left\| \left\{ 2^{k\alpha} \left\| \left(\sum_{j=-\infty}^{\infty} |T(f\chi_j)| \right) \chi_k \right\|_{L^\varphi} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \left\| \left\{ 2^{k\alpha} \left\| \left(\sum_{j=-\infty}^{k-2} |T(f\chi_j)| \right) \chi_k \right\|_{L^\varphi} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\quad + \left\| \left\{ 2^{k\alpha} \left\| \left(\sum_{j=k-1}^{k+1} |T(f\chi_j)| \right) \chi_k \right\|_{L^\varphi} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\quad + \left\| \left\{ 2^{k\alpha} \left\| \left(\sum_{j=k+2}^{\infty} |T(f\chi_j)| \right) \chi_k \right\|_{L^\varphi} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &=: (I_1 + I_2 + I_3). \end{aligned}$$

For I_2 , using the φ -boundedness of T , we obtain

$$\begin{aligned} I_2 &\lesssim \left\| \left\{ 2^{k\alpha} \sum_{j=k-1}^{k+1} \|T(f\chi_j)\chi_k\|_{L^\varphi} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \left\| \left\{ 2^{k\alpha} \sum_{j=k-1}^{k+1} \|T(f\chi_j)\|_{L^\varphi} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \left\| \left\{ 2^{k\alpha} \sum_{j=k-1}^{k+1} \|f\chi_j\|_{L^\varphi} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \|f\|_{\dot{K}_\varphi^{\alpha,p}}. \end{aligned}$$

For I_1 , by the fact that $j \leq k-2$, $x \in A_k$, and Lemma 2.1, we have

$$|T(f\chi_j)(x)| \lesssim 2^{-kn} \|f\chi_j\|_{L^1} \lesssim 2^{-kn} \|f\chi_j\|_{L^\varphi} \|\chi_{B_j}\|_{L^{\varphi^*}}.$$

Then

$$\begin{aligned} I_1 &\lesssim \left\| \left\{ 2^{k(\alpha-n)} \left\| \left(\sum_{j=-\infty}^{k-2} \|f\chi_j\|_{L^\varphi} \|\chi_{B_j}\|_{L^{\varphi^*}} \right) \chi_k \right\|_{L^\varphi} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \left\| \left\{ 2^{k(\alpha-n)} \|\chi_k\|_{L^\varphi} \left(\sum_{j=-\infty}^{k-2} \|f\chi_j\|_{L^\varphi} \|\chi_{B_j}\|_{L^{\varphi^*}} \right) \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \left\| \left\{ \sum_{j=-\infty}^{k-2} \left(2^{k\alpha} \|f\chi_j\|_{L^\varphi} 2^{-kn} \|\chi_{B_j}\|_{L^{\varphi^*}} \|\chi_k\|_{L^\varphi} \right) \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p}. \end{aligned}$$

Due to Lemma 2.2 and Lemma 2.4, there exists $\delta_{\varphi^*} \in (0, 1)$ such that

$$2^{-kn} \|\chi_{B_j}\|_{L^{\varphi^*}} \|\chi_k\|_{L^\varphi} \leq 2^{-kn} \|\chi_{B_j}\|_{L^{\varphi^*}} \|\chi_{B_k}\|_{L^\varphi} \lesssim \frac{\|\chi_{B_j}\|_{L^{\varphi^*}}}{\|\chi_{B_k}\|_{L^{\varphi^*}}} \lesssim 2^{n\delta_{\varphi^*}(j-k)}.$$

Then

$$I_1 \lesssim \left\| \left\{ \sum_{j=-\infty}^{k-2} 2^{j\alpha} \|f\chi_j\|_{L^\varphi} 2^{(n\delta_{\varphi^*}-\alpha)(j-k)} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p}.$$

If $\alpha \in (-\infty, n\delta_{\varphi^*})$,

$$I_1 \lesssim \left\| \left\{ 2^{j\alpha} \|f\chi_j\|_{L^\varphi} \right\}_{j \in \mathbb{Z}} \right\|_{\ell^p} = \|f\|_{\dot{K}_\varphi^{\alpha,p}}.$$

For I_3 , we use $j \geq k+2$ and Lemma 2.1, we have

$$|T(f\chi_j)(x)| \chi_k(x) \lesssim 2^{-jn} \|f\chi_j\|_{L^1} \lesssim 2^{-jn} \|f\chi_j\|_{L^\varphi} \|\chi_{B_j}\|_{L^{\varphi^*}} \chi_k(x).$$

It follows that

$$I_3 \lesssim \left\| \left\{ \sum_{j=k+2}^{\infty} \left(2^{k\alpha-jn} \|f\chi_j\|_{L^\varphi} \|\chi_{B_j}\|_{L^{\varphi^*}} \|\chi_k\|_{L^\varphi} \right) \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p}.$$

By Lemmas 2.2 and 2.4, there exists $\delta_\varphi \in (0, 1)$ such that

$$2^{-jn} \|\chi_{B_j}\|_{L^{\varphi^*}} \|\chi_k\|_{L^\varphi} \leq 2^{-jn} \|\chi_{B_j}\|_{L^{\varphi^*}} \|\chi_{B_k}\|_{L^\varphi} \lesssim \frac{\|\chi_{B_k}\|_{L^\varphi}}{\|\chi_{B_j}\|_{L^\varphi}} \lesssim 2^{n\delta_\varphi(k-j)}.$$

Then

$$I_3 \lesssim \left\| \left\{ \sum_{j=k+2}^{\infty} 2^{j\alpha} \|f\chi_j\|_{L^\varphi} 2^{(n\delta_\varphi+\alpha)(k-j)} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p}.$$

If $\alpha \in (-n\delta_\varphi, \infty)$,

$$I_3 \lesssim \left\| \left\{ 2^{j\alpha} \|f\chi_j\|_{L^\varphi} \right\}_{j \in \mathbb{Z}} \right\|_{\ell^p} = \|f\|_{\dot{K}_\varphi^{\alpha,p}}.$$

Combining the estimates on I_1, I_2 and I_3 , we obtain

$$\|Tf\|_{\dot{K}_\varphi^{\alpha,p}} \lesssim \|f\|_{\dot{K}_\varphi^{\alpha,p}}.$$

This finishes the proof of Theorem 1.2. \square

3. The proof of Theorems 1.5 and 1.6

In this section we aim to prove Theorems 1.5 and 1.6. For $0 < \sigma < n$, we say φ_1 and φ_2 are σ -connected if $\varphi_1, \varphi_2 \in \Phi(\mathbb{R}^n)$ and

$$|B|^{\sigma/n} \|\chi_B\|_{L^{\varphi_2}(\mathbb{R}^n)} \lesssim \|\chi_B\|_{L^{\varphi_1}(\mathbb{R}^n)}.$$

The condition is reasonable because it is nearly a necessary and sufficient condition for the boundedness of fractional integral operators between two different Musielak-Orlicz Hardy spaces, for more details please see [25].

Next, we prove Theorem 1.5 only, because the Theorem 1.6 is also similar to the proof of Theorem 1.5.

Proof of Theorem 1.5. First of all we have

$$\begin{aligned} \|T_\sigma(f)\|_{\dot{K}_{\varphi_2}^{\alpha,p}} &= \left\| \left\{ 2^{k\alpha} \|T_\sigma(f)\chi_k\|_{L^{\varphi_2}} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \left\| \left\{ 2^{k\alpha} \left\| \left(\sum_{j=-\infty}^{\infty} |T_\sigma(f\chi_j)| \right) \chi_k \right\|_{L^{\varphi_2}} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \left\| \left\{ 2^{k\alpha} \left\| \left(\sum_{j=-\infty}^{k-2} |T_\sigma(f\chi_j)| \right) \chi_k \right\|_{L^{\varphi_2}} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\quad + \left\| \left\{ 2^{k\alpha} \left\| \left(\sum_{j=k-1}^{k+1} |T_\sigma(f\chi_j)| \right) \chi_k \right\|_{L^{\varphi_2}} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\quad + \left\| \left\{ 2^{k\alpha} \left\| \left(\sum_{j=k+2}^{\infty} |T_\sigma(f\chi_j)| \right) \chi_k \right\|_{L^{\varphi_2}} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &=: II_1 + II_2 + II_3. \end{aligned}$$

For II_2 , using the fact that $T_\sigma(f)$ maps $L^{\varphi_1}(\mathbb{R}^n)$ into $L^{\varphi_2}(\mathbb{R}^n)$, we obtain

$$II_2 \lesssim \left\| \left\{ 2^{k\alpha} \sum_{j=k-1}^{k+1} \|f\chi_j\|_{L^{\varphi_1}} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \lesssim \|f\|_{\dot{K}_{\varphi_1}^{\alpha,p}}.$$

For II_1 , notice that $j \leq k-2$, by the condition $(I)_\sigma$, we have

$$|T_\sigma(f\chi_j)(x)| \chi_k(x) \lesssim 2^{-k(n-\sigma)} \|f\chi_j\|_{L^1} \chi_k(x).$$

By Lemma 2.1,

$$\begin{aligned} II_1 &\lesssim \left\| \left\{ 2^{k\alpha} \left\| \left(\sum_{j=-\infty}^{k-2} 2^{-k(n-\sigma)} \|f\chi_j\|_{L^{\varphi_1}} \|\chi_{B_j}\|_{L^{\varphi_1^*}} \right) \chi_k \right\|_{L^{\varphi_2}} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &= \left\| \left\{ \sum_{j=-\infty}^{k-2} \left[2^{k(\alpha-n+\sigma)} \|f\chi_j\|_{L^{\varphi_1}} \|\chi_k\|_{L^{\varphi_2}} \|\chi_{B_j}\|_{L^{\varphi_1^*}} \right] \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \left\| \left\{ \sum_{j=-\infty}^{k-2} \left(2^{k(\alpha-n)} \|f\chi_j\|_{L^{\varphi_1}} \|\chi_{B_k}\|_{L^{\varphi_1^*}} \|\chi_{B_k}\|_{L^{\varphi_1}} \frac{\|\chi_{B_j}\|_{L^{\varphi_1^*}}}{\|\chi_{B_k}\|_{L^{\varphi_1^*}}} \right) \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p}. \end{aligned}$$

Due to Lemma 2.4, we have

$$\begin{aligned} II_1 &\lesssim \left\| \left\{ \sum_{j=-\infty}^{k-2} 2^{j\alpha} \|f\chi_j\|_{L^{\varphi_1}} 2^{(n\delta_{\varphi_1^*}-\alpha)(j-k)} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \left\| \left\{ 2^{j\alpha} \|f\chi_j\|_{L^{\varphi_1}} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} = \|f\|_{\dot{K}_{\varphi_1}^{\alpha,p}}, \end{aligned}$$

where the last inequality is due to $\alpha < n\delta_{\varphi_1^*}$.

For II_3 , notice that $j \geq k+2$, by the condition $(I)_\sigma$ and Lemma 2.1, we have

$$|T_\sigma(f\chi_j)(x)| \chi_k(x) \lesssim 2^{-j(n-\sigma)} \|f\chi_j\|_{L^1} \lesssim 2^{-j(n-\sigma)} \|f\chi_j\|_{L^{\varphi_1}} \|\chi_{B_j}\|_{L^{\varphi_1^*}} \chi_k(x).$$

By Lemma 2.4 and $\alpha > -n\delta_{\varphi_2}$, we deduce that

$$\begin{aligned} II_3 &\lesssim \left\| \left\{ \sum_{j=k+2}^{\infty} \left[2^{k\alpha-j(n-\sigma)} \|f\chi_j\|_{L^{\varphi_1}} \|\chi_{B_j}\|_{L^{\varphi_1^*}} \|\chi_k\|_{L^{\varphi_2}} \right] \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \left\| \left\{ \sum_{j=k+2}^{\infty} \left(2^{k\alpha-jn} \|f\chi_j\|_{L^{\varphi_1}} \|\chi_{B_j}\|_{L^{\varphi_1^*}} \|\chi_{B_j}\|_{L^{\varphi_1}} \frac{\|\chi_{B_k}\|_{L^{\varphi_2}}}{\|\chi_{B_j}\|_{L^{\varphi_2}}} \right) \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \end{aligned}$$

$$\begin{aligned} &\lesssim \left\| \left\{ \sum_{j=k+2}^{\infty} 2^{j\alpha} \|f\chi_j\|_{L^{\varphi_1}} 2^{(n\delta_{\varphi_2} + \alpha)(k-j)} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} \\ &\lesssim \left\| \left\{ 2^{j\alpha} \|f\chi_j\|_{L^{\varphi_1}} \right\}_{k \in \mathbb{Z}} \right\|_{\ell^p} = \|f\|_{\dot{K}_{\varphi_1}^{\alpha,p}}. \end{aligned}$$

Combining II_1 , II_2 and II_3 , we obtain

$$\|Tf\|_{\dot{K}_{\varphi_2}^{\alpha,p}} \lesssim \|f\|_{\dot{K}_{\varphi_1}^{\alpha,p}}.$$

This finishes the proof of Theorem 1.5. \square

Acknowledgement

The authors would like to thank the anonymous referee for his suggestions.

References

- [1] ALMEIDA, A.; DRIHEM, D. Maximal, potential and singular type operators on Herz spaces with variable exponents. *J. Math. Anal. Appl.* **394** (2012), no. 2, 781–795. [MR2927498](#), [Zbl 1250.42077](#), doi: [10.1016/j.jmaa.2012.04.043](#). 1289
- [2] BAERNSTEIN, A.; SAWYER, E. Embedding and multiplier theorems for $H^p(\mathbb{R}^n)$. *Mem. Amer. Math. Soc.* **53** (1985), no. 318, iv+82 pp. [MR0776176](#), [Zbl 0566.42013](#), doi: [10.1090/memo/0318](#). 1287, 1288
- [3] BEURLING, A. Construction and analysis of some convolution algebras. *Ann. Inst. Fourier* **14** (1964), no. 2, 1–32. [MR0182839](#), [Zbl 0133.07501](#), doi: [10.5802/aif.172](#). 1287
- [4] DRIHEM, D.; HERAIZ, R. Boundedness of some bilinear operators on variable Herz-type Hardy spaces. *J. Pseudo-Differ. Oper. Appl.* **10** (2019), no. 3, 601–648. [MR3990262](#), [Zbl 1430.42014](#), doi: [10.1007/s11868-018-0258-8](#). 1289
- [5] DIENING, L.; HARJULEHTO, P.; HÄSTÖ, P.; RŮŽIČKA, M. Lebesgue and Sobolev Spaces with Variable Exponents. Lecture Notes in Mathematics, 2017. Springer, Heidelberg, 2011. x+509 pp. ISBN: 978-3-642-18362-1. [MR2790542](#), [Zbl 1222.46002](#), doi: [10.1007/978-3-642-18363-8](#). 1292, 1293
- [6] DONG, B.; XU, J. Herz-Morrey type Besov and Triebel-Lizorkin spaces with variable exponents. *Banach J. Math. Anal.* **9** (2015), no. 1, 75–101. [MR3296087](#), [Zbl 1333.46026](#), doi: [10.15352/bjma/09-1-7](#). 1289
- [7] FEICHTINGER, H. An elementary approach to Wiener's third Tauberian theorem on Euclidean n -space. in *Symposia Mathematica Vol. XXIX*, pp. 267–301. Academic Press, New York, 1987. ISBN: 0-12-612229-6. [MR0951188](#). 1287
- [8] FLETT, T. Some elementary inequalities for integrals with applications to Fourier transforms. *Proc. Lond. Math. Soc.* **29** (1974), no. 3, 538–556. [MR0360959](#), [Zbl 0318.46044](#), doi: [10.1112/plms/s3-29.3.538](#). 1287
- [9] FEICHTINGER, H.; WEISZ, F. Wiener amalgams and pointwise summability of Fourier transforms and Fourier series. *Math. Proc. Camb. Philos. Soc.* **140** (2006), no. 3, 509–536. [MR2225645](#), [Zbl 1117.43001](#), doi: [10.1017/S0305004106009273](#). 1288
- [10] FEICHTINGER, H.; WEISZ, F. Herz spaces and summability of Fourier transforms. *Math. Nachr.* **281** (2008), no. 3, 309–324. [MR2392115](#), [Zbl 1189.42001](#), doi: [10.1002/mana.200510604](#). 1288, 1289
- [11] GARCÍA-CUERVA, J. Hardy spaces and Beurling algebras. *J. Lond. Math. Soc.* **39** (1989), no. 3, 499–513. [MR1002462](#), [Zbl 0681.42014](#), doi: [10.1112/jlms/s2-39.3.499](#). 1287

- [12] GARCÍA-CUERVA, J.; HERRERO, M. A theory of Hardy spaces associated to the Herz spaces. *Proc. Lond. Math. Soc.* **69** (1994), no. 3, 605–628. [MR1289865](#), [Zbl 0831.42012](#), doi: [10.1112/plms/s3-69.3.605](#). [1287](#)
- [13] HARJULEHTO, P.; HÄSTÖ, P. Orlicz spaces and generalized Orlicz spaces. Lecture Notes in Mathematics, 2236. Springer, Cham, 2019. x+167 pp. ISBN:978-3-030-15099-0. [MR3931352](#), [Zbl 1436.46002](#), doi: [10.1007/978-3-030-15100-3](#). [1290](#)
- [14] HERNÁNDEZ, E.; WEISS, G.; YANG, D. The φ -transform and wavelet characterizations of Herz-type spaces. *Collect. Math.* **47** (1996), no. 3, 285–320. [MR1437657](#), [Zbl 0878.42011](#). [1289](#)
- [15] HERNÁNDEZ, E.; YANG, D. Interpolation of Herz spaces and applications. *Math. Nachr.* **205** (1999), no. 1, 69–87. [MR1709163](#), [Zbl 0936.41001](#), doi: [10.1002/mana.3212050104](#). [1289](#)
- [16] HERZ, C. Lipschitz spaces and Bernstein's theorem on absolutely convergent Fourier transforms. *J. Math. Mech.* **18** (1968), 283–323. [MR0438109](#), [Zbl 0177.15701](#), doi: [10.1512/iumj.1969.18.18024](#). [1287](#)
- [17] HOFMANN, S. Singular integrals with power weights. *Proc. Amer. Math. Soc.* **110** (1990), no. 2, 343–353. [MR1021899](#), [Zbl 0717.42018](#), doi: [10.2307/2048076](#). [1288](#), [1289](#)
- [18] HO, K. Sublinear operators on Herz-Hardy spaces with variable exponents. *Math. Nachr.* **295** (2022), no. 5, 876–889. [MR4453941](#), [Zbl 07747274](#), doi: [10.1002/mana.202000161](#). [1289](#)
- [19] HO, K. Extrapolation to Herz spaces with variable exponents and applications. *Rev. Mat. Complut.* **33** (2020), no. 2, 437–463. [MR4082594](#), [Zbl 1465.42015](#), doi: [10.1007/s13163-019-00320-3](#). [1289](#)
- [20] HO, K. Spherical maximal function, maximal Bochner-Riesz mean and geometrical maximal function on Herz spaces with variable exponents. *Rend. Circ. Math. Palermo II. Ser.* **70** (2021), no. 1, 559–574. [MR4234327](#), [Zbl 1462.42030](#), doi: [10.1007/s12215-020-00511-8](#). [1289](#)
- [21] HO, K. Young's inequalities and Hausdorff-Young inequalities on Herz spaces. *Boll. Unione Mat. Ital.* **11** (2018), no. 4, 469–481. [MR3869581](#), [Zbl 1403.42002](#), doi: [10.1007/s40574-017-0147-8](#). [1289](#)
- [22] HU, G. Boundedness of sublinear operators on the homogeneous Herz spaces. *Publ. Mat.* **47** (2003), no. 1, 143–159. [MR1970898](#), [Zbl 1043.42008](#), doi: [10.5565/PUBLMAT_47103_07](#). [1289](#)
- [23] HUANG, L.; WEISZ, F.; YANG, D.; YUAN, W. Summability of Fourier transforms on mixed-norm Lebesgue spaces via associated Herz spaces. *Anal. Appl. (Singap.)* **21** (2023), no. 2, 279–328. [MR4539218](#), [Zbl 1518.42015](#), doi: [10.1142/S0219530521500135](#). [1288](#)
- [24] HUMBERTO, R.; STEFAN, S. Riesz potential operator in continual variable exponents Herz spaces. *Math. Nachr.* **288** (2015), no. 4, 465–475. [MR3320460](#), [Zbl 1325.46034](#), doi: [10.1002/mana.201300270](#). [1289](#)
- [25] HUY, D.; KY, L. Boundedness of fractional integral operators on Musielak-Orlicz Hardy spaces. *Math. Nachr.* **294** (2021), no. 12, 2340–2354. [MR4371302](#), [Zbl 07747395](#), doi: [10.1002/mana.201900392](#). [1296](#)
- [26] IZUKI, M.; NOI, T. Boundedness of fractional integrals on weighted Herz spaces with variable exponent. *J. Inequal. Appl.* (2016), Paper No. 199, 15 pp. [MR3537090](#), [Zbl 1346.42014](#), doi: [10.1186/s13660-016-1142-9](#). [1289](#)
- [27] IZUKI, M.; NOI, T. An intrinsic square function on weighted Herz spaces with variable exponent. *J. Math. Inequal.* **11** (2017), no. 3, 799–816. [MR3732815](#), [Zbl 1375.42037](#), doi: [10.7153/jmi-2017-11-62](#). [1289](#)
- [28] IZUKI, M. Boundedness of sublinear operators on Herz spaces with variable exponent and application to wavelet characterization. *Anal. Math.* **36** (2010), no. 1, 33–50. [MR2606575](#), [Zbl 1224.42025](#), doi: [10.1007/s10476-010-0102-8](#). [1289](#)

- [29] LU, S.; YANG, D. The decomposition of weighted Herz spaces on \mathbb{R}^n and its application. *Sci. China, Ser. A* **38** (1995), no. 2, 147–158. [MR1338138](#), [Zbl 0830.42014](#). [1289, 1291](#)
- [30] LI, X.; YANG, D. Boundedness of some sublinear operators on Herz spaces. *Illinois J. Math.* **40** (1996), no. 3, 484–501. [MR1407632](#), [Zbl 0956.46025](#), doi: [10.1215/ijm/1255986021](#). [1288, 1289](#)
- [31] LI, Y.; YANG, D.; HUANG, L. Real-variable theory of Hardy spaces associated with generalized Herz spaces of Rafeiro and Samko. Lecture Notes in Mathematics, 2320. *Springer, Singapore*, 2022. xix+647 pp. ISBN: 978-981-19-6787-0. [MR4628102](#), [Zbl 07634795](#), doi: [10.1007/978-981-19-6788-7](#). [1289](#)
- [32] LU, S.; YANG, D.; HU, G. Herz type spaces and their applications. *Science Press, Beijing*, 2008. ISBN: 978-7-03-020909-2. [1288, 1289, 1291](#)
- [33] LU, Y.; ZHU, Y. Boundedness of multilinear Calderón-Zygmund singular operators on Morrey-Herz spaces with variable exponents. *Acta Math. Sin.* **30** (2014), no. 7, 1180–1194. [MR3226766](#), [Zbl 1304.42042](#), doi: [10.1007/s10114-014-3410-2](#). [1289](#)
- [34] RAFEIRO, H.; SAMKO, S. Riesz potential operator in continual variable exponents Herz spaces. *Math. Nachr.* **288** (2015), no. 4, 465–475. [MR3320460](#), [Zbl 1325.46034](#), doi: [10.1002/mana.201300270](#). [1289](#)
- [35] SAMKO, S. Variable exponent Herz spaces. *Mediterr. J. Math.* **10** (2013), no. 4, 2007–2025. [MR3119348](#), [Zbl 1285.46027](#), doi: [10.1007/s00009-013-0285-x](#). [1289](#)
- [36] STEIN, E. Note on singular integrals. *Proc. Amer. Math. Soc.* **8** (1957), 250–254. [MR0088606](#), [Zbl 0077.27301](#), doi: [10.2307/2033721](#). [1288, 1289](#)
- [37] STEIN, E.; WEISS, G. Introduction to Fourier analysis on Euclidean spaces. *Princeton University Press, Princeton, NJ*, 1971. x+297 pp. ISBN: 0-691-08078-X. [MR0304972](#), [Zbl 0232.42007](#). [1288](#)
- [38] SORIA, F.; WEISS, G. A remark on singular integrals and power weights. *Indiana Univ. Math. J.* **43** (1994), no. 1, 187–204. [MR1275458](#), [Zbl 0803.42004](#), doi: [10.1512/iumj.1994.43.43009](#). [1288, 1289](#)
- [39] SHI, C.; XU, J. Herz type Besov and Triebel-Lizorkin spaces with variable exponent. *Front. Math. China* **8** (2013), no. 4, 907–921. [MR3057227](#), [Zbl 1297.46029](#), doi: [10.1007/s11464-012-0248-8](#). [1289](#)
- [40] TORCHINSKY, A. Real variable methods in harmonic analysis. Pure and Applied Mathematics, 123. *Academic Press, Orlando, FL*, 1986. xii+462 pp. ISBN: 0-12-695460-7. [MR0869816](#), [Zbl 0621.42001](#). [1288](#)
- [41] WANG, H.; LIU, Z. The Herz-type Hardy spaces with variable exponent and their applications. *Taiwanese J. Math.* **16** (2012), no. 4, 1363–1389. [MR2951143](#), [Zbl 1260.42012](#), doi: [10.11650/twjm/1500406739](#). [1289](#)
- [42] WANG, L.; TAO, S. Parameterized Littlewood-Paley operators and their commutators on Herz spaces with variable exponents. *Turk. J. Math.* **40** (2016), no. 1, 122–145. [MR3438791](#), [Zbl 1424.42032](#), doi: [10.3906/mat-1412-52](#). [1289](#)
- [43] XU, J.; YANG, D. Herz-type Triebel-Lizorkin spaces. *Acta Math. Sin., Engl. Ser.* **21** (2005), no. 3, 643–654. [MR2145918](#), [Zbl 1092.42013](#), doi: [10.1007/s10114-004-0424-1](#). [1289](#)
- [44] YANG, D.; LIANG, Y.; KY, L. Real-variable theory of Musielak-Orlicz Hardy spaces. Lecture Notes in Mathematics, 2182. *Springer, Cham*, 2017. xiii+466 pp. ISBN: 978-3-319-54360-4. [MR3586020](#), [Zbl 1375.42038](#), doi: [10.1007/978-3-319-54361-1](#). [1289](#)

(Baohua Dong) CENTER FOR APPLIED MATHEMATICS OF JIANGSU PROVINCE, DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY OF INFORMATION SCIENCE AND TECHNOLOGY, NANJING 210044, CHINA

baochuadong@nuist.edu.cn

(Yu Li) DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY OF INFORMATION SCIENCE AND TECHNOLOGY, NANJING 210044, CHINA

wllly992021@163.com

(Jingshi Xu) CENTER FOR APPLIED MATHEMATICS OF GUANGXI, GUANGXI COLLEGES AND UNIVERSITIES KEY LABORATORY OF DATA ANALYSIS AND COMPUTATION, SCHOOL OF MATHEMATICS AND COMPUTING SCIENCE, GUILIN UNIVERSITY OF ELECTRONIC TECHNOLOGY, GUILIN 541004, CHINA

jingshixu@126.com

This paper is available via <http://nyjm.albany.edu/j/2023/29-49.html>.