PUBLICATIONS DE L'INSTITUT MATHÉMATIQUE
Nouvelle série tome 56 (70), 1994, 34–40

GRAPHICAL COMPOSITIONS AND WEAK CONGRUENCES

Miroslav Ploščica

Communicated by Žarko Mijajlović

Abstract. Graphical compositions of equivalences were introduced (independently) by B. Jónsson and H. Werner in order to determine whether a subset of $\mathrm{Eq}(X)$ (the set of all equivalences on the set X) is the set of all congruences of some algebra defined on X. Namely, a complete sublattice L of $\mathrm{Eq}(X)$ is the congruence lattice of some algebra defined on X if and only if L is closed under all graphical compositions. We generalize this result and prove that a similar characterization is possible for weak congruences (i. e. symmetric and transitive compatible relations).

Weak congruences were introduced and investigated by B. Šešelja, G. Vojvodić and A. Tepavčević in [**2**]–[**4**] and other papers. Let us recall basic concepts.

An algebra $\mathcal{A} = (A, F)$ is a set A (called the underlying set) endowed with some set F of finitary operations (called the basic operations of $\mathcal{A}$). A finitary function $f:\ A^n \longrightarrow A$ is called a polynomial of $\mathcal{A}$ if it can be obtained from projections, constant functions and basic operations of $\mathcal{A}$ by means of compositions.

Let X be any set. A weak equivalence on X is any symmetric and transitive binary relation. We denote by $\mathrm{Eq}(X)$, $\mathrm{E}_w(X)$ and $\mathrm{Rel}(X)$ the sets of all equivalences, weak equivalences and binary relations on the set X, respectively. Let $f:\ X^n \longrightarrow X$ be any function. We say that f preserves a relation $\rho \in \mathrm{Rel}(X)$ if $(x_1, y_1), \dots, (x_n, y_n) \in \rho$ implies $(f(x_1, \dots, x_n), f(y_1, \dots, y_n)) \in \rho$. A nullary function f (i.e. a costant $f \in X$) preserves $\rho \in \mathrm{Rel}(X)$ if $(f, f) \in \rho$. A binary relation $\rho \in \mathrm{Rel}(A)$ is called compatible with the algebra $\mathcal{A} = (A, F)$ if every $f \in F$ preserves ρ. Such a compatible relation is a (weak) congruence of $\mathcal{A}$ if it is a (weak) equivalence. It is easy to see that a relation ρ is a weak congruence of $\mathcal{A}$ if and only if it is a congruence of some subalgebra of $\mathcal{A}$. We denote by $\mathrm{Con}(\mathcal{A})$ and $\mathrm{C}_w(\mathcal{A})$ the sets of all congruences and weak congruences of $\mathcal{A}$, respectively.

The domain of a relation $\alpha \in \mathrm{Rel}(X)$ is the set $\mathrm{dom}(\alpha) = \{x \in X \mid (x, x) \in \alpha\}$. For any $\alpha \in \mathrm{Rel}(X)$ we consider its restrictions to its domain $\alpha \restriction \mathrm{dom}(\alpha) = \alpha \cap (\mathrm{dom}(\alpha))^2 = \{(x, y) \in \alpha \mid (x, x) \in \alpha,\ (y, y) \in \alpha\}$. It is easy to see that the

---

AMS Subject Classification (1991): Primary 08 A 30, 08 A 40 symmetric and transitive closure of $\alpha \restriction \mathrm{dom}(\alpha)$ is always a weak equivalence and we call it the weak equivalence generated by α. We stress that we form the closure of $\alpha \restriction \mathrm{dom}(\alpha)$ and not of α itself. In fact, the symmetric and transitive closure of any relation is a weak equivalence.

LEMMA 1. *Let α be a weak equivalence on an algebra $\mathcal{A} = (A, F)$. Then $\alpha \in \mathrm{C}_w(\mathcal{A})$ if and only if the following conditions hold:*

(i) $\mathrm{dom}(\alpha)$ *is a subalgebra of $\mathcal{A}$;*

(ii) *every unary polynomial f of the algebra* $\mathrm{dom}(\alpha)$ *preserves α.*

Proof. It is easy to see that any weak congruence satisfies (i) and (ii). Conversely, suppose that (i) and (ii) hold for $\alpha \in \mathrm{Eq}(A)$. Let $f : A^n \longrightarrow A$ be any of the basic operations of the algebra $\mathcal{A}$. Suppose that $(a_i, b_i) \in \alpha$ for $i = 1, \ldots, n$. Then clearly $a_i \in \mathrm{dom}(\alpha)$, $b_i \in \mathrm{dom}(\alpha)$ for every i. Let us consider the unary polynomial $f_1(x) = f(x, a_2, \ldots, a_n)$. Because of (ii), $(a_1, b_1) \in \alpha$ implies that $(f_1(a_1), f_1(b_1)) \in \alpha$ and therefore $(f(a_1, \ldots, a_n), f(b_1, a_2, \ldots, a_n)) \in \alpha$. Similarly, $(f(b_1, \ldots, b_i, a_{i+1}, \ldots, a_n), f(b_1, \ldots, b_{i+1}, a_{i+2}, \ldots, a_n)) \in \alpha$ holds for any $i = 0, 1, \ldots, n-1$. From the transitivity of α we infer that
$(f(a_1, \ldots, a_n), f(b_1, \ldots, b_n)) \in \alpha$. $\square$

LEMMA 2. *Let α be a compatible binary relation on an algebra $\mathcal{A} = (A, F)$. Then the weak equivalence generated by α is also compatible. (And hence it is a weak congruence.)*

Proof. Let $\beta \in \mathrm{E}_w(A)$ be generated by α. We prove that β satisfies (i), (ii) from Lemma 1.

It is easy to see that $\mathrm{dom}(\beta) = \mathrm{dom}(\alpha)$. Let $f : A^n \longrightarrow A$ be any of the basic operations of $\mathcal{A}$, let $\{a_1, \ldots, a_n\} \subseteq \mathrm{dom}(\beta)$. Then $(a_i, a_i) \in \alpha$ for every i and since α is compatible we obtain that $(f(a_1, \ldots, a_n), f(a_1, \ldots, a_n)) \in \alpha$, hence $f(a_1, \ldots, a_n) \in \mathrm{dom}(\alpha) = \mathrm{dom}(\beta)$.

To prove (ii), let f be a unary polynomial of the algebra $\mathrm{dom}(\beta)$. (That is, the constants used in f belong to $\mathrm{dom}(\beta)$.) Let $(x, y) \in \beta$. Then we have a finite sequence $x = z_0, z_1, \ldots, z_k = y$ such that, for every $i = 1, \ldots, k$, $(z_{i-1}, z_i) \in \alpha \restriction \mathrm{dom}(\alpha)$ or $(z_i, z_{i-1}) \in \alpha \restriction \mathrm{dom}(\alpha)$. Since $\mathrm{dom}(\alpha)$ is closed under f (the first part of this proof) and the relation α is compatible, it follows that $(f(z_{i-1}), f(z_i)) \in \alpha \restriction \mathrm{dom}(\alpha)$ or $(f(z_i), f(z_{i-1})) \in \alpha \restriction \mathrm{dom}(\alpha)$. Now $f(x) = f(z_0), f(z_1), \ldots, f(z_k) = f(y)$ is the sequence showing that $(f(x), f(y)) \in \beta$. $\square$

Now we recall the definition of graphical compositions. A (undirected) graph is a pair (V, E) of sets V and E, whose elements are called vertices and edges, together with a map $\nu : E \longrightarrow \mathcal{P}_1(V) \cup \mathcal{P}_2(V)$, where $\mathcal{P}_1(V)$ and $\mathcal{P}_2(V)$ are the sets of all one element subsets and of all two element subsets of V, respectively. If $\nu(e) = \{x, y\}$, we say that e is an edge between x and y. If $\nu(e) = \{x\}$, we say that e is a loop on x. Hence, we admit several edges with the same endpoints. (In **[5]** Werner excludes loops, but in the case of weak congruences they are useful.)

Let $G = (V, E)$ be a graph and let $\varphi : E \longrightarrow \mathrm{Rel}(X)$ be a mapping. A function $f : V \longrightarrow X$ is called a φ-compatible labelling if, for every $e \in E$,

$e = \{x, y\}$ implies that $(f(x), f(y)) \in \varphi(e)$. (The case $x = y$ is included.) For every two distinguished vertices $0, 1 \in V$ we define a relation

$$S_{G,0,1}(\varphi) = \{(a, b) \in X^2 \mid a = f(0), b = f(1) \text{ for some } \varphi\text{-compatible labelling } f\}.$$

Thus, $S_{G,0,1}$ is a mapping $\mathrm{Rel}(X)^E \longrightarrow \mathrm{Rel}(X)$. We define a mapping

$$P_{G,0,1} : \mathrm{E}_w(X)^E \longrightarrow \mathrm{E}_w(X)$$

by the rule that $P_{G,0,1}(\varphi)$ is the weak equivalence generated by $S_{G,0,1}(\varphi)$. Hence, every graph with two distinguished vertices determines a $|E|$-ary operation on the set $\mathrm{E}_w(X)$. Any such operation is called a graphical composition.

As an illustration, let us present two simple examples. (More examples can be found in [**5**].)

First, let G and φ be as follows.



(In pictures like this, each edge e is labelled by $\varphi(e)$.) It is easy to see that $S_{G,0,1}(\varphi) = \alpha \cap \beta$. If α and β are weak equivalences, then $S_{G,0,1}(\varphi)$ is also a weak equivalence and therefore $P_{G,0,1}(\varphi) = S_{G,0,1}(\varphi)$. Hence, this graphical composition is the usual intersection of two relations. By adding more edges between 0 and 1 we obtain a graphical composition that describes the intersection of arbitrarily many (even of infinitely many) relations.

As the second example we consider the following graph.



Suppose that $\alpha, \beta \in \mathrm{E}_w(X)$ and denote $Y = \mathrm{dom}(\alpha) \cap \mathrm{dom}(\beta)$. The restrictions $\alpha \restriction Y$ and $\beta \restriction Y$ are equivalences on the set Y. We claim that $P_{G,0,1}(\varphi)$ is the least equivalence on Y containing $\alpha \restriction Y$ and $\beta \restriction Y$ (the join in the lattice $\mathrm{Eq}(Y)$). First, it is easy to see that $S_{G,0,1}(\varphi)$ is equal to the relational product

$$\alpha \cdot \beta = \{(x, y) \in X^2 \mid (x, z) \in \alpha,\ (z, y) \in \beta \text{ for some } z \in X\}.$$

Since α and β are weak equivalences, it follows that $\mathrm{dom}(P_{G,0,1}(\varphi)) = \mathrm{dom}(S_{G,0,1}(\varphi)) = Y$, hence $P_{G,0,1}(\varphi) \in \mathrm{Eq}(Y)$. Further, $\alpha \restriction Y \subseteq P_{G,0,1}(\varphi)$. Indeed, if $(x, y) \in \alpha \restriction Y$, then $(y, y) \in \beta$, which shows that $(x, y) \in \alpha \cdot \beta \subseteq P_{G,0,1}(\varphi)$. For similar reasons, $\beta \restriction Y \subseteq P_{G,0,1}(\varphi)$. On the other hand, if θ is any weak equivalence containing both $\alpha \restriction Y$ and $\beta \restriction Y$, then also $S_{G,0,1}(\varphi) \restriction Y = \alpha \restriction Y \cdot \beta \restriction$

$Y \subseteq \theta$. Since $P_{G,0,1}(\varphi)$ is, by the definition, the least weak equivalence containing $S_{G,0,1}(\varphi) \restriction Y$, it follows that $P_{G,0,1}(\varphi) \subseteq \theta$.

If $\alpha, \beta \in \mathrm{Eq}(X)$, then $Y = X$ and $P_{G,0,1}(\varphi)$ is the usual join of equivalence relations. Hence, this graphical composition can be regarded as a generalization of the join operation for weak equivalences.

LEMMA 3. *Let $\mathcal{A} = (A, F)$ be an algebra, let $G = (V, E)$ be a graph, $0, 1 \in V$. Suppose that $\varphi : E \longrightarrow \mathrm{Rel}(A)$ is such that $\varphi(e)$ is a compatible relation on $\mathcal{A}$ for every $e \in E$. Then $S_{G,0,1}(\varphi)$ is also a compatible relation on $\mathcal{A}$.*

Proof. Let $(a_i, b_i) \in S_{G,0,1}(\varphi)$ for $i = 1, \ldots, k$. Let $g : A^k \longrightarrow A$ be any of the basic operations of $\mathcal{A}$. For every i we have a φ-compatible labelling $f_i : V \longrightarrow A$ with $f_i(0) = a_i$, $f_i(1) = b_i$. Define a function $f : V \longrightarrow A$ by $f(x) = g(f_1(x), \ldots, f_k(x))$. Then $f(0) = g(a_1, \ldots, a_k)$, $f(1) = g(b_1, \ldots, b_k)$. It remains to show that f is a φ-compatible labelling.

Let $e \in E$, $e = \{x, y\}$. Then $(f_i(x), f_i(y)) \in \varphi(e)$ for every $i = 1, \ldots, k$. Since, by the assumption, the relation $\varphi(e)$ is compatible, we obtain that $(f(x), f(y)) = (g(f_1(x), \ldots, f_k(x)), g(f_1(y), \ldots, f_k(y))) \in \varphi(e)$. $\square$

As a consequence of Lemma 2 and Lemma 3 we obtain the following assertion.

LEMMA 4. *Let $\mathcal{A} = (A, F)$ be an algebra. Then $\mathrm{C}_w(\mathcal{A})$ is a subset of $\mathrm{E}_w(A)$ closed under all graphical compositions (i.e. for any graph G with two distinguished vertices, if $\varphi : E \longrightarrow \mathrm{C}_w(\mathcal{A})$ then also $P_{G,0,1}(\varphi) \in \mathrm{C}_w(\mathcal{A})$).* $\square$

Now we show that closedness under all graphical compositions is not sufficient for characterization of those subsets of $\mathrm{E}_w(X)$ that are equal to $\mathrm{C}_w(\mathcal{A})$ for some algebra $\mathcal{A}$ defined on the set X. (Similarly as in the case of usual congruence relations.)

To see this, let $\mathcal{A}$ be an algebra whose congruence lattice looks as follows.

$$
\begin{array}{c}
\nabla \\
| \\
\psi \\
\vdots \\
| \\
\theta_2 \\
| \\
\theta_1 \\
| \\
\Delta
\end{array}
$$

Such an algebra $\mathcal{A}$ certainly exists, since the lattice is algebraic. We can assume that all elements of $\mathcal{A}$ are nullary operations (constants), so that $\mathrm{C}_w(\mathcal{A}) =$

$\mathrm{Con}(\mathcal{A})$ and our example serves both the case of usual and weak congruences. Let us consider the family $\mathcal{F} = \{\Delta, \nabla, \theta_1, \theta_2, \dots\} = \mathrm{Con}(\mathcal{A}) \setminus \{\psi\}$. This family cannot be the set of all (weak) congruences of any algebra, since it is not a complete sublattice of $\mathrm{Eq}(A)$. (It is not closed under infinite joins.) However, we claim that $\mathcal{F}$ is closed under all graphical compositions.

To see this, let $G = (V, E)$ be a graph, $0, 1 \in V$, $\varphi : E \longrightarrow \mathcal{F}$. Without loss of generality we can assume that $\nabla \notin \varphi(E)$. Indeed, if $\nabla \in \varphi(E)$, then we consider the graph $G' = (V, E')$, where $E' = \{e \in E \mid \varphi(e) \neq \nabla\}$, and the restriction $\varphi' = \varphi \restriction E'$. It is easy to see that $S_{G,0,1}(\varphi) = S_{G',0,1}(\varphi')$.

Thus, suppose that $\nabla \notin \varphi(E)$. We distinguish two cases. First, suppose that there is a path $(e_1, \dots, e_k)$ in E connecting 0 and 1. Then $P_{G,0,1}(\varphi)$ cannot be greater (in the sense of set inclusion) then the greatest relation among $\varphi(e_1), \dots, \varphi(e_k)$. Hence, $P_{G,0,1}(\varphi)$ is equal to Δ or to some θ_i.

The second possibility is that there is no path between 0 and 1. Then it is not difficult to see that $P_{G,0,1}(\varphi) = S_{G,0,1}(\varphi) = A^2 = \nabla$.

We have proved that $P_{G,0,1}(\varphi)$ cannot be equal to ψ, which means that $\mathcal{F}$ must be closed under all graphical compositions.

The example above suggests what we should add to graphical compositions. A family $\mathcal{F} \subseteq \mathrm{E}_w(X)$ is called up-directed if for every $\alpha, \beta \in \mathcal{F}$ there is a $\gamma \in \mathcal{F}$ with $\alpha \cup \beta \subseteq \gamma$. It is easy to see that if $\mathcal{F}$ is such an up-directed family, then the set-theoretical union $\bigcup \mathcal{F}$ is a weak equivalence. Further, if all relations in $\mathcal{F}$ are compatible with some algebraic structure on X, then $\bigcup \mathcal{F}$ is also compatible (and hence a weak congruence). We obtain the following assertion.

LEMMA 5. *For any algebra $\mathcal{A}$, the set $\mathrm{C}_w(\mathcal{A})$ is closed under unions of up-directed families $\mathcal{F} \subseteq \mathrm{C}_w(\mathcal{A})$.* $\square$

Now we are going to prove the converse of Lemmas 4 and 5. Let us suppose that $\mathcal{F} \subseteq \mathrm{E}_w(X)$ is closed under all graphical compositions and up-directed unions.

First notice that $\mathcal{F}$ is closed under intersections. (See the example preceding Lemma 2. In accordance with this example, the intersection of the empty family of relations is equal to the greatest relation X^2.) Hence, for every $\alpha \in \mathrm{Rel}(X)$ there is a smallest $\beta \in \mathcal{F}$ with $\alpha \subseteq \beta$. We use the notation $\beta = \alpha^{\mathcal{F}}$.

We shall use some special graphs. Let G be the graph, whose set of vertices is X and the number of edges between vertices x and y is equal to the number of all $\alpha \in \mathcal{F}$ containing (x, y). (This applies also to loops.) Formally, the set E of edges can be expressed as $E = \{(\{x, y\}, \alpha) \mid x, y \in X,\ \alpha \in \mathcal{F} \text{ and } (x, y) \in \alpha\}$ and the map ν is defined by $\nu((\{x, y\}, \alpha)) = \{x, y\}$.

Similarly we define the graph G^n (the n-th power of G). The set of vertices of G^n will be X^n and we put an edge $(\{\overline{x}, \overline{y}\}, \alpha)$ between $\overline{x} = (x_1, \dots, x_n)$ and $\overline{y} = (y_1, \dots, y_n)$ whenever $(x_i, y_i) \in \alpha$ for every $i = 1, \dots, n$. Hence, $G = G^1$.

The importance of the graphs defined above lies in the following easy fact.

LEMMA 6. *Let $\varphi : E \longrightarrow \mathcal{F}$ be defined by $\varphi((\{\overline{x}, \overline{y}\}, \alpha)) = \alpha$. A function*

$f : X^n \longrightarrow X$ is a φ-compatible labelling on G^n if and only if f preserves all $\alpha \in \mathcal{F}$. $\square$

Now we are ready to define an algebra on the set X whose set of all equivalences equals $\mathcal{F}$.

LEMMA 7. *Let $\mathcal{G}$ be the set of all finitary operations on X that preserve every $\alpha \in \mathcal{F}$. Let $\mathcal{A}$ be the algebra with X as the underlying set and $\mathcal{G}$ as the set of basic operations. Then $\mathcal{F} = \mathrm{C}_w(\mathcal{A})$.*

Proof. By the definition, every basic operation of $\mathcal{A}$ preserves all relations in $\mathcal{F}$, hence $\mathcal{F} \subseteq \mathrm{C}_w(\mathcal{A})$.

To prove the other inclusion, let $\alpha \in \mathrm{C}_w(\mathcal{A})$, i.e. α is a weak equivalence on X that is preserved by all $f \in \mathcal{G}$.

Let $\beta = \{(x_1, y_1), \dots, (x_k, y_k)\}$ be an arbitrary finite subset (subrelation) of α. Consider the graph G^n with $n = 3k$. We distinguish the vertices
$0 = (x_1, \dots, x_k, y_1, \dots, y_k, x_1, \dots, x_k)$,
$1 = (x_1, \dots, x_k, y_1, \dots, y_k, y_1, \dots, y_k)$.
Let φ be defined by $\varphi((\{\overline{x}, \overline{y}\}, \alpha)) = \alpha$. By our assumption, the relation $P_{G^n,0,1}(\varphi)$ belongs to $\mathcal{F}$. It is easy to see that for every $i = 1, \dots, n$ the i-th projection $f_i : X^n \longrightarrow X$ (i.e. $f_i(z_1, \dots, z_n) = z_i$) is a φ-compatible labelling. For $i = 1, \dots, k$ we obtain that $(f_i(0), f_i(1)) = (x_i, x_i) \in S_{G^n,0,1}(\varphi)$, $(f_{i+k}(0), f_{i+k}(1)) = (y_i, y_i) \in S_{G^n,0,1}(\varphi)$, $(f_{i+2k}(0), f_{i+2k}(1)) = (x_i, y_i) \in S_{G^n,0,1}(\varphi)$. Consequently, $\beta \subseteq S_{G^n,0,1}(\varphi) \restriction \mathrm{dom}(S_{G^n,0,1}(\varphi))$, hence $\beta \subseteq P_{G^n,0,1}(\varphi)$.

Further, for every $(x, y) \in S_{G^n,0,1}(\varphi)$ there is a φ-compatible labelling $f : X^n \longrightarrow X$ with $f(0) = x$, $f(1) = y$. By Lemma 6, $f \in \mathcal{G}$ and by our assumption, f preserves α. Since for every $i = 1, \dots, k$ we have $(x_i, x_i) \in \alpha$, $(y_i, y_i) \in \alpha$, $(x_i, y_i) \in \alpha$, it follows that $(x, y) = (f(0), f(1)) \in \alpha$. We have shown that $S_{G^n,0,1}(\varphi) \subseteq \alpha$. Since α is a weak equivalence, we obtain that $P_{G^n,0,1}(\varphi) \subseteq \alpha$.

Hence, for every such β there is $\gamma \in \mathcal{F}$ with $\beta \subseteq \gamma \subseteq \alpha$ (namely, $\gamma = P_{G^n,0,1}(\varphi)$, where the number n and the vertices 0, 1 depend on β). Then clearly $\beta^{\mathcal{F}} \subseteq \alpha$. The family

$$\{\beta^{\mathcal{F}} \mid \beta \text{ is a finite subset of } \alpha\}$$

is an up-directed subset of $\mathcal{F}$ and its union is α. Since $\mathcal{F}$ is closed under up-directed unions, we obtain that $\alpha \in \mathcal{F}$, which was to prove. $\square$

From Lemmas 4, 5 and 7 we obtain our main result.

THEOREM. *A family $\mathcal{F} \subseteq \mathrm{E}_w(X)$ is the set of all weak congruences of some algebra if and only if $\mathcal{F}$ is closed under all graphical compositions and up-directed unions.* $\square$

REFERENCES

[1] B. Jónsson, *Topics in Universal Algebra*, Lecture Notes in Math. **250**, Springer-Verlag, 1972.

[2] B. Šešelja, A. Tepavčević, *Infinitely distributive elements in the lattice of weak congruences, in: General Algebra 6*, Elsevier (North-Holland), 1990, 241–253.

[3] G. Vojvodić, B. Šešelja, *On the lattice of weak congruence relations*, Algebra Universalis **25** (1988) 121–130.

[4] G. Vojvodić, B. Šešelja, *The diagonal relation in the lattice of weak congruences and the representation of lattices*, Review of Research Fac. Sci. Univ. Novi Sad **19** (1989), 167–178.

[5] H. Werner, *Which partition lattices are congruence lattices?*, Coll. Math. Soc. J. Bólyai **14** (Lattice Theory), North-Holland, 1976, pp. 433–453.

Mathematical Institute
Slovak Academy of Sciences
Grešákova 6
04001 Košice
Slovakia

(Received 03 08 1994)
(Revised 07 02 1995)