PUBLICATIONS DE L'INSTITUT MATHÉMATIQUE
Nouvelle série, tome 92(106) (2012), 145–156 DOI: 10.2298/PIM1206145A

ON APPROXIMATION OF THE RIEMANN–STIELTJES INTEGRAL AND APPLICATIONS

Mohammad Wajeeh Alomari

Communicated by Gradimir Milovanović

Abstract. Several inequalities of Grüss type for the Stieltjes integral with various type of integrand and integrator are introduced. Some improvements inequalities are proved. Applications to the approximation problem of the Riemann–Stieltjes integral are also pointed out.

1. Introduction

In 2002, Guessab and Schmeisser [3], incorporate the mid-point and the trapezoid inequality together, and they proved the following companion of Ostrowski's inequality:

Theorem 1. *Assume that the function $f:[a,b]\to\mathbb{R}$ is of r-H-Hölder type, where $r\in(0,1]$ and $H>0$ are given, i.e., $|f(t)-f(s)|\leqslant H|t-s|^r$, for any $t,s\in[a,b]$. Then, for each $x\in[a,(a+b)/2]$, one has the inequality*

$$\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right| \leqslant H\left[\frac{2^{r+1}(x-a)^{r+1}+(a+b-2x)^{r+1}}{2^r(r+1)(b-a)}\right]. \tag{1.1}$$

This inequality is sharp for each admissible x. Equality is obtained if and only if $f=\pm Hf_*+c$, with $c\in\mathbb{R}$ and

$$f_*(t)=\begin{cases}(x-t)^r, & a\leqslant t\leqslant x\\ t(t-x)^r, & x\leqslant t\leqslant (a+b)/2\\ f_*(a+b-x), & (a+b)/2\leqslant t\leqslant b\end{cases}$$

In [**11**] Dragomir proved the following companion of the Ostrowski inequality for mappings of bounded variation.

2010 *Mathematics Subject Classification:* Primary 26D15, 26D20; Secondary 41A55.
Key words and phrases: Ostrowski's inequality, bounded variation.

THEOREM 2. *Let $f : [a,b] \to \mathbb{R}$ be a mapping of bounded variation on $[a,b]$. Then we have the inequalities:*

$$\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right| \leqslant \left[\frac{1}{4}+\left|\frac{x-(3a+b)/4}{b-a}\right|\right]\bigvee_a^b(f), \tag{1.2}$$

for any $x \in [a,(a+b)/2]$, where $\bigvee_a^b(f)$ denotes the total variation of f on $[a,b]$. The constant $1/4$ is best possible.

Also, Dragomir in [**12**] proved some companions of Ostrowski's integral inequality for absolutely continuous mappings. Among others, our interest is incorporated in the following result:

THEOREM 3. *Let $f : I \subset \mathbb{R} \to \mathbb{R}$ be an absolutely continuous function on $[a,b]$ such that $f' \in L_\infty[a,b]$. Then for all $x \in [a,(a+b)/2]$ we have the inequality*

$$\begin{aligned}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right| \\ \leqslant \left[\frac{1}{8}+2\left(\frac{x-(3a+b)/4}{b-a}\right)^2\right](b-a)\left\|f'\right\|_\infty.\end{aligned} \tag{1.3}$$

By Guessab–Schmeisser functional we mean the functional

$$\mathcal{GS}(f;u) := \int_a^{(a+b)/2}\frac{f(x)+f(a+b-x)}{2}du(x)-\frac{u((a+b)/2)-u(a)}{b-a}\int_a^b f(t)\,dt,$$

provided that the Stieltjes integral $\int_a^b \frac{1}{2}\big(f(x)+f(a+b-x)\big)\,du(x)$, and the Riemann integral $\int_a^b f(t)\,dt$ exist.

Motivated by Guessab–Schmeisser companion of Ostrowski's inequality (1.1), the author of this paper, has established the functional $\mathcal{GS}(f;u)$ in [**1**], and he has proved the following results in estimating $\mathcal{GS}(f;u)$.

THEOREM 4. *Let $f : [a,b] \to \mathbb{R}$ be an r-H-Hölder type mapping on $[a,b]$, where r and $H > 0$ are given, and $u : [a,b] \to \mathbb{R}$ be a mapping of bounded variation on $[a,b]$. Then the following inequality holds*

$$|\mathcal{GS}(f;u)| \leqslant \frac{H}{r+1}(b-a)^r\bigvee_a^{(a+b)/2}(u). \tag{1.4}$$

THEOREM 5. *Let $f : [a,b] \to \mathbb{R}$ be an r-H-Hölder type mapping on $[a,b]$, and $u : [a,b] \to \mathbb{R}$ be an L-Lipschitzian mapping on $[a,b]$, where r and $H, L > 0$ are given. Then the following inequality holds*

$$|\mathcal{GS}(f;u)| \leqslant \frac{LH}{(r+1)(r+2)}(b-a)^{r+1} \tag{1.5}$$

In this paper we point out several bounds for the functional $\mathcal{GS}(f;u)$ with various type of integrand and integrator. Improvements bounds for $\mathcal{GS}(f;u)$ are proved.

Finally, we apply the obtained results to approximate the Riemann–Stieltjes integral

$$\int_a^{(a+b)/2} \frac{f(x)+f(a+b-x)}{2}\,du(x)$$

in terms of the Riemann integral $\int_a^b f(t)\,dt$

2. The case of bounded variation integrators

2.1. The case of bounded variation integrands.

THEOREM 6. *Let $u:[a,b]\to\mathbb{R}$ be a mapping of bounded variation on $[a,b]$ and $f:[a,b]\to\mathbb{R}$ be continuous and of bounded variation on $[a,b]$. Then we have the inequality:*

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{2}\bigvee_a^b(f)\bigvee_a^{(a+b)/2}(u). \tag{2.1}$$

PROOF. Using the fact that for a continuous function $p:[a,b]\to\mathbb{R}$ and a function $\nu:[a,b]\to\mathbb{R}$ of bounded variation, one has the inequality

$$\left|\int_a^b p(t)\,d\nu(t)\right| \leqslant \sup_{t\in[a,b]}|p(t)|\bigvee_a^b(\nu). \tag{2.2}$$

As u is of bounded variation on $[a,b]$ and f is continuous, by (2.2) we have

$$\begin{aligned}|\mathcal{GS}(f;u)| &= \left|\int_a^{(a+b)/2}\left[\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right]du(x)\right| \\ &\leqslant \sup_{x\in[a,(a+b)/2]}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right|\bigvee_a^{(a+b)/2}(u).\end{aligned}$$

Since f is of bounded variation, then using the companion of Ostrowski type inequality (1.2), we may state that

$$\begin{aligned}&\sup_{x\in[a,(a+b)/2]}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right| \\ &\qquad\leqslant \sup_{x\in[a,(a+b)/2]}\left[\frac{1}{4}+\left|\frac{x-(3a+b)/4}{b-a}\right|\right]\bigvee_a^b(f)\leqslant\frac{1}{2}\bigvee_a^b(f).\end{aligned}$$

It follows that

$$\begin{aligned}|\mathcal{GS}(f;u)| &\leqslant \sup_{x\in[a,(a+b)/2]}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right|\bigvee_a^{(a+b)/2}(u) \\ &\leqslant \frac{1}{2}\bigvee_a^b(f)\bigvee_a^{(a+b)/2}(u),\end{aligned}$$

and the theorem is proved. □

REMARK 1. If $\bigvee_a^{(a+b)/2}(u) = \bigvee_{(a+b)/2}^b(u)$, then (2.1) becomes

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{4}\bigvee_a^b(f)\bigvee_a^b(u)$$

COROLLARY 1. *Let u be as in Theorem 6.*

(1) *If $f:[a,b]\to\mathbb{R}$ be an L-Lipschitzian mapping on $[a,b]$, then*

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{2}L(b-a)\bigvee_a^{(a+b)/2}(u).$$

(2) *If $f\in C^{(1)}[a,b]$, then $|\mathcal{GS}(f;u)| \leqslant \frac{1}{2}\bigvee_a^{(a+b)/2}(u)\cdot\|f'\|_{1,[a,b]}$.*

(3)*If $f:[a,b]\to\mathbb{R}$ be a monotonic mapping, then*

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{2}\bigvee_a^{(a+b)/2}(u)\cdot|f(b)-f(a)|,$$

where $\|\cdot\|_1$ is the L_1 norm, namely $\|f'\|_{1,[a,b]} := \int_a^b|f'(t)|dt$.

COROLLARY 2. *Let f be as in Theorem 6.*

(1) *If $u:[a,b]\to\mathbb{R}$ be an K-Lipschitzian mapping on $[a,b]$, then*

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{4}K(b-a)\bigvee_a^b(f).$$

(2) *If $u\in C^{(1)}[a,b]$, then $|\mathcal{GS}(f;u)| \leqslant \frac{1}{2}\bigvee_a^b(f)\cdot\|u'\|_{1,[a,(a+b)/2]}$.*

(3) *If $u:[a,b]\to\mathbb{R}$ is a monotonic mapping, then*

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{2}\bigvee_a^b(f)\cdot\Big|u\Big(\frac{a+b}{2}\Big)-u(a)\Big|,$$

where $\|\cdot\|_1$ is the L_1 norm, namely $\|u'\|_{1,[a,(a+b)/2]} := \int_a^{(a+b)/2}|u'(t)|dt$.

REMARK 2. In Corollary1, we have the following cases:

(1) If f is L-Lipschitzian mapping on $[a,b]$ and

(a) u is K-Lipschitzian mapping on $[a,b]$, then

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{4}KL(b-a)^2. \tag{2.3}$$

(b) $u\in C^{(1)}[a,b]$, then

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{2}L(b-a)\,\|u'\|_{1,[a,(a+b)/2]}. \tag{2.4}$$

(c) u is monotonic on $[a,b]$, then

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{2}L(b-a)\,\left|u\left((a+b)/2\right)-u(a)\right|. \tag{2.5}$$

(2) If $f \in C^{(1)}[a,b]$, and
(a) u is K-Lipschitzian mapping on $[a,b]$, then

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{4}K(b-a)\|f'\|_{1,[a,b]}. \tag{2.6}$$

(b) $u \in C^{(1)}[a,b]$, then we have the inequality:

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{2}\|f'\|_{1,[a,b]}\|u'\|_{1,[a,(a+b)/2]}. \tag{2.7}$$

(c) u is monotonic on $[a,b]$, then

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{2}\|f'\|_{1,[a,b]}\left|u\Big(\frac{a+b}{2}\Big) - u(a)\right|. \tag{2.8}$$

(3) If f is monotonic on $[a,b]$, and
(a) u is K-Lipschitzian mapping on $[a,b]$, then

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{4}K(b-a)|f(b)-f(a)|. \tag{2.9}$$

(b) $u \in C^{(1)}[a,b]$, then

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{2}|f(b)-f(a)|\|u'\|_{1,[a,(a+b)/2]}. \tag{2.10}$$

(c) u is monotonic on $[a,b]$, then

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{2}|f(b)-f(a)|\left|u\Big(\frac{a+b}{2}\Big) - u(a)\right|. \tag{2.11}$$

Remark 3. In Corollary 2 we have the following cases:
(1) If u is K-Lipschitzian mapping on $[a,b]$ and
(a) f is L-Lipschitzian mapping on $[a,b]$, then inequality (2.3) holds.
(b) $f \in C^{(1)}[a,b]$, then inequality (2.6) holds.
(c) f is monotonic on $[a,b]$, then inequality (2.9) holds.
(2) If $u \in C^{(1)}[a,b]$, and
(a) f is L-Lipschitzian mapping on $[a,b]$, then inequality (2.4) holds.
(b) $f \in C^{(1)}[a,b]$, then inequality (2.7) holds.
(c) f is monotonic on $[a,b]$, then inequality (2.10) holds.
(3) If u is monotonic on $[a,b]$, and
(a) f is L-Lipschitzian mapping on $[a,b]$, then inequality (2.5) holds.
(b) $f \in C^{(1)}[a,b]$, then inequality (2.8) holds.
(c) f is monotonic on $[a,b]$, then inequality (2.11) holds.

2.2. The case of r-H-Hölder type integrands.

Theorem 7. *Let $u : [a,b] \to \mathbb{R}$ be a mapping of bounded variation on $[a,b]$ and $f : [a,b] \to \mathbb{R}$ be of r-H-Hölder type mapping on $[a,b]$. Then*

$$|\mathcal{GS}(f;u)| \leqslant \frac{H}{2^r(r+1)}(b-a)^r \bigvee_{a}^{(a+b)/2}(u). \tag{2.12}$$

PROOF. As u is of bounded variation on $[a,b]$ and f is of r-H-Hölder type on $[a,b]$, by (2.2) we have

$$|\mathcal{GS}(f;u)| = \left| \int_a^{(a+b)/2} \left[\frac{f(x)+f(a+b-x)}{2} - \frac{1}{b-a}\int_a^b f(t)\,dt \right] du(x) \right|$$
$$\leqslant \sup_{x\in[a,(a+b)/2]} \left| \frac{f(x)+f(a+b-x)}{2} - \frac{1}{b-a}\int_a^b f(t)\,dt \right| \bigvee_a^{(a+b)/2}(u).$$

Using the companion of Ostrowski's type inequality (1.1), we may state that

$$\sup_{x\in[a,(a+b)/2]} \left| \frac{f(x)+f(a+b-x)}{2} - \frac{1}{b-a}\int_a^b f(t)\,dt \right|$$
$$\leqslant H \sup_{x\in[a,(a+b)/2]} \left[\frac{2^{r+1}(x-a)^{r+1}+(a+b-2x)^{r+1}}{2^r(r+1)(b-a)} \right] \leqslant H\frac{(b-a)^r}{2^r(r+1)}.$$

It follows that

$$|\mathcal{GS}(f;u)| \leqslant \sup_{x\in[a,(a+b)/2]} \left| \frac{f(x)+f\,(a+b-x)}{2} - \frac{1}{b-a}\int_a^b f(t)\,dt \right| \bigvee_a^{(a+b)/2}(u)$$
$$\leqslant H\frac{(b-a)^r}{2^r(r+1)} \bigvee_a^{(a+b)/2}(u),$$

and the theorem is proved. □

REMARK 4. Inequality (2.12) improves inequality (1.4) by the constant $\frac{1}{2^r}$, and therefore, (2.12) is better than (1.4).

COROLLARY 3. *Let $u:[a,b]\to\mathbb{R}$ be a mapping of bounded variation on $[a,b]$ and $f:[a,b]\to\mathbb{R}$ be of L-Lipschitzian type mapping on $[a,b]$. Then*

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{4}L(b-a) \bigvee_a^{(a+b)/2}(u).$$

COROLLARY 4. *Let $f:[a,b]\to\mathbb{R}$ be of r-H-Hölder type mapping on $[a,b]$.*

(1) *If u is K-Lipschitzian on $[a,b]$, then $|\mathcal{GS}(f;u)| \leqslant \frac{HK}{2^{r+1}(r+1)}(b-a)^{r+1}$.*
(2) *If $u\in C^{(1)}[a,b]$, then $|\mathcal{GS}(f;u)| \leqslant \frac{H}{2^r(r+1)}(b-a)^r \|u'\|_{1,[a,(a+b)/2]}$.*
(3) *If u is monotonic on $[a,b]$, then $|\mathcal{GS}(f;u)| \leqslant \frac{H}{2^r(r+1)}(b-a)^r \left|u\left(\frac{a+b}{2}\right)-u(a)\right|$.*

Therefore, we may deduce the following result.

COROLLARY 5. *Let $f:[a,b]\to\mathbb{R}$ be L-Lipschitzian mapping on $[a,b]$.*

(1) *If u is K-Lipschitzian on $[a,b]$, then $|\mathcal{GS}(f;u)| \leqslant \frac{1}{8}KL(b-a)^2$.*
(2) *If $u\in C^{(1)}[a,b]$, then $|\mathcal{GS}(f;u)| \leqslant \frac{1}{4}L(b-a)\|u'\|_{1,[a,(a+b)/2]}$.*
(3) *If u is monotonic on $[a,b]$, then $|\mathcal{GS}(f;u)| \leqslant \frac{1}{4}L(b-a)\left|u\left(\frac{a+b}{2}\right)-u(a)\right|$.*

2.3. The case of absolutely continuous integrands.

THEOREM 8. *Let $u : [a,b] \to \mathbb{R}$ be a mapping of bounded variation on $[a,b]$ and $f : [a,b] \to \mathbb{R}$ is absolutely continuous on $[a,b]$. Then*

$$|\mathcal{GS}(f;u)| \leqslant \frac{1}{4}(b-a)\|f'\|_{\infty,[a,b]} \bigvee_{a}^{(a+b)/2}(u).$$

PROOF. As u is of bounded variation on $[a,b]$ and f is continuous, by (2.2) we have

$$\begin{aligned}|\mathcal{GS}(f;u)| &= \left| \int_a^{(a+b)/2} \left[\frac{f(x)+f(a+b-x)}{2} - \frac{1}{b-a}\int_a^b f(t)\,dt \right] du(x) \right| \\ &\leqslant \sup_{x\in[a,(a+b)/2]} \left| \frac{f(x)+f(a+b-x)}{2} - \frac{1}{b-a}\int_a^b f(t)\,dt \right| \bigvee_{a}^{(a+b)/2}(u).\end{aligned}$$

Since f is absolutely continuous on $[a,b]$, then using the companion of Ostrowski type inequality (1.3), we may state that

$$\begin{aligned}&\sup_{x\in[a,(a+b)/2]} \left| \frac{f(x)+f(a+b-x)}{2} - \frac{1}{b-a}\int_a^b f(t)\,dt \right| \\ &\leqslant \sup_{x\in[a,(a+b)/2]} \left[\frac{1}{8} + 2\Big(\frac{x-(3a+b)/4}{b-a} \Big)^2 \right] (b-a)\|f'\|_{\infty,[a,b]} \leqslant \frac{1}{4}(b-a)\|f'\|_{\infty,[a,b]}.\end{aligned}$$

It follows that

$$\begin{aligned}|\mathcal{GS}(f;u)| &\leqslant \sup_{x\in[a,(a+b)/2]} \left| \frac{f(x)+f(a+b-x)}{2} - \frac{1}{b-a}\int_a^b f(t)\,dt \right| \bigvee_{a}^{(a+b)/2}(u) \\ &\leqslant \frac{1}{4}(b-a)\|f'\|_{\infty,[a,b]} \bigvee_{a}^{(a+b)/2}(u),\end{aligned}$$

and the theorem is proved. □

COROLLARY 6. *Let f be as in Theorem 8.*

(1) *If u is K-Lipschitzian on $[a,b]$, then $|\mathcal{GS}(f;u)| \leqslant \frac{1}{8}K(b-a)^2\|f'\|_{\infty,[a,b]}$.*
(2) *If $u \in C^{(1)}[a,b]$, then $|\mathcal{GS}(f;u)| \leqslant \frac{1}{4}(b-a)\|f'\|_{\infty,[a,b]} \cdot \|u'\|_{1,[a,(a+b)/2]}$.*
(3) *If u is monotonic on $[a,b]$, then $|\mathcal{GS}(f;u)| \leqslant \frac{1}{4}(b-a)\|f'\|_{\infty,[a,b]} \cdot \big|u\big(\frac{a+b}{2}\big) - u(a)\big|$.*

3. The case of Lipschitzian integrators

3.1. The case of bounded variation integrands.

THEOREM 9. *Let $u : [a,b] \to \mathbb{R}$ be an K-Lipschitzian on $[a,b]$ and $f : [a,b] \to \mathbb{R}$ be of bounded variation on $[a,b]$. Then $|\mathcal{GS}(f;u)| \leqslant \frac{3}{16}K(b-a)\bigvee_a^b(f)$.*

PROOF. It is well-known that for a Riemann integrable function $p:[a,b]\to\mathbb{R}$ and L-Lipschitzian function $\nu:[a,b]\to\mathbb{R}$, one has the inequality

$$\left|\int_a^b p(t)d\nu(t)\right|\leqslant L\int_a^b |p(t)|dt. \tag{3.1}$$

Therefore, as u is K-Lipschitzian on $[a,b]$, by (3.1) we have

$$\begin{aligned}|\mathcal{GS}(f;u)| &= \left|\int_a^{(a+b)/2}\left[\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right]du(x)\right|\\ &\leqslant K\int_a^{(a+b)/2}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right|dx\end{aligned}$$

Since f is of bounded variation, then using the companion of Ostrowski type inequality (1.2), we may state that

$$\begin{aligned}&\int_a^{(a+b)/2}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right|dx\\ &\qquad\leqslant \bigvee_a^b(f)\cdot\int_a^{(a+b)/2}\left[\frac{1}{4}+\left|\frac{x-(3a+b)/4}{b-a}\right|\right]dx\leqslant\frac{3}{16}(b-a)\bigvee_a^b(f).\end{aligned}$$

It follows that

$$\begin{aligned}|\mathcal{GS}(f;u)| &\leqslant K\int_a^{(a+b)/2}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right|dx\\ &\leqslant\frac{3}{16}K(b-a)\bigvee_a^b(f),\end{aligned}$$

and the theorem is proved. □

COROLLARY 7. *Let u be as in Theorem 9.*

(1) *If f is L-Lipschitzian on $[a,b]$, then $|\mathcal{GS}(f;u)|\leqslant\frac{3}{16}KL(b-a)^2$.*
(2) *If $f\in C^{(1)}[a,b]$, then $|\mathcal{GS}(f;u)|\leqslant\frac{3}{16}K(b-a)\|f'\|_{1,[a,b]}$.*
(3) *If f is monotonic on $[a,b]$, then $|\mathcal{GS}(f;u)|\leqslant\frac{3}{16}K(b-a)\cdot|f(b)-f(a)|$.*

3.2. The case of r-H-Hölder type integrands.

THEOREM 10. *Let $u:[a,b]\to\mathbb{R}$ be an K-Lipschitzian on $[a,b]$ and $f:[a,b]\to\mathbb{R}$ be of r-H-Hölder type mapping on $[a,b]$. Then*

$$|\mathcal{GS}(f;u)|\leqslant KH\frac{(b-a)^{r+1}}{2^r(r+1)(r+2)}. \tag{3.2}$$

PROOF. As u is K-Lipschitzian on $[a,b]$ and f is continuous, by (3.1) we have

$$|\mathcal{GS}(f;u)|\leqslant K\int_a^{(a+b)/2}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right|dx$$

Using the companion of Ostrowski type inequality (1.1), we may state that

$$\begin{aligned}&\int_a^{(a+b)/2}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right|dx\\&\leqslant H\int_a^{(a+b)/2}\left[\frac{2^{r+1}(x-a)^{r+1}+(a+b-2x)^{r+1}}{2^r(r+1)(b-a)}\right]dx\\&\leqslant H\frac{(b-a)^{r+1}}{2^r(r+1)(r+2)}.\end{aligned}$$

It follows that

$$\begin{aligned}|\mathcal{GS}(f;u)|&\leqslant K\int_a^{(a+b)/2}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right|dx\\&\leqslant KH\frac{(b-a)^{r+1}}{2^r(r+1)(r+2)},\end{aligned}$$

and the theorem is proved. □

REMARK 5. Inequality (3.2) improves inequality (1.5) by the constant $\frac{1}{2^r}$, and therefore (3.2) is better than (1.5).

COROLLARY 8. *Let u be as in Theorem 10, and $f:[a,b]\to\mathbb{R}$ be of L-Lipschitzian type mapping on $[a,b]$. Then $|\mathcal{GS}(f;u)|\leqslant\frac{1}{12}KL(b-a)^2$.*

3.3. The case of absolutely continuous integrands.

THEOREM 11. *Let $u:[a,b]\to\mathbb{R}$ be a mapping of bounded variation on $[a,b]$ and $f:[a,b]\to\mathbb{R}$ be absolutely continuous on $[a,b]$. Then*

$$|\mathcal{GS}(f;u)|\leqslant\frac{1}{12}K(b-a)^2\|f'\|_{\infty,[a,b]}.$$

PROOF. As u is K-Lipschitzian on $[a,b]$ and f is continuous, by (3.1) we have

$$|\mathcal{GS}(f;u)|\leqslant K\int_a^{(a+b)/2}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right|dx$$

Using the companion of Ostrowski's type inequality (1.3), we may state that

$$\begin{aligned}&\int_a^{(a+b)/2}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right|dx\\&\leqslant(b-a)\|f'\|_{\infty,[a,b]}\int_a^{(a+b)/2}\left[\frac{1}{8}+2\Big(\frac{x-(3a+b)/4}{b-a}\Big)^2\right]dx\leqslant\frac{1}{12}(b-a)^2\|f'\|_{\infty,[a,b]}.\end{aligned}$$

It follows that

$$\begin{aligned}|\mathcal{GS}(f;u)|&\leqslant K\int_a^{(a+b)/2}\left|\frac{f(x)+f(a+b-x)}{2}-\frac{1}{b-a}\int_a^b f(t)\,dt\right|dx\\&\leqslant\frac{1}{12}K(b-a)^2\|f'\|_{\infty,[a,b]},\end{aligned}$$

and the theorem is proved. □

4. A Numerical quadrature formula for the Riemann–Stieltjes integral

In this section, we use the results from the previous sections to approximate the Riemann–Stieltjes integral $\int_a^{(a+b)/2}\left[\frac{f(x)+f(a+b-x)}{2}\right]du(x)$, in terms of the Riemann integral $\int_a^b f(t)\,dt$.

THEOREM 12. *Let f, u be as in Theorem 6 and consider*

$$I_h := \{a = x_0 < x_1 < \cdots < x_{n-1} < x_n = b\},$$

be a partition of $[a,b]$. Denote $h_i = x_{i+1} - x_i$, $i = 1, 2, \ldots n-1$. Then we have

$$\int_a^{(a+b)/2} \frac{f(x)+f(a+b-x)}{2}du(x) = A_n(f,u,I_h) + R_n(f,u,I_h) \tag{4.1}$$

where,

$$A_n(f,u,I_h) = \sum_{i=0}^{n-1} \frac{u\left((x_{i+1}+x_i)/2\right) - u(x_i)}{h_i} \times \int_{x_i}^{(x_{i+1}+x_i)/2} f(t)\,dt \tag{4.2}$$

and the remainder $R_n(f,u,I_h)$ satisfies the estimation

$$|R_n(f,u,I_h)| \leqslant \frac{1}{2}\cdot \max_{i=\overline{0,n-1}}\left\{\bigvee_{x_i}^{x_{i+1}}(f)\right\}\bigvee_{a}^{(a+b)/2}(u).$$

PROOF. Applying Theorem 6 on the intervals $[x_i, x_{i+1}]$, $i = 1, 2, \cdots n-1$, we get

$$\left|\int_{x_i}^{(x_{i+1}+x_i)/2} \frac{f(x)+f(a+b-x)}{2}du(x) - \frac{u\left((x_{i+1}+x_i)/2\right)-u(x_i)}{h_i}\int_{x_i}^{(x_{i+1}+x_i)/2} f(t)\,dt\right| \leqslant \frac{1}{2}\bigvee_{x_i}^{x_{i+1}}(f)\bigvee_{x_i}^{(x_{i+1}+x_i)/2}(u).$$

Summing the above inequality over i from 0 to $n-1$ and using the generalized triangle inequality, we deduce that

$$\begin{aligned}\left|\int_a^{(a+b)/2} \frac{f(x)+f(a+b-x)}{2}du(x) - A_n(f,u,I_h)\right| &\leqslant \frac{1}{2}\sum_{i=0}^{n-1}\left[\bigvee_{x_i}^{x_{i+1}}(f)\bigvee_{x_i}^{(x_{i+1}+x_i)/2}(u)\right]\\ &= \frac{1}{2}\cdot\max_{i=\overline{0,n-1}}\left\{\bigvee_{x_i}^{x_{i+1}}(f)\right\}\sum_{i=0}^{n-1}\bigvee_{x_i}^{(x_{i+1}+x_i)/2}(u)\\ &= \frac{1}{2}\cdot\max_{i=\overline{0,n-1}}\left\{\bigvee_{x_i}^{x_{i+1}}(f)\right\}\bigvee_{a}^{(a+b)/2}(u),\end{aligned}$$

and the theorem is proved. □

THEOREM 13. *Let f, u be as in Theorem* 10 *and I_h as above. Then*

$$\int_a^{(a+b)/2} \frac{f(x)+f(a+b-x)}{2} du(x) = A_n(f,u,I_h) + R_n(f,u,I_h)$$

where, $A_n(f,u,I_h)$ is defined in (4.2) and the remainder $R_n(f,u,I_h)$ satisfies the estimation

$$|R_n(f,u,I_h)| \leqslant \frac{KH}{2^r(r+1)(r+2)} [\nu(h)]^r (b-a)$$

where, $\nu(h) = \max_{i=\overline{0,n-1}}\{h_i\}$.

PROOF. Applying Theorem 10 on the intervals $[x_i, x_{i+1}]$, $i = 1, 2, \ldots n-1$, we get

$$\left| \int_{x_i}^{(x_{i+1}+x_i)/2} \frac{f(x)+f(a+b-x)}{2} du(x) \right.$$
$$\left. - \frac{u\left((x_{i+1}+x_i)/2\right) - u(x_i)}{h_i} \int_{x_i}^{(x_{i+1}+x_i)/2} f(t)\, dt \right| \leqslant \frac{KH}{2^r(r+1)(r+2)} h_i^{r+1}.$$

Summing the above inequality over i from 0 to $n-1$ and using the generalized triangle inequality, we deduce that

$$\left| \int_a^{(a+b)/2} \frac{f(x)+f(a+b-x)}{2} du(x) - A_n(f,u,I_h) \right| \leqslant \frac{KH}{2^r(r+1)(r+2)} \sum_{i=0}^{n-1} h_i^{r+1}$$
$$\leqslant \frac{KH}{2^r(r+1)(r+2)} \left[\max_{i=\overline{0,n-1}} \{h_i\} \right]^r \sum_{i=0}^{n-1} h_i$$
$$\leqslant \frac{KH}{2^r(r+1)(r+2)} [\nu(h)]^r (b-a),$$

and the theorem is proved. □

REMARK 6. In order to approximate the Riemann–Stieltjes integral (4.1), one may state several interesting error estimations for the remainder $R_n(f,u,I_h)$ under various assumptions using the inequalities from Sections 2 and 3. We omit the details.

References

1. M. W. Alomari, *A companion of Grüss type inequality for Riemann-Stieltjes integral and applications*, avaliable as a prprin at*RGMIA Research Report Collection*, 14 (2011), Article 96, 11 pp. [http://ajmaa.org/RGMIA/papers/v14/v14a96.pdf].
2. N. S. Barnett, S. S. Dragomir and I. Gomma, *A companion for the Ostrowski and the generalised trapezoid inequalities*, Mathematical and Computer Modelling **50** (2009), 179–187.
3. N. S. Barnett, W.-S. Cheung, S. S. Dragomir, A. Sofo, *Ostrowski and trapezoid type inequalities for the Stieltjes integral with Lipschitzian integrands or integrators*, Comp. Math. Appl. **57** (2009), 195–201.
4. P. Cerone, W. S. Cheung, S. S. Dragomir, *On Ostrowski type inequalities for Stieltjes integrals with absolutely continuous integrands and integrators of bounded variation*, Comp. Math. Appl. **54** (2007), 183–191.

5. P. Cerone, S. S. Dragomir, *New bounds for the three-point rule involving the Riemann-Stieltjes integrals*, in: C. Gulati, et al. (Eds.), *Advances in Statistics Combinatorics and Related Areas*, World Science Publishing, 2002, pp. 53–62.
6. P. Cerone, S. S. Dragomir, *Approximating the Riemann–Stieltjes integral via some moments of the integrand*, Mathematical and Computer Modelling **49** (2009), 242–248.
7. S. S. Dragomir and Th.GS. Rassias (Ed.), *Ostrowski Type Inequalities and Applications in Numerical Integration*, Kluwer, Dordrecht, 2002.
8. S. S. Dragomir, *Inequalities of Grüss type for the Stieltjes integral and applications*, Kragujevac J. Math., **26** (2004), 89–112.
9. S. S. Dragomir, *On the Ostrowski inequality for Riemann–Stieltjes integral* $\int_a^b f(t)du(t)$ *where* f *is of Hölder type and* u *is of bounded variation and applications*, J. KSIAM, **5** (2001), 35–45.
10. S. S. Dragomir, *On the Ostrowski's inequality for Riemann-Stieltes integral and applications*, Korean J. Comput.& Appl. Math. **7** (2000), 611–627.
11. S. S. Dragomir, *A companion of Ostrowski's inequality for functions of bounded variation and applications*, RGMIA Preprint, Vol. 5 Supp. (2002) article No. 28. [http://ajmaa.org/RGGSIA/papers/v5e/COIFBVApp.pdf]
12. S. S. Dragomir, *Some companions of Ostrowski's inequality for absolutely continuous functions and applications*, Bull. Korean Math. Soc. **42**(2) (2005), 213–230.
13. S. S. Dragomir, *Some inequalities of midpoint and trapezoid type for the Riemann-Stieltjes integral*, Nonlinear Anal. **47**(4) (2001), 2333–2340.
14. S. S. Dragomir, *Approximating the RiemannŰStieltjes integral in terms of generalised trapezoidal rules*, Nonlinear Anal. TMA **71** (2009), e62–e72.
15. S. S. Dragomir, *Approximating the Riemann-Stieltjes integral by a trapezoidal quadrature rule with applications*, Mathematical and Computer Modelling **54** (2011), 243–260.
16. S. S. Dragomir, C. Buşe, M. V. Boldea, L. Braescu, *A generalisation of the trapezoid rule for the Riemann-Stieltjes integral and applications*, Nonlinear Anal. Forum **6**(2) (2001), 33–351.
17. S. S. Dragomir, I. Fedotov, *A Grüss type inequality for mappings of bounded variation and applications to numerical analysis*, Nonlinear Funct. Anal. Appl. **6**(3) (2001), 425–433.
18. S. S. Dragomir, I. Fedotov, *An inequality of Grüss type for Riemann-ŰStieltjes integral and applications for special means*, Tamkang J. Math. **29**(4) (1998), 287–292
19. A. Guessab, G. Schmeisser, *Sharp integral inequalities of the Hermite–Hadamard type*, J. Approx. Th. **115** (2002), 260–288.
20. Z. Liu, *Refinement of an inequality of Grüss type for Riemann-Stieltjes integral*, Soochow J. Math. **30**(4) (2004), 483–489.

Department of Mathematics
Faculty of Science, Jerash University
26150 Jerash, Jordan
mwomath@gmail.com

(Received 21 02 2012)