PUBLICATIONS DE L'INSTITUT MATHÉMATIQUE
Nouvelle série, tome 96 (110) (2014), 23–29 DOI: 10.2298/PIM1410023B

ON L^p-CONVERGENCE OF BERNSTEIN–DURRMEYER OPERATORS WITH RESPECT TO ARBITRARY MEASURE

Elena E. Berdysheva and Bing-Zheng Li

Abstract. We consider Bernstein–Durrmeyer operators with respect to arbitrary measure on the simplex in the space $\mathbb{R}^d$. We obtain estimates for rate of convergence in the corresponding weighted L^p-spaces, $1 \leqslant p < \infty$.

1. Introduction

We consider Bernstein–Durrmeyer operators with respect to arbitrary measure. These are positive linear operators defined for functions on a d-dimensional simplex. We start with notation. Let

$$\mathbb{S}^d := \{x = (x_1, \dots, x_d) \in \mathbb{R}^d : \ 0 \leqslant x_1, \dots, x_d \leqslant 1, \quad 0 \leqslant x_1 + \cdots + x_d \leqslant 1\}$$

denote the standard simplex in $\mathbb{R}^d$. We denote by $\partial\mathbb{S}^d$ the boundary of $\mathbb{S}^d$. We will also use barycentric coordinates on the simplex which we denote by the boldface symbol $\mathbf{x} = (x_0, x_1, \dots, x_d)$, $x_0 := 1 - x_1 - \cdots - x_d$. We will use standard multiindex notation such as

$$\mathbf{x}^\alpha := x_0^{\alpha_0}\, x_1^{\alpha_1} \cdots x_d^{\alpha_d} \qquad \text{and} \qquad \frac{\alpha}{n} := \Big(\frac{\alpha_0}{n}, \frac{\alpha_1}{n}, \cdots, \frac{\alpha_d}{n}\Big)$$

for $\mathbf{x} = (x_0, x_1, \dots, x_d)$, $\alpha = (\alpha_0, \alpha_1, \dots, \alpha_d) \in \mathbb{R}^{d+1}$, $n \in \mathbb{N}$. Functions defined on $\mathbb{S}^d$ are understood as functions of a point that can be given alternatively in cartesian or in barycentric coordinates.

The spaces $L^p(\mathbb{S}^d, \rho)$, $1 \leqslant p < \infty$, are defined in the standard way as spaces of (equivalence classes) of real-valued functions f for which $|f|^p$ is integrable with respect to a measure ρ with the norm

$$\|f\|_{L^p(\mathbb{S}^d,\rho)} := \Big(\int_{\mathbb{S}^d} |f(x)|^p d\rho(x) \Big)^{1/p}.$$

The space $L^\infty(\mathbb{S}^d, \rho)$ is the space of essentially bounded functions with the norm $\|f\|_{L^\infty(\mathbb{S}^d,\rho)} := \operatorname{ess\,sup}_{x\in\mathbb{S}^d} |f(x)|$. We will also consider the space $C(\mathbb{S}^d)$ of continuous bounded functions on $\mathbb{S}^d$ with the norm $\|f\|_{C(\mathbb{S}^d)} := \max_{x\in\mathbb{S}^d} |f(x)|$.

2010 *Mathematics Subject Classification:* Primary 41A36, 41A63.
Key words and phrases: positive operators, Bernstein type operators, rate of convergence.
The paper is dedicated to Giuseppe Mastroianni on the occasion of his retirement.

An important building stone of our construction are the *Bernstein basis polynomials* of degree $n \in \mathbb{N}$ on the simplex

$$B_\alpha(x) := \binom{n}{\alpha} \mathbf{x}^\alpha = \frac{n!}{\alpha_0!\alpha_1!\cdots\alpha_d!}\,(1 - x_1 - \cdots - x_d)^{\alpha_0}\, x_1^{\alpha_1}\cdots x_d^{\alpha_d},$$

with $\alpha = (\alpha_0, \alpha_1, \ldots, \alpha_d)$, where $\alpha_0, \alpha_1, \ldots, \alpha_d$ are nonnegative integers and $|\alpha| := \alpha_0 + \alpha_1 + \cdots + \alpha_d = n$. Here, and in similar expressions later, 0^0 means 1. The Bernstein basis polynomials are nonnegative on $\mathbb{S}^d$, and

$$\sum_{|\alpha|=n} B_\alpha(x) = 1.$$

The polynomials $\{B_\alpha\}_{|\alpha|=n}$ constitute a basis of the space of real algebraic polynomials in d variables of total degree at most n.

Definition 1.1. Let ρ be a nonnegative bounded Borel measure on $\mathbb{S}^d$ such that

$$\operatorname{supp}\rho \setminus \partial\mathbb{S}^d \neq \emptyset. \tag{1.1}$$

The Bernstein–Durrmeyer operator with respect to the measure ρ is defined for $f \in C(\mathbb{S}^d)$ or $f \in L^p(\mathbb{S}^d, \rho)$, $1 \leqslant p \leqslant \infty$, by

$$\mathbf{M}_{n,\rho} f := \sum_{|\alpha|=n} \frac{\int_{\mathbb{S}^d} f B_\alpha d\rho}{\int_{\mathbb{S}^d} B_\alpha d\rho}\, B_\alpha, \qquad n \in \mathbb{N}. \tag{1.2}$$

Note that ρ is regular (being a nonnegative bounded Borel measure on a metric space), and thus polynomials are dense in the spaces $L^p(\mathbb{S}^d, \rho)$, $1 \leqslant p \leqslant \infty$. Condition (1.1) guarantees that $\int_{\mathbb{S}^d} B_\alpha d\rho > 0$ for all Bernstein basis polynomials B_α.

The operator $\mathbf{M}_{n,\rho}$ is linear and positive, and it reproduces constant functions. It is a variant of the Bernstein polynomial operator $\mathbf{B}_n$ for integrable functions. The latter is defined as follows.

Definition 1.2. The Bernstein operator is defined for $f \in C(\mathbb{S}^d)$ by

$$\mathbf{B}_n f := \sum_{|\alpha|=n} f\Big(\frac{\alpha}{n}\Big)\, B_\alpha, \qquad n \in \mathbb{N}. \tag{1.3}$$

This is a linear positive operator that reproduces linear functions. The operator $\mathbf{B}_n$ was introduced by Bernstein [**7**] in the one-dimensional case in order to give a constructive proof of the Weierstrass Approximation Theorem. Many variants and generalizations of operator (1.3) were studied in hundreds of papers.

The operator $\mathbf{M}_{n,\rho}$ without weight (i.e., when ρ is the Lebesgue measure) was defined in [**12, 17**] and studied in [**8, 9**]. In the special case when ρ is the Jacobi weight, $\mathbf{M}_{n,\rho}$ was introduced in [**18, 6**]. It is very well understood; see, e.g., [**11**]. See also [**5**] for properties and further references.

Operators (1.2) in full generality were for the first time systematically studied in [**4**], to our knowledge. The motivation came from learning theory; Jetter and Zhou [**14**] used the univariate Bernstein–Durrmeyer operators of type (1.2) to obtain

bias-variance estimates for support vector machine classifiers. In [**16**], the second named author applied multivariate operators (1.2) as a tool for proving learning rates of least-square regularized regression with polynomial kernels.

In this paper, we continue our investigations on convergence of operators (1.2). In [**2**], the first named author considered uniform convergence of operators $\mathbf{M}_{n,\rho}$. She proved that

$$\lim_{n\to\infty} \|f - \mathbf{M}_{n,\rho}\, f\|_{C(\mathbb{S}^d)} = 0 \quad \text{for every} \quad f \in C(\mathbb{S}^d)$$

if and only if ρ is strictly positive on $\mathbb{S}^d$ (i.e., $\operatorname{supp}\rho = \mathbb{S}^d$). In [**3**], she considered pointwise convergence on the support of the measure. She showed that $(\mathbf{M}_{n,\rho}\, f)(x) \to f(x)$ as $n \to \infty$ at each point $x \in \operatorname{supp}\rho$ if f is bounded on $\operatorname{supp}\rho$ and continuous at x. Moreover, the convergence is uniform on any compact set in the interior of $\operatorname{supp}\rho$. Her method does not lead to estimates for rates of convergence.

The second named author studied the weighted L^p-convergence of operators (1.2). In [**16**], she proved that

$$\lim_{n\to\infty} \|f - \mathbf{M}_{n,\rho}\, f\|_{L^p(\mathbb{S}^d,\rho)} = 0$$

for every $f \in L^p(\mathbb{S}^d,\rho)$, $1 \leqslant p < \infty$. Note that no additional assumptions on ρ are required. Moreover, she obtained estimates for the rate of convergence in the spaces $L^p(\mathbb{S}^d,\rho)$, $1 \leqslant p < \infty$, in terms of the following K-functional. Let $C^1(\mathbb{S}^d)$ be the space of functions $g \in C(\mathbb{S}^d)$ with continuous partial derivatives $\partial_i g := \frac{\partial g}{\partial x_i}$, $i = 1, \dots, d$, endowed with the seminorm

$$\|\nabla g\|_{C(\mathbb{S}^d)} := \max_{i=1,\dots,d} \|\partial_i g\|_{C(\mathbb{S}^d)}.$$

The K-functional used in [**16**] is defined by

$$\mathcal{K}(f,t)_p := \inf_{g\in C^1(\mathbb{S}^d)} \bigl\{\|f-g\|_{L^p(\mathbb{S}^d,\rho)} + t\,\|\nabla g\|_{C(\mathbb{S}^d)}\bigr\}, \qquad 1 \leqslant p \leqslant \infty.$$

The following estimates were proved in [**16**]. If $f \in L^p(\mathbb{S}^d,\rho)$, $1 \leqslant p < \infty$, then

$$\|f - \mathbf{M}_{n,\rho}\, f\|_{L^p(\mathbb{S}^d,\rho)} \leqslant 2d\mathcal{K}\bigl(f, n^{-1/2p}\bigl[\rho(\mathbb{S}^d)\bigr]^{1/p}\bigr)_p, \qquad 1 \leqslant p < 2, \tag{1.4}$$

$$\|f - \mathbf{M}_{n,\rho}\, f\|_{L^p(\mathbb{S}^d,\rho)} \leqslant 2d\mathcal{K}\bigl(f, n^{-1/p}\bigl[\rho(\mathbb{S}^d)\bigr]^{1/p}\bigr)_p, \qquad 2 \leqslant p < \infty. \tag{1.5}$$

In this paper, we improve the rates given in estimates (1.4) and (1.5). Namely, by a modification of the method of [**16**], we obtain the following result.

THEOREM 1.1. *Let ρ be a nonnegative bounded Borel measure on $\mathbb{S}^d$ such that $\operatorname{supp}\rho \setminus \partial\mathbb{S}^d \neq \emptyset$, and let $f \in L^p(\mathbb{S}^d,\rho)$, $1 \leqslant p < \infty$. Then*

$$\|f - \mathbf{M}_{n,\rho}\, f\|_{L^p(\mathbb{S}^d,\rho)} \leqslant 2\mathcal{K}\bigl(f, C_p\, n^{-1/2} d\bigl[\rho(\mathbb{S}^d)\bigr]^{1/p}\bigr)_p, \qquad 1 \leqslant p < \infty,$$

where C_p is a constant that depends only on p. It holds $C_p \leqslant C_{\tilde p}$ for $p \leqslant \tilde p$. Moreover, one can take $C_p = \frac{1}{2}$ for $1 \leqslant p \leqslant 2$.

2. Proof of Theorem 1.1

Denote $\varphi_i(x) := x_i$, $i = 1, \dots, d$, and for $1 \leqslant p \leqslant \infty$

$$\Delta_{n,p} := \sum_{i=1}^{d} \left\| \mathbf{M}_{n,\rho}\big(|\varphi_i(x) - \varphi_i(\cdot)|\big) \right\|_{L^p(\mathbb{S}^d,\rho)}.$$

It is easy to see that

$$\|\mathbf{M}_{n,\rho} f - f\|_{L^p(\mathbb{S}^d,\rho)} \leqslant 2\mathcal{K}(f, \Delta_{n,p}/2)_p \tag{2.1}$$

for $f \in L^p(\mathbb{S}^d, \rho)$, $1 \leqslant p \leqslant \infty$ (see [**4**, Theorem 4.5] or [**16**, Theorem 2.1]). Thus, the key to proving estimates for the rate of convergence of the operator $\mathbf{M}_{n,\rho}$ is to study the behaviour of $\Delta_{n,p}$.

We were able to obtain estimates for $\Delta_{n,p}$ in case when $1 \leqslant p < \infty$. Theorem 1.1 is a direct consequence of the lemma given below.

LEMMA 2.1. *Let ρ be a nonnegative bounded Borel measure on $\mathbb{S}^d$ such that $\operatorname{supp}\rho \setminus \partial\mathbb{S}^d \neq \emptyset$, and let $f \in L^p(\mathbb{S}^d, \rho)$, $1 \leqslant p < \infty$. Then*

$$\|\mathbf{M}_{n,\rho}\big(|\varphi_i(x) - \varphi_i(\cdot)|\big)\|_{L^p(\mathbb{S}^d,\rho)} \leqslant c_p\, n^{-1/2} \big[\rho(\mathbb{S}^d)\big]^{1/p}, \qquad i = 1, \dots, d, \tag{2.2}$$

where c_p is a constant that depends only on p. It holds $c_p \leqslant c_{\tilde{p}}$ for $p \leqslant \tilde{p}$. Moreover, one can take $c_p = 1$ for $1 \leqslant p \leqslant 2$.

PROOF. Denote $\theta_\alpha := \int_{\mathbb{S}^d} B_\alpha \, d\rho$. Following [**16**], we write

$$\begin{aligned}
\mathbf{M}_{n,\rho}\big(|\varphi_i(x) - \varphi_i(\cdot)|\big)(x) &= \sum_{|\alpha|=n} \frac{1}{\theta_\alpha} \int_{\mathbb{S}^d} |\varphi_i(x) - \varphi_i(t)|\, B_\alpha(t)\, d\rho(t)\, B_\alpha(x) \\
&= \sum_{|\alpha|=n} \Big|\varphi_i(x) - \frac{\alpha_i}{n}\Big| B_\alpha(x) + \sum_{|\alpha|=n} \frac{1}{\theta_\alpha} \int_{\mathbb{S}^d} \Big|\frac{\alpha_i}{n} - \varphi_i(t)\Big| B_\alpha(t)\, d\rho(t)\, B_\alpha(x) \\
&= \mathbf{B}_n\big(|\varphi_i(x) - \varphi_i(\cdot)|\big)(x) + I(x),
\end{aligned}$$

where $\mathbf{B}_n$ is the Bernstein operator (1.3), and

$$I(x) := \sum_{|\alpha|=n} \frac{1}{\theta_\alpha} \int_{\mathbb{S}^d} \Big|\frac{\alpha_i}{n} - \varphi_i(t)\Big| B_\alpha(t)\, d\rho(t)\, B_\alpha(x).$$

By Cauchy–Schwarz inequality for positive operators (e.g., [**13**]), we have

$$\mathbf{B}_n\big(|\varphi_i(x) - \varphi_i(\cdot)|\big)(x) \leqslant \left(\mathbf{B}_n\left([\varphi_i(x) - \varphi_i(\cdot)]^2\right)(x)\right)^{1/2} \left(\mathbf{B}_n(1)(x)\right)^{1/2}.$$

It is well known and easy to prove that

$$\mathbf{B}_n\left([\varphi_i(x) - \varphi_i(\cdot)]^2\right)(x) = \frac{\varphi_i(x)\,(1 - \varphi_i(x))}{n} \leqslant \frac{1}{4n}, \tag{2.3}$$

and $\mathbf{B}_n(1) = 1$. Thus, $\mathbf{B}_n\big(|\varphi_i(x) - \varphi_i(\cdot)|\big)(x) \leqslant \frac{1}{2\sqrt{n}}$, and

$$\|\mathbf{B}_n\big(|\varphi_i(x) - \varphi_i(\cdot)|\big)\|_{L^p(\mathbb{S}^d,\rho)} \leqslant \frac{1}{2\sqrt{n}} \big[\rho(\mathbb{S}^d)\big]^{1/p}. \tag{2.4}$$

Next we obtain an estimate for $\|I\|_{L^p(\mathbb{S}^d,\rho)}$. Take q such that $\frac{1}{p}+\frac{1}{q}=1$. Applying the Hölder inequality two times, we obtain

$$\begin{aligned}
\|I\|^p_{L^p(\mathbb{S}^d,\rho)} &= \int_{\mathbb{S}^d}\Big\{\sum_{|\alpha|=n}\frac{1}{\theta_\alpha}\int_{\mathbb{S}^d}\Big|\frac{\alpha_i}{n}-\varphi_i(t)\Big|B_\alpha(t)\,d\rho(t)\,B_\alpha^{\frac{1}{p}+\frac{1}{q}}(x)\Big\}^p d\rho(x)\\
&\leqslant \int_{\mathbb{S}^d}\sum_{|\alpha|=n}\frac{1}{\theta_\alpha^p}\Big(\int_{\mathbb{S}^d}\Big|\frac{\alpha_i}{n}-\varphi_i(t)\Big|B_\alpha(t)\,d\rho(t)\Big)^p B_\alpha(x)\,d\rho(x)\\
&= \sum_{|\alpha|=n}\frac{1}{\theta_\alpha^{p-1}}\Big(\int_{\mathbb{S}^d}\Big|\frac{\alpha_i}{n}-\varphi_i(t)\Big|B_\alpha^{\frac{1}{p}+\frac{1}{q}}(t)\,d\rho(t)\Big)^p\\
&\leqslant \sum_{|\alpha|=n}\frac{1}{\theta_\alpha^{p-1}}\int_{\mathbb{S}^d}\Big|\frac{\alpha_i}{n}-\varphi_i(t)\Big|^p B_\alpha(t)\,d\rho(t)\,\theta_\alpha^{p/q}\\
&= \int_{\mathbb{S}^d}\sum_{|\alpha|=n}\Big|\frac{\alpha_i}{n}-\varphi_i(t)\Big|^p B_\alpha(t)\,d\rho(t)\\
&= \big\|\mathbf{B}_n\big(|\varphi_i(x)-\varphi_i(\cdot)|^p\big)\big\|_{L^1(\mathbb{S}^d,\rho)}.
\end{aligned}$$

First suppose that $p\geqslant 1$ is an even integer. In this case, the expression in the last line of the previous formula is the $L^1(\mathbb{S}^d,\rho)$-norm of a moment of the Bernstein operator (1.3), namely, of $\mathbf{B}_n\big([\varphi_i(x)-\varphi_i(\cdot)]^p\big)(x)$. First we note that the value of this moment is independent of the dimension d. To see this, consider without loss of generality $i=1$. Then

$$\begin{aligned}
\mathbf{B}_n\big([\varphi_1(x)-\varphi_1(\cdot)]^p\big)(x) &= \sum_{|\alpha|=n}\Big(x_1-\frac{\alpha_1}{n}\Big)^p B_\alpha(x)\\
&= \sum_{|\alpha|=n}\Big(x_1-\frac{\alpha_1}{n}\Big)^p\binom{n}{\alpha_1}x_1^{\alpha_1}(1-x_1)^{n-\alpha_1}\\
&\qquad\times\frac{(n-\alpha_1)!}{\alpha_2!\cdots\alpha_0!}\Big(\frac{x_2}{1-x_1}\Big)^{\alpha_2}\cdots\Big(\frac{x_0}{1-x_1}\Big)^{\alpha_0}\\
&= \sum_{\alpha_1=0}^{n}\Big(x_1-\frac{\alpha_1}{n}\Big)^p\binom{n}{\alpha_1}x_1^{\alpha_1}(1-x_1)^{n-\alpha_1}\\
&\qquad\times\sum_{|(\alpha_2,\dots,\alpha_d,\alpha_0)|=n-\alpha_1}B_{(\alpha_2,\dots,\alpha_d,\alpha_0)}\Big(\frac{x_2}{1-x_1},\cdots,\frac{x_d}{1-x_1}\Big)\\
&= \sum_{\alpha_1=0}^{n}\Big(x_1-\frac{\alpha_1}{n}\Big)^p\binom{n}{\alpha_1}x_1^{\alpha_1}(1-x_1)^{n-\alpha_1}
\end{aligned}$$

which is the p-th moment of the one-dimensional Bernstein operator. Estimates for these moments are well known an can be found, e.g., in [**10**, Chapter 10, §1]. It follows from Corollary to Theorem 1.1 of this chapter that there is a constant A_p depending only on p such that

$$\sum_{\alpha_1=0}^{n}\Big(x_1-\frac{\alpha_1}{n}\Big)^p\binom{n}{\alpha_1}x_1^{\alpha_1}(1-x_1)^{n-\alpha_1}\leqslant A_p\,n^{-p/2}.$$

Consequently,

$$\|I\|_{L^p(\mathbb{S}^d,\rho)} \leqslant \left\{\|\mathbf{B}_n([\varphi_i(x)-\varphi_i(\cdot)]^p)\|_{L^1(\mathbb{S}^d,\rho)}\right\}^{1/p} \leqslant n^{-1/2}A_p^{1/p}\ \left[\rho(\mathbb{S}^d)\right]^{1/p}. \tag{2.5}$$

For an arbitrary $p \geqslant 1$, take $\tilde{p}$ to be the smallest even integer with $p \leqslant \tilde{p}$. The $L^p(\mathbb{S}^d,\rho)$-and $L^{\tilde{p}}(\mathbb{S}^d,\rho)$-norms are connected by the inequality

$$\|\cdot\|_{L^p(\mathbb{S}^d,\rho)} \leqslant \|\cdot\|_{L^{\tilde{p}}(\mathbb{S}^d,\rho)}\ \left[\rho(\mathbb{S}^d)\right]^{\frac{1}{p}-\frac{1}{\tilde{p}}} \tag{2.6}$$

(e.g., [**15**, Chapter IV, §3, Theorem 6]). This estimate and (2.5) yield

$$\|I\|_{L^p(\mathbb{S}^d,\rho)} \leqslant n^{-1/2}A_{\tilde{p}}^{1/\tilde{p}}\left[\rho(\mathbb{S}^d)\right]^{1/p}.$$

Combining this with (2.4), we obtain

$$\begin{aligned}\|\mathbf{M}_{n,\rho}\big(|\varphi_i(x)-\varphi_i(\cdot)|\big)\|_{L^p(\mathbb{S}^d,\rho)} &\leqslant \|\mathbf{B}_n\big(|\varphi_i(x)-\varphi_i(\cdot)|\big)\|_{L^p(\mathbb{S}^d,\rho)} + \|I\|_{L^p(\mathbb{S}^d,\rho)}\\ &\leqslant n^{-1/2}c_p\left[\rho(\mathbb{S}^d)\right]^{1/p}\end{aligned}$$

with $c_p = \frac{1}{2} + A_{\tilde{p}}^{1/\tilde{p}}$, which is (2.2).

It follows from (2.6) that for all $p \leqslant \tilde{p}$ it holds

$$\|\mathbf{M}_{n,\rho}\big(|\varphi_i(x)-\varphi_i(\cdot)|\big)\|_{L^p(\mathbb{S}^d,\rho)} \leqslant n^{-1/2}c_{\tilde{p}}\left[\rho(\mathbb{S}^d)\right]^{1/p}.$$

Thus, $c_p \leqslant c_{\tilde{p}}$ for $p \leqslant \tilde{p}$.

Finally, consider $p = 2$. In this case $A_2 = \frac{1}{4}$ (see (2.3)). Thus, $c_2 = 1$, and we also can take $c_p = 1$ for $1 \leqslant p \leqslant 2$. □

Remark 2.1. Representations of general moments of the multivariate Bernstein operator (1.3) of the form $\mathbf{B}_n\big(\prod_{i=1}^d(\varphi_i(x)-\varphi_i(\cdot))^{p_i}\big)$ with nonnegative integers p_i, $i = 1,\dots,d$, in terms of Stirling numbers are given by Abel and Ivan [**1**]. They also estimated the order of these moments.

Remark 2.2. Alternatively, we could use in Theorem 1.1 the estimate

$$\|\mathbf{M}_{n,\rho}f - f\|_{L^p(\mathbb{S}^d,\rho)} \leqslant \max\{2,d\}\,\mathcal{K}\left(f,\Delta_{n,p}\right)_p$$

instead of (2.1). This inequality leads to the estimate

$$\|\mathbf{M}_{n,\rho}f - f\|_{L^p(\mathbb{S}^d,\rho)} \leqslant \max\{2,d\}\,\mathcal{K}\big(f, n^{-1/2}c_p\left[\rho(\mathbb{S}^d)\right]^{1/p}\big)_p, \qquad 1 \leqslant p < \infty,$$

with a constant c_p like in Lemma 2.1.

Remark 2.3. Our method does not lead to estimates for the rates of convergence of the operator $\mathbf{M}_{n,\rho}$ in the space $L^\infty(\mathbb{S}^d,\rho)$, or to pointwise estimates. These are important and interesting open questions.

Acknowledgement. The research was initiated while the first named author was visiting Mathematics Research Institute, The Ohio State University, Columbus, OH, USA. She acknowledges the hospitality of this organization and financial support provided by them.

References

1. U. Abel, M. Ivan, *Asymptotic expansion of the multivariate Bernstein polynomials on a simplex*, Approx. Theory Appl. **16** (2000), 85–93.
2. E. E. Berdysheva, *Uniform convergence of Bernstein-Durrmeyer operators with respect to arbitrary measure*, J. Math. Anal. Appl. **394** (2012), 324–336.
3. ———, *Bernstein-Durrmeyer operators with respect to arbitrary measure, II: pointwise convergence*, J. Math. Anal. Appl. **418** (2014), 734–752.
4. E. E. Berdysheva, K. Jetter, *Multivariate Bernstein-Durrmeyer operators with arbitrary weight functions*, J. Approx. Theory **162** (2010), 576–598.
5. E. Berdysheva, K. Jetter, J. Stöckler, *Durrmeyer operators and their natural quasi-interpolants*, in: K. Jetter at al. (ed.), *Topics in Multivariate Approximation and Interpolation*, Elsevier, Amsterdam, 2006, 1–21.
6. H. Berens, Y. Xu, *On Bernstein-Durrmeyer polynomials with Jacobi weights*, in: C. K. Chui (ed.),*Approximation Theory and Functional Analysis*, Academic Press, Boston, 1991, 25–46.
7. S. Bernstein, *Démonstration du théorème de Weierstrass, fondée sur le calcul des probabilités*, Commun. Soc. Math. Kharkow (2) **13** (1912–13), 1–2.
8. M.-M. Derriennic, *Sur l'approximation de fonctions intégrables sur* $[0,1]$ *par des polynômes de Bernstein modifies*, J. Approx. Theory **31** (1981), 325–343.
9. ———, *On multivariate approximation by Bernstein-type polynomials*, J. Approx. Theory **45** (1985), 155–166.
10. R. DeVore, G. G. Lorentz, *Constructive Approximation*, Springer-Verlag, Berlin, Heidelberg, New York, 1993.
11. Z. Ditzian, *Multidimensional Jacobi-type Bernstein-Durrmeyer operators*, Acta Sci. Math. (Szeged) **60** (1995), 225–243.
12. J.-L. Durrmeyer, *Une formule d'inversion de la transformée de Laplace: Applicationsà la théorie des moments*, Thèse de 3e cycle, Faculté des Sciences de l'Université de Paris, 1967.
13. H. Gonska, P. Piţul, I. Raşa, *On Peano's form of the Taylor remainder, Voronovskaja's theorem and the commutator of positive linear operators*, in: O. Agratini et al. (eds.), *Proc. Internat. Conf. Numer. Anal. Approx. Theory, Cluj-Napoca, Romania, July 4–8, 2006*, Casa Cărtii de Ştiinta, Cluj-Napoca, 2006, 55–80.
14. K. Jetter, D.-X. Zhou, *Approximation with polynomial kernels and SVM classifiers*, Adv. Comput. Math. **25** (2006), 323–344.
15. L. V. Kantorovich, G. P. Akilov, *Functional Analysis*, 2nd ed., Oxford: Pergamon Press, 1982.
16. B.-Z. Li, *Approximation by multivariate Bernstein-Durrmeyer operators and learning rates of least-square regularized regression with multivariate polynomial kernels*, J. Approx. Theory **173** (2013), 33–55.
17. A. Lupaş, *Die Folge der Betaoperatoren*, Dissertation, Universität Stuttgart, 1972.
18. R. Păltănea, *Sur un opérateur polynomial defini sur l'ensemble des fonctions intégrables*, "Babeş-Bolyai" Univ., Fac. Math., Res. Semin. **2** (1983), 101–106.

German University of Technology in Oman
Muscat, Oman
elena.berdysheva@gutech.edu.om

Zhejiang University
Hangzhou, China
libingzheng@zju.edu.cn