Fourteenth International Conference on Geometry, Integrability and Quantization June 8–13, 2012, Varna, Bulgaria Ivaïlo M. Mladenov, Andrei Ludu and Akira Yoshioka, Editors **Avangard Prima**, Sofia 2013, pp 253–267 doi: 10.7546/giq-14-2013-253-267



## EXPANSIONS OVER ADJOINT SOLUTIONS FOR THE CAUDREY-BEALS-COIFMAN SYSTEM WITH $\mathbb{Z}_p$ REDUCTIONS OF MIKHAILOV TYPE

## ALEXANDAR B. YANOVSKI

Department of Mathematics and Applied Mathematics, University of Cape Town Cape Town, Rondebosch 7700, South Africa

Abstract. We consider the Caudrey-Beals-Coifman linear problem and the theory of the Recursion Operators (Generating Operators) related to it in the presence of  $\mathbb{Z}_p$  reduction of Mikhailov type

## 1. Introduction

## 1.1. The Generalized Zakharov-Shabat and Caudrey-Beals-Coifman Systems

As it is well known **nonlinear evolution equations** (NLEEs) of soliton type are equations (systems) that can be written into the form [L, A] = 0 (Lax represenstation) where L, A are linear operators on  $\partial_x, \partial_t$  depending also on some functions  $q_\alpha(x,t), 1 \le \alpha \le s$  (called **potentials**) and the **spectral parameter**  $\lambda$ . The corresponding system is of a course system of partial differential equations on  $q_\alpha(x,t)$ . Usually the equation is a part of a hierarchy of NLEEs related to  $L\psi = 0$  (auxiliary linear problem) which consists of the equations that can be obtained by changing A and fixing L, [7, 15]. The soliton equations possess many interesting properties but for our purposes we shall mention only that they can be solved explicitly through various schemes, most of which share the property that the Lax representation permits to pass from the original evolution to the evolution of some spectral data related to the problem  $L\psi = 0$ . The **Caudrey-Beals-Coifman** (CBC) system, called the **Generalized Zakharov-Shabat** (GZS) system in the case when the element J is real, is one of the best known auxiliary linear problems

$$L\psi = (i\partial_x + q(x) - \lambda J)\psi = 0.$$
(1)